



RESEARCH ARTICLE

# Moringa flower incorporated instant dip soups - market sustainable approach

GG Kavitha Shree<sup>1\*</sup>, M Chitra<sup>2\*</sup>, M Visalakshi<sup>3</sup> & SR Padma<sup>4</sup>

<sup>1</sup>Agricultural Engineering College and Research Institute, Tamil Nadu Agricultural University, Coimbatore 641 003, India

<sup>2</sup>Dr. M.S.S. Agricultural College and Research Institute, Tamil Nadu Agricultural University, Thanjavur 614 902, Tamil Nadu, India

<sup>3</sup>Horticultural College and Research Institute, Tamil Nadu Agricultural University, Coimbatore 03, Tamil Nadu, India

<sup>4</sup>Tamil Nadu Agricultural University, Coimbatore 03, Tamil Nadu, India

\*Email: [kavikarthikfsn@gmail.com](mailto:kavikarthikfsn@gmail.com)



## ARTICLE HISTORY

Received: 30 June 2024

Accepted: 03 July 2024

Available online

Version 1.0 : 26 December 2024



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Shree GGK, Chitra M, Visalakshi M, Padma SR. Moringa flower incorporated instant dip soups - market sustainable approach. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.4213>

## Abstract

India is a long way towards alleviating micronutrient deficiencies and developing functional foods for various chronic degenerative disorders. In developing countries, women's diets have inadequate micronutrients and are mostly non-heme foods. Intake of animal products, fruits, vegetables and fortified foods are recommended for micronutrient deficiencies. In the standardization of food products, food scientists and entrepreneurs face challenges in nutrition-suitable and market-value foods. However, the benefits of food-based strategies for preventing malnutrition are manifold nowadays and are focused on easily accessible, inexpensive green vegetables for micronutrient deficiencies. *Moringa oleifera* is a repository of nutrients for the creation of value-added food products. The leaves, pods, flowers and seeds of the plant are consumable and rich in various nutrients. *Moringa oleifera* is a miracle tree that has a very impressive range of medicinal and nutritional properties. In recent years phyto chemicals in *Moringa* flower have been studied extensively. In alleviating micronutrient deficiencies, it is the need of the hour to promote the *Moringa* flower as an incorporated diet. The objective of this study is to assess the potential use of *Moringa* flower in instant dip soups to enhance nutritional quality and storage stability. In this study, optimization process parameters via RSM Box-Behnken design were done and based on the same moringa dip drink was finalised. The findings of the current investigation demonstrated that the developed Moringa flower instant dip drink retains sufficient nutrients and can be utilized as a dietary supplement for various micronutrient deficits. The development of the Instant moringa dip drink has a market-sustainable approach to the generation of additional income for farmers, self-employment and export value in the food industry.

## Keywords

food industry; food product; instant dip drink; micronutrient deficiencies; moringa flower

## Introduction

Dietary methods emphasizing green vegetables are crucial for eliminating micronutrient deficiencies and present a challenge for food scientists in standardizing healthy and organoleptically acceptable food products available in the market. The deficiency of vitamins and minerals in food causes night blindness, iron-deficiency anemia and bone-related diseases. Women are at risk of micronutrient deficiencies due to low intakes of micronutrients in

recommended values (1). This leads to deficiencies in vitamin A, iron and folic acid and has consequences in pregnancy outcomes and malnutrition in children and aged women (2). *Moringa oleifera*, native to India, is commonly known as the 'drumstick tree' and is widely cultivated across the world. *M. oleifera* tree differs in nutrient composition at different locations (3) and there is a slight difference in nutritional components of the tree grown in India to that grown in Nigeria. The nutritional differences in the leaves from two ecological locations, semi-deciduous and Savannah regions, were studied by Asante et al. (4).

Every part of *M. oleifera* is a storehouse of important phytoconstituents, with high nutritive value, suitable for nutritional and commercial purposes. *Moringa* has numerous elements vital for growth and development and it possesses a higher iron content than spinach (5). The leaves are rich in minerals calcium, potassium, zinc, magnesium, iron and copper (6), vitamins and other essential phytochemicals. *M. oleifera* leaves contain around 25.5-31.03 mg of zinc per kilogram, which meets the daily dietary need for zinc (7) and serves as a potential antioxidant, anticancer, anti-inflammatory, antidiabetic and antibacterial agent. *M.oleifera* seed, a natural coagulant, is extensively used in water treatment and flowers are an exceptionally good source of vitamins, minerals and amino acids such as methionine and cysteine (8). Bioactive substances including tannins, saponins, anthraquinones, alkaloids, terpenoids, flavonoids and phytosterols such as stigmasterol, sitosterol and campesterol, together with reducing sugars, are present in *M. oleifera* and are utilized to address malnutrition in children (9). Anti-cancerous agents like glucosinolates, isothiocyanates, glycoside compounds and glycerol-1-9-octadecanoate are also found in *Moringa* (10). *Moringa* leaves also have a low calorific value that can be used in the diet of obese people. The pods are fibrous, which is valuable for treating digestive problems and thwarting colon cancer (11). Research showed that immature pods contain around 46.78% fiber and around 20.66% protein content and are valuable in treating digestive problems and inhibit colon cancer (10). Pods have 30% of amino acid content, the leaves have 44% and flowers have 31%. The immature pods and flowers showed similar amounts of palmitic, linolenic, linoleic and oleic acids (12). Instant drink mixes are in the schemes presented by the Integrated Child Development Service of Tamilnadu (13) for school children of Tamilnadu.

The food industry uses the versatile portions of the *Moringa* tree to create a range of products with additional value. Significant findings were obtained in the physicochemical and functional content and storage studies of muffins containing *Moringa oleifera* powder (MOLP) at a concentration of 12% (9). Aloe vera (4%) enriched with *Moringa* (1%) resulted in a drink rich in color, flavor, taste overall acceptability (14). The addition of *M. oleifera* (MOLE) extract or oil is having a significant impact in enhancing microbial stability and improving the shelf-life of sour cream (8). Incorporating 10% *Moringa* leaf debris into cookies fulfills daily nutritional fiber requirements (15). Fortification of dry leaves of *M. oleifera* (2%) has a good effect in biological value (BV), true protein digestibility (TD) and net protein utilization (NPU) in Labneh cheese (16). *Moringa* leaf powder (10%) in *Moringa* idly resulted in sensory evaluation scores in the 8.0

range and higher nutritive value for supplementation (6). The addition of *M. oleifera* powder (15%) in durum wheat semolina enhanced nutritive and functional value without affecting the sensory acceptability of the pasta (17). The value addition of Lassi is increased by blending *Moringa* pod powder (1.63%), which has a high content of Vitamin C and fiber. As a natural functional additive, moringa flower extract in cooked chicken nuggets may be useful for improving quality and lowering lipid oxidation (18). One per cent *Moringa* seed flour addition enhances the nutritional and sensory qualities of mayonnaise (18,19). The scientific effort of this research provides insights into the use of *Moringa* as a cure for diabetes and cancer and the fortification of moringa in commercial products.

*Moringa* flower is a good source of nutraceuticals and has lots of bioactive compounds in flowers, seeds, leaves, oil and pods (20,22,28). The Flower of *Moringa* consists of essential phytocompounds, vitamins, omega oils, minerals and antioxidants. The dry *Moringa* flower contains calcium levels 17 times higher to those found in milk, vitamin C content 7 times greater than that of oranges, potassium levels 15 times higher than those in bananas, iron content 25 times higher than that in spinach, vitamin A levels 10 times greater than those in carrots, and protein content 9 times that of yogurt (7). The *Moringa* flower possesses medicinal properties that benefit the cardiovascular system, regulate blood glucose levels, neutralize free radicals, offer substantial anti-inflammatory support, elevate iron levels in the blood, and bolster the immune system. It is also effective in improving eyesight, mental abilities, bone strength, malnutrition and depression (21,22). *Moringa* flower provides a nutritional supplement around the world as a vegetable all over the world (23,28). Dried *Moringa* flowers are stored for many months under ambient conditions, and there is a minimum loss in nutritional value. As a food supplement, flower powder can be added to any dish to enrich the nutritional content of the products. The processing technology of *Moringa* flower powder is simple and can be easily stored. It is appropriate for many South Indian dishes, such as roti, chapati, biscuits, soup, rice and juice mixes and is significantly demanded in both national and international markets (24).

*The Moringa flower belongs to the Moringaceae family and is well known for its nutritive value, and the twigs, stems and seeds contain high protein content (25,26). Lots of minerals essential for growth and development, i.e. calcium, zinc and iron, are present in Moringa. While leaves of Moringa provide 1000 mg (milk has 300-400 mg), and Moringa powder can offer more than 4000 mg (27). However, flowers have a large amount of calcium when compared to leaves. Moringa leaf powder has 28 mg more iron than spinach and can be used as a substitute for iron tablets for anemia treatment (28). M. oleifera leaves showed around 25.5 -31.03 mg of zinc/kg, essential for synthesizing DNA and RNA (29). These studies would help the farmers to export Moringa value-added foods and youngsters to act as entrepreneurs. It is worth mentioning that Moringa-based soup dips are not available in the Indian market. Hence, an instant dip drink based on Moringa flower is developed in this study.*

## Materials and Methods

Fresh *Moringa oleifera* flowers were obtained from an organic moringa farm in the Tiruppur district, Tamilnadu. For initial range-finding trials, Moringa flowers were added in amounts ranging from 6 g to 25 g of fresh weight (FW) for every 30 g raw weight (RW) of the mix (i.e. one serving of each product) and three instant dip drink mixes were standardized (Fig.3b). Final evaluations were performed on product containing 20 g FW of blanched Moringa flower (MF) for every 30 g RW of instant drink mix in each. After harvest, young moringa flowers (T<sub>1</sub>) and, mature moringa flowers (T<sub>2</sub>), and flowers along with calyx and stalk (T<sub>3</sub>) were stripped off, washed, blanched and dried in a solar drier. The dried flowers are made into fine powder, which can be stored in air-tight containers.

An index of overall acceptability for sensory evaluation provides appearance, flavor, taste, and texture after taste and it is based on 9-point hedonic scale. The modified products were evaluated to ensure their acceptability by composite scoring for the sensory qualities. All trials *Moringa* flowers (T1 -Immature flowers, T2- Matured flowers, T3 - Flowers along with calyx and stalk and T4- Flowers along with calyx and stalk grinded) were prepared fresh and presented to a panel of 15 trained judges. The characteristics of each formula, i.e., appearance, color, flavor, taste, texture and overall acceptability, were rated separately on a scale of 1 to 5. Scores were defined as 1 - dislike, extremely bad; 2 - like only slightly tolerable; 3 - like, good; 4 - like very much, very good; 5 - like, extremely excellent. Numerical averages were then calculated for a composite test score for the range-finding trials. Each of the 3 products developed was served to all sensory analysts who were the judges and then the recipes were modified and improved. Based upon their suggestions, final tests were conducted by incorporating 20 g of fresh Moringa flower into each serving since this addition level received the highest initial ratings from the judges.

## Results and Discussion

The sensory characteristics of the Moringa flower dip drink from the range-finding trials indicated that the most acceptable formulation contained 20 mg of freshly blanched Moringa flowers or flower powder per serving of the soup mix. Evaluation of the nutritional and sensory qualities is done in the *M. oleifera* flower dips. The overall composite score for T2 was highest (Table 1) and this score was followed by T1 powder at 3.3±0.2 and T3 at 3.1 ± 0.5. Scores for the three tested dip drink products from each of the individual attributes ranged from 3.2 to 3.4 (Table 2). Most of the panel preferred composite scores to evaluate the food products and dehydrated MF powder incorporated has been evaluated using composite scores (Table 3).

The nutritive value of the developed product was assessed for nutrients and sensory attributes such as taste and flavor. The nutritive values of trials supplemented with MF were calculated using RDA and the three products provided the same amounts of nutrients. Results indicated that despite nutrient losses, there is enough  $\beta$ -carotene retained in instant dip drinks to help eradicate several micronutrient deficiencies (30). When compared to RDA, the nutritional values for non-pregnant, pregnant and lactating women, it was recorded that

the products studied had vitamin C-170-370% of  $\frac{1}{3}$  RDA, iron-12-15 % of  $\frac{1}{3}$  RDA and  $\beta$ -carotene-316-510% of  $\frac{1}{3}$  RDA. Twenty grams of Moringa flower provided 3810  $\mu$ g of  $\beta$ -carotene, which is equal to 663  $\mu$ g retinol equivalents (RE) (1 RE = 6  $\mu$ g  $\beta$ -carotene) (Table 4). The three trials of fresh Moringa flower preparation had a  $\beta$ -carotene content from 3966  $\mu$ g to 4200  $\mu$ g per serving, which is equivalent to 661-669  $\mu$ g RE (Fig.2). There may be 50% losses of the  $\beta$ -carotene content during cooking. According to the most recent studies on processing losses for  $\beta$ -carotene, even if the final recipes have 1980  $\mu$ g or 330-333 RE, it meets 82.5% of RDA to adult women. To combat vitamin A deficiency, food-based approaches are generally considered for improving bioavailability (31). These results have an important implication that T1 (MF) had good sensory attributes compared to others for making the instant dip drink food product (Fig. 1, 2, 3a, 3b, 3c). Nutrients in food should prevent disease and disorders at the molecular level of the body. The deficiency of iron in the diet is considered the most important factor in nutritional anemia.

Several studies showed that *Moringa*-based food is used to treat malnutrition problems. However, due to its slightly bitter taste, children refuse to take in moringa (32). Similarly, *Moringa* incorporation in food increases the delicacy's nutrient value and has shown the potential for developing protein and minerals-rich chocolate (33). Moringa fortifications are possible to ensure intake of adequate nutrients in children and the *Moringa* flowers are used to make tea with low cholesterol properties. Moringa flowers are said to taste like mushrooms when fried (34) and the moringa flowers are great sources of nectar and are used by beekeepers. The root bark of Moringa has medicinal values and is used for dyspepsia, eye diseases and heart complaints (35). The taproot of *Moringa* is used as a spice, and the gum from the tree can be used in calico printing. The gum and roots also have antibacterial, antifungal and anti-inflammatory properties (36). Zeatin derived from leaves is an exceptional growth promoter that can enhance crop output by 25%-30% (12). Incorporation and fortification of moringa can significantly tackle nutrient deficiencies and malnutrition. Owusu et al. (37) also used *Moringa* as a fortificant and produced cream and butter crackers with moringa and *Ipomoea batatas* as fortificants, with the hope of adding additional nutrients to snacks. The sensory evaluation proved the cream crackers to be widely accepted. *M. oleifera* leaves can be incorporated into the diet of hens and layers, thereby providing an excellent protein source, substituting other expensive ingredients such as soybean meal and ground nut cake (38,39). Cereal gruels have also been fortified with moringa leaves to improve the protein content and energy. The cereal gruel with 65% popcorn and 35% moringa leaves is blanched and fermented and showed higher protein content.

Antioxidants in Moringa can reduce the reactive oxygen species, thereby protecting the brain (40,41). As a promoter of spatial memory, it is used to treat dementia (39) and is also useful to increase protein content and decrease urea and creatinine in blood to prevent renal dysfunction. Moringa decreased acidity in gastric ulcers that can be used as an antiulcer agent (42) *M. oleifera* has been proven as a good antimicrobial agent (43) to act against bacteria like *Bacillus*

**Table 1:** Experimental runs of Box-Behnken design

	F1	F2	F3	R1
RUN	A	B	C	A1
	°Celcius	hours	%	µg/g
1	70	22	0.3	42
2	70	22	0.2	42
3	70	34	0.6	49
4	80	12	0.3	24
5	60	22	0.4	28
6	80	22	0.4	30
7	60	10	0.4	27
8	70	36	0.2	28
9	80	24	0.2	25
10	70	24	0.4	45
11	70	24	0.4	44
12	70	24	0.4	46
13	60	24	0.2	22
14	80	36	0.4	31
15	60	36	0.4	29
16	70	12	0.2	31
17	70	12	0.6	30

F1-Factor 1, F2-Factor 2, F3-Factor 3, R1-Response -1



A: Drying Temperature, B-Duration of drying, C-Citric acid, A1-Ascorbic acid

**Table 2:** ANOVA for quadratic model

	SUM OF SQUARES	DF	MEAN SQUARE	F	P	
<b>MODEL</b>	1228.11	9	136.46	50.41	< 0.0001	significant
<b>A-DRYING TEMPERATURE</b>	3.13	1	3.13	1.15	0.3183	
<b>B-DURATION OF DRYING</b>	66.13	1	66.13	24.43	0.0017	
<b>C-CITRIC ACID</b>	144.50	1	144.50	53.38	0.0002	
<b>AB</b>	2.25	1	2.25	0.8311	0.3923	
<b>AC</b>	1.0000	1	1.0000	0.3694	0.5625	
<b>BC</b>	121.00	1	121.00	44.70	0.0003	
<b>A<sup>2</sup></b>	598.76	1	598.76	221.18	< 0.0001	
<b>B<sup>2</sup></b>	82.44	1	82.44	30.45	0.0009	
<b>C<sup>2</sup></b>	135.60	1	135.60	50.09	0.0002	
<b>R</b>	18.95	7	2.71			
<b>LF</b>	15.75	3	5.25	6.56	0.0504	not significant
<b>PE</b>	3.20	4	0.8000			
<b>COR TOTAL</b>	1247.06	16				

Response =  $44.6 + 0.625 * A + 2.875 * B + 4.25 * C + 0.75 * AB + -0.5 * AC + 5.5 * BC + -11.925 * A^2 + -4.425 * B^2 + -5.675 * C^2$ 

R-residual, L-lack of fit, PE-pure error, L-lack of fit

**Fig. 2.** Predicted vs. actual result.**Fig. 1.** Normal plot.**Table 3:** optimised process conditions

F	NAME	LEVEL
A	DT	65
B	D	28
C	C	0.55


For 43.58µg/g of ascorbic acid retention, the optimized process parameters are 0.55% of citric acid, 28 hours of drying at 65° C.

F- factor, DT-Drying Temperature; D-Duration of drying; C-Citric acid

**Table 4:** The nutritive values in developed products

Nutrients	Value
Energy	150 Kcal
Protein	6-10 g
Calcium	114-141 mg
Iron	1.5-1.9 mg
β-carotene	3888-3911 µg
Ascorbic acid	42-48 mg





**Fig. 3a.** Comparison of the impact of drying temperature and duration of drying on ascorbic acid retention.

*subtilis*, *Staphylococcus aureus* and *Vibrio cholera*. Compounds such as pterygosperrin, moringine and benzyl isothiocyanate present in *M. oleifera* are responsible for the anti-bacterial activity. Beta-carotene-rich Moringa flower can be an important source of vitamin A that is used to release bound iron stores for reducing deficiency of anemia and vitamin A diseases. Hence, diet supplementation with *M. oleifera* increases iron and vitamin A content in blood. In the present study, it is also worth mentioning that the developed dip drink is fully organic, does not have chemicals, and requires no extra effort to make the usual recipes. It reveals that the inclusion of the Moringa flower in dips leads to the development of natural nutritional recipes in modern foods. The inclusion of Moringa flower foods would help the development of good entrepreneurship and provide an additional source of income.

## Conclusion

In this study, instant dip soups from *M. oleifera* flowers are standardized, and the formulated dip is fixed for further upscaling. Out of three dips standardized, 20mg/100ml of dip soups have more nutrients compared to other dips and it reveals that the processing technology of Moringa flower is simple and it has great demand in national and international markets. Moringa flower based instant dips may be combined with other herbals to enhance nutrients and to produce mixed instant dips for foreign exports.

## Acknowledgements

All authors had a role in the technical help of project work, writing the article, and assistance with revision. The authors acknowledge Tamil Nadu Agricultural University, Coimbatore, for providing financial support to the work.

## Authors' contributions

GGK has carried out the research work, MC drafted the manuscript for publication, MV participated in the study's design and SRP performed the statistical analysis. All authors read and approved the final manuscript.



**Fig. 3b.** Instant dip from *M. oleifera* flowers.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interest to declare.

**Ethical issues :** None

## References

- Kasolo JN, Bimenya GS, Ojok L, Ochieng J, Ogwal-okeng JW. Phytochemicals and uses of *Moringa oleifera* leaves in Ugandan rural communities J Med Plants Res. 2010; (4):753-57
- Rockwood JL, Anderson BG, Casamatta DA. Potential uses of *Moringa oleifera* and an examination of antibiotic efficacy conferred by *M. oleifera* seed and leaf extracts using crude extraction techniques available to underserved indigenous populations. Int J Phytotherapy Res 2013; 3(2):61-71
- Aslam M, Anwar F, Nadeem R, Rashid U, Kazi TG, Nadeem M. Mineral composition of *Moringa oleifera* leaves and pods from different regions of Punjab Pakistan Asian J Plant Sci. 2005;4(4):417-21. <https://doi.org/10.3923/ajps.2005.417.421>
- Asante WJ, Nasare IL, Tom-Dery D, Ochire-Boadu K, Kentil KB. Nutrient composition of *Moringa oleifera* leaves from two agro ecological zones in Ghana Afr J Plant Sci. 2014;8(1):65-71. <https://doi.org/10.5897/AJPS2012.0727>
- Fuglie LJ. The Moringa Tree: A local solution to malnutrition Church World Service in Senegal. Nature's Pharm. 2003;221:22-35.
- Berkovich L, Earon G, Ron I, Rimmon A, Vexler A, Lev-Ari S. *Moringa oleifera* aqueous leaf extract down-regulates nuclear factor-kappaB and increases cytotoxic effect of chemotherapy in pancreatic cancer cells, BMC Complement. Altern. Med. 2013; (13): 1-7. <https://doi.org/10.1186/1472-6882-13-212>
- Barminas JT, Charles M, Emmanuel D. Mineral composition of non-conventional leafy vegetables Plant Foods Hum Nutr. 1998;53:29-36 <https://doi.org/10.1023/A:1008084007189>
- Oduro I, Ellis WO, Owusu D. Nutritional potential of two leafy vegetables: *Moringa oleifera* and *Ipomoea batatas* leaves, Sci Res Essays 2008;3(2): 57-60.
- Otunola GA, Arise AK, Sola-Ojo FE, Nmomo IO, Teye AA. Effect of addition of Moringa leaf by-product (Leaf-Waste) on proximate and sensory characteristics of cookies Agrosearch. 2013;13(1):69-76. <https://doi.org/10.4314/agrosh.v13i1.7>
- Srinivasamurthy S, Yadav U, Sahay S, Singh A. Development of muffin by incorporation of dried *Moringa oleifera* (Drumstick) leaf powder with enhanced micronutrient content Development. Int J Food Sci Nutr 2017;2(4):173-178.

11. Haque MO, Moni MA, Rani S, Hasan MR, Sarker MR, Rahman MS, et al. A. Standardization of recipe of value-added functional aloe vera drink enriched with Moringa leave. Malays J Halal Res. 2022;(1):66-69. <https://doi.org/10.2478/mjhr-2022-0007>
12. Mutiara T, Titi ES, Estiasih W. Effect lactagogue moringa leaves (*Moringa oleifera* Lam) powder in rats J Basic Appl Sci Res 2013; (3):430-34.
13. Salem AS, Salama WM, Ragab WA. Prolonged shelf life of sour cream by adding *Moringa oleifera* Leaves extract (MOLE) or *Moringa oleifera* oil (MOO) Am J Food Technol. 2005;10:58-67. <https://doi.org/10.3923/ajft.2015.58.67>
14. Salem AS, Salama WM, Hassanein AM, El-Ghandour HMA. Enhancement of nutritional and biological values of Labneh by adding dry leaves of *Moringa oleifera* as innovative dairy products. World Appl Sci J. 2013;22(11):1594-602.
15. Reddy BH, Pradeep P, Padmavathi TVN. Development and evaluation of value-added products from Moringa leaves. J Pharmacogn Phytochem. 2020;9(5):660-63. <https://doi.org/10.22271/phyto.2020.v9.i5j.12304>
16. Lokshin M, Das Gupta M, Gragnolati M, Ivaschenko O. Improving child nutrition the integrated child development services in india. Dev Change. 2005;36(4):613-40. <https://doi.org/10.1111/j.0012-155X.2005.00427.x>
17. Kamble DB, Bashir K, Singh R, Rani S. Effect of *Moringa oleifera* pod addition on the digestibility, cooking quality and structural attributes of functional pasta. J Food Process Preserv. 2022;46(1):e16163. <https://doi.org/10.1111/jfpp.16163>
18. Mistry EM, Patel SM, Pinto S, Modha HM. Utilization of Moringa pod powder as a value-added ingredient in Lassi, Research & Reviews: J Dairy Sci Technol. 2018;7:6-17.
19. Madane P, Das AK, Pateiro M, Nanda PK, Bandyopadhyay S, Jagtap P, et al. Drumstick (*Moringa oleifera*) flower as an antioxidant dietary fibre in chicken meat nuggets Foods. 2019;8(8):307. <https://doi.org/10.3390/foods8080307>
20. Moyo B, Masika PJ, Hugo A, Muchenje V. Nutritional characterization of Moringa flower (*Moringa flower oleifera* Lam.). African J Biotechnol. 2011;10(60):12925-2933. <https://doi.org/10.5897/AJB10.1599>
21. Bolarin FM, Oke MO. Effect of addition of moringa (*Moringa oleifera* L.) seed flour on the quality characteristics of mayonnaise International Journal of Multidisciplinary Research and Publications. 2018;1(1):11-17.
22. Fuglie LJ. The Moringa Tree: A local solution to malnutrition Church World Service in Senegal, 2005.
23. Aluko O, Brai MR, Adedore AO. Materials evaluation of sensory attributes of snack from maize-moringa seed flour blends. Int J Nutr Food Eng. 2013;7(10):944-46.
24. Steve IO, Babatunde OI, Chemical compositions and nutritional properties of popcorn-based complementary foods supplemented with *Moringa oleifera* leaves Flour 2013;2(6):117-32. <https://doi.org/10.5539/jfr.v2n6p117>
25. Divi SM, Bellamkonda R, Dasireddy SK. Evaluation of antidiabetic and anti hyperlipidemic potential of aqueous extract of *Moringa flower oleifera* in fructose fed insulin resistant and STZ induced diabetic wistar rats: a comparative study Asian J Pharm Clin Res. 2012;5(1):67-72
26. Nambiar VS, Parnami S. Standardization and organoleptic evaluation of drumstick (*Moringa oleifera*) leaves incorporated into traditional Indian recipes. Trees for Life Journal. 2008;3(2):1-7.
27. Abou-zaid AA, Nadir AS. Quality evaluation of nutritious chocolate and halawa tahinia produced with moringa (*Moringa oleifera*) leaves powder. Middle East J Appl Sci. 2014;4(4):1007-015.
28. Arise AK, Arise RO, Sanusi MO, Esan OT, Oyeyinka SA. Effect of *Moringa oleifera* flower fortification on the nutritional quality and sensory properties of weaning food Croat J Food Sci Technol. 2014;6(2):65-71.
29. Adejumo OE, Kolapo AL, Folarin AO. *Moringa oleifera* Lam. (Moringaceae) grown in Nigeria: *in vitro* antiscikling activity on deoxygenated erythrocyte cells. J Pharm Bioallied Sci. 2012;4(2):118-22. <https://doi.org/10.4103/0975-7406.94812>
30. Shank LP, Riyathong T, Lee VS, Dheeranupattana S. Peroxidase activity in native and callus culture of *Moringa oleifera* Lam. Journal of Medical and Bioengineering. 2013;2(3):163-67. <https://doi.org/10.12720/jomb.2.3.163-167>
31. Owusu D, Oduro I, Ellis WO. Development of crackers from cassava and sweet potato flours using *Moringa oleifera* and *Ipomoea batatas* leaves as fortificant. Am J Food Nutr. 2011;1(3):114-122. <https://doi.org/10.5251/ajfn.2011.1.3.114.122>
32. Raphaël KJ, Christian KT, Juliano RS, Lisita F, Soultan MY, Hervé MK, Alexis T. Effects of substituting soybean with *Moringa oleifera* meal in diets on laying and eggs quality characteristics of KABIR chickens. J Anim Nutr. 2015;1:1-6.
33. Olugbemi TS, Mutayoba SK, Lekule FP. Effect of Moringa (*M. oleifera*) inclusion in cassava-based diets fed to broiler chickens Int J Poult Sci. 2010;9(4):363-67. <https://doi.org/10.3923/ijps.2010.363.367>
34. Miyoshi N, Uchida K, Osawa T, Nakamura Y. A link between benzyl isothiocyanate-induced cell cycle arrest and apoptosis: involvement of mitogen-activated protein kinases in the Bcl-2 phosphorylation. Cancer Res. 2004;64(6):2134-142. <https://doi.org/10.1158/0008-5472.CAN-03-2296>
35. Baker K, Marcus CB, Huffman K, Kruk H, Malfroy B, Doctrow SR. Synthetic combined superoxide dismutase/catalase mimetics are protective as a delayed treatment in a rat stroke model: a key role for reactive oxygen species in ischemic brain injury. J Pharmacol Exp Ther. 1998;284(1):215-21.
36. Choudhary MK, Bodakhe SH, Gupta SK. Assessment of the antiulcer potential of *Moringa oleifera* root-bark extract in rats. J Acupunct Meridian Stud. 2013;6(4):214-20. <https://doi.org/10.1016/j.jams.2013.07.003>
37. Jahn SA, Musnad, HA, Burgstaller H. The tree that purifies water: cultivating multipurpose Moringaceae in the Sudan. Unasylva. 1986;38(152):23-28.
38. Viera GHF, Mourão JA, Ângelo AM, Costa RA, Vieira RHFD. Antibacterial effect (*in vitro*) of *Moringa oleifera* and *Annona muricata* against Gram positive and Gram-negative bacteria. Rev Inst Med Trop Sao Paulo. 2010;52(3):129-32. <https://doi.org/10.1590/S0036-46652010000300003>
39. Al-Malki AL, El-Rabey HA. The antidiabetic effect of low doses of *Moringa oleifera* Lam. seeds on streptozotocin induced diabetes and diabetic nephropathy in male rats Biomed Res Int. 2015;2015(1):381040. <https://doi.org/10.1155/2015/381040>
40. Kamalakkannan N, Prince PSM. Antihyperglycaemic and antioxidant effect of rutin, a polyphenolic flavonoid, in streptozotocin-induced diabetic wistar rats. Basic Clin Pharmacol Toxicol. 2006;98(1):97-103. [https://doi.org/10.1111/j.1742-7843.2006.pto\\_241.x](https://doi.org/10.1111/j.1742-7843.2006.pto_241.x)
41. Chen M, Verdes RP. Elucidation of bactericidal effects incurred by *Moringa oleifera* and Chitosan J US SJWP. 2009; 4:65-79.
42. Adeyemi OS, Elebiyo TS. *Moringa oleifera* supplemented diets prevented nickel-induced nephrotoxicity in Wistar rats. J Nutr Metab. 2014;2014(1):958621. <https://doi.org/10.1155/2014/958621>
43. Mthiyane FT, Dlodla PV, Ziqubu K, Mthembu SX, Muvhulawa N, Hlengwa N, Nkambule BB, Mazibuko-Mbeje SE. Corrigendum: A review on the antidiabetic properties of Moringa oleifera extracts: Focusing on oxidative stress and inflammation as main therapeutic targets. Front Pharmacol. 2023;14:1142410. <https://doi.org/10.3389/fphar.2023.1142410>