



REVIEW ARTICLE

Effect of landfill leachates on urban soil: A review

Selvakumar Jagasri¹, J Kannan^{2*}, R Jayashree³, S Sheeba¹, K Prabakaran⁴, R M Jayabalakrishnan¹, R Murugaragavan¹, C Poornachandra³ & S Madhusree⁵

¹Department of Soils and Environment, Agricultural College and Research Institute, Madurai - 625104, India

²Department of Environmental Science, ICAR- Krishi Vigyan Kendra, Aruppukottai, Virudhunagar - 626107, India

³Department of Environmental Science, Tamil Nadu Agricultural University, Coimbatore - 641003, India

⁴Department of Agricultural Economics, Agricultural College and Research Institute, Madurai - 625104, India

⁵Department of Agronomy, Anbil Dharmalingam, Agricultural College and Research Institute, Trichy - 62009, India

*Email: kannan.j@tnau.ac.in

 OPEN ACCESS

ARTICLE HISTORY

Received: 22 August 2024

Accepted: 15 September 2024

Available online

Version 1.0 : 09 October 2024

Version 2.0 : 12 October 2024



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Jagasri S, Kannan J, Jayashree R, Sheeba S, Prabakaran K, Jayabalakrishnan RM, Murugaragavan R, Poornachandra C, Madhusree S. Effect of Landfill Leachates on Urban Soil: A Review. Plant Science Today. 2024; 11(4): 485-493. <https://doi.org/10.14719/pst.4767>

Abstract

The increasing generation of Municipal Solid Waste (MSW) is a significant global concern, with landfills receiving around 1.4 billion tonnes of MSW yearly. Inadequate landfill management contributes to environmental degradation, with landfill leachate being a substantial outcome of MSW decomposition. Leachate contains inorganic nutrients, volatile and dissolved organic molecules and heavy metals and its properties vary depending on waste composition, moisture content and seasonal elements. Heavy metals found in leachate include Pb, Cu, Cr, Ni, Mn, Hg, Fe, Zn and Cd and Emerging Organic Contaminants (EOCs) such as Persistent Organic Pollutants (POPs), Endocrine Disrupting Chemicals (EDC), pharmaceuticals and Personal Care Products (PCPs) are also prevalent. Microplastics (MPs) have been found in raw leachate samples at concentrations ranging from 49.0 ± 24.3 to 507.6 ± 37.3 items/L. Landfill leachate production ranks among the most aggressive pollutants to the environment, particularly to soil and poses a danger of contaminating both surface and groundwater. This review examines the potential impacts of landfill leachate on soil quality and the broader implications of this phenomenon, summarizing recent scientific studies and presenting the direct and indirect effects of leachate on soil based on the literature. Bibliometric analysis of publications in the Scopus database reveals a growing scholarly interest in this topic, with the number of publications in the Science Citation Index (SCI) database increasing dramatically to over 464 articles between 2009 and 2024.

Keywords

Emerging Organic Contaminants (EOCs); environmental degradation; heavy metal; Municipal Solid Waste (MSW); soil contamination

Introduction

Worldwide, the increasing generation of Municipal Solid Waste (MSW) is becoming a significant concern. On average, about 1.04 kilograms of waste is produced globally per person daily. Waste generation rates differ significantly across countries, ranging from 0.5 to 2.3 kilograms per person daily. By 2050, 3.40 billion tonnes of MSW will be generated globally, with 19 % and 40 % growth rates in industrialized and developing countries, respectively (1). The statistics on global MSW generation are shown in Table 1 (2). Top 10 Indian City's waste generation are shown in Table 2 (3). Most

Table 1. Statistics of global MSW generation

Region	Total MSW (million tonnes)	MSW per capita (kg/person/day)
North America	320	2.3
Central America and the Caribbean	80	0.9
South America	140	0.95
Northern Europe	60	1.3
Western Europe	110	1.4
Southern Europe	80	1.2
Eastern Europe	120	1.0
West Asia and North Africa	150	0.8
Sub-Saharan Africa	220	0.55
Central and South Asia	280	0.5
East and South-East Asia	580	0.75
Oceania	15	0.5
Australia and New Zealand	20	1.4

Source: UNEP, 2024

MSW comes from everyday activities, including residential, commercial and institutional sources. The increasing volume of solid waste highlights the need for safe landfills. Many cities find landfills to be an unsuitable technique for safe disposing of MSW. Landfills receive around 1.4 billion tonnes of MSW annually, accounting for 70 % of total MSW. Landfills in India need 1240 acres of land annually, with just 21% of MSW being adequately managed and disposed of. However, the remaining MSW is disposed of in unsanitary landfills without sufficient treatment (4), which degrades the environment. According to (5) and (6), inadequate landfill management can lead to environmental degradation. The health impacts studied included mortality, adverse birth and neonatal outcomes, cancer, respiratory problems, gastroenteritis, vector-borne illnesses, mental health issues and cardiovascular diseases. However, occupational risks were not considered in the assessment (7).

The significant outcome of MSW decomposition is the generation of landfill leachate, which is the aqueous effluent produced from solid waste due to its physical, chemical and biological transformation within landfills (8). Municipal solid waste (MSW) composition varies widely across different regions but generally consists of a combination of biodegradable and non-biodegradable materials derived from organic and inorganic sources. MSW is typically collected from residential areas, offices, institutions and commercial establishments, comprising items such as organic waste (e.g., food scraps and yard trimmings), paper, plastics, metals, glass and a variety of other materials, including electronic waste, inert substances, pharmaceuticals and debris from construction, demolition and renovations. The approach to managing MSW differs by locality but generally follows three key stages: (i) waste generation at the source, (ii) collection and transportation and (iii) disposal, processing and treatment (9).

Table 2. Top 10 Indian Cities and Their Waste Generation Patterns

Cities	MSW generation in tons per day (TPD) in India, CPCB				
	1971	1999	2004	2010	2015
Mumbai	2039	5355	5320	6500	11000
Delhi	766	4000	5922	6800	8700
Chennai	508	3124	3036	4500	5000
Hyderabad	593	1556	2187	4200	4000
Kolkata	1574	3692	2653	3670	4000
Bangalore	529	2000	1669	3700	3700
Ahmedabad	381	1683	1302	2300	2500
Surat	74	900	1000	1200	1680
Pune	205	700	1175	1300	1600
Jaipur	178	580	904	810	1000

Source: Dutta, 2020

MSWs contain organic biodegradable components and compacted waste layers, creating an anaerobic environment in landfills (10). Most landfills receive and dispose of municipal, commercial and mixed industrial garbage. One tonne of landfilled waste produces approximately 0.2 m³ of landfill leachate during decomposition (11). Leachates from various landfills have similar constituents (12) and contain inorganic nutrients, volatile and dissolved organic molecules and heavy metals, which occur when water flows through a landfill and absorbs dissolved elements from degraded garbage (13). A well-designed landfill can reduce leachate leaking into the soil. To improve landfills, surface runoff should be altered and proper vegetation and leachate should be collected and pumped to a treatment facility (14).

Landfill leachate is characterized using standard criteria such as COD, TOC, BOD, suspended particles, pH, ammonia and heavy metal concentrations. The BOD 5/ COD and COD/TOC ratios indicate the biodegradability and oxidation of organic carbon. Several variables influence landfill leachate quality, including waste type, operational conditions, climate, hydrogeology and landfill age (15). Landfill leachate properties vary depending on waste composition, moisture content and seasonal elements such as temperature and precipitation (16). Microplastics (MPs) concentration in raw leachate samples ranged from 49.0 ± 24.3 to 507.6 ± 37.3 items/L. A potential correlation was found between the concentration of MPs in raw leachate samples from landfill sites and the annual leachate (17). Heavy metals found in leachate include Pb, Cu, Cr, Ni, Mn, Hg, Fe, Zn and Cd (18), with different concentrations for each landfill. Heavy metals remain in polluted sites for an extended period and, unlike other pollutants, cannot be degraded chemically or biologically (19). Recent years have seen a lot of attention paid to Emerging Organic Contaminants (EOCs), like Persistent Organic Pollutants (POPs), Endocrine Disrupting Chemicals (EDC), pharmaceuticals, Personal Care Products (PCPs), antibiotic resistance genes and disinfection by-products, due to their prevalence in landfill leachate and their potential for harm to the environment and people

(20). (17) found MP concentrations in raw leachate samples ranging from 49.0 ± 24.3 to 507.6 ± 37.3 items/L. Over the past two decades, 172 pharmaceutical and PCPs have been found in landfill leachate worldwide, including antibiotics, anti-inflammatories, stimulants and beta-blockers (21). Due to its properties and content, landfill leachate production ranks among the most aggressive pollutants in the environment today, mostly in soil and poses a danger of contaminating both surface and groundwater (22).

This review aims to examine the potential impacts of landfill leachate on soil quality and the broader implications of this phenomenon. The main aims of this review were to (i) summarise the most recent scientific studies on landfill leachate and (ii) present the direct and indirect impacts of leachate on soil based on the literature. Studies in the literature have examined the impact of landfill leachates on soil physical, chemical and biological properties. Modern remedial techniques to treat soil degradation from landfill leachate are also presented here.

Scientific focus on leachate impact on soil

The Scopus database was selected to methodically monitor the effects of landfill leachate on soil among reputable publications because of its consistency in citation records. Only peer-reviewed English-language literature was the subject of this literature search. Bibliometric data were gathered by deciding on the best sources of information, establishing search parameters and creating the dataset. Following data cleaning and anomaly identification, bibliometric analysis was performed (Fig. 1). About 653 publications with titles, abstracts, and keywords like "landfill leachate impact on soil"-such as "landfill AND leachate AND impact AND on AND soil"-were analyzed when they were retrieved on July 3, 2024. After that, 168 publications were found using Boolean search terms like "landfill AND leachate AND impact AND on AND soil." The growing number of papers about the effects of landfill leachate on soil during the previous 16 years (from 2009 to 2024) indicates a growing scholarly interest in this topic (Kurniawan et al., 2021d). Consequently, the total number of publications in the Science Citation Index (SCI) database (2009-2024) concerning the effect of landfill leachate on soil increased dramatically to over 464 articles (Fig. 2) (23).




Fig. 2. Trends of landfill leachate-related publications in the body of knowledge (2009-2024)

Bibliometric analysis on current hotspot

A bibliometric analysis was carried out using data gathered from Scopus and VOS viewer to visualize the network, as shown in Fig 3.

Searching the Scopus database with the keywords "landfill AND leachate AND impact AND on AND soil," about 168 documents were found. All key terms were used as the unit of analysis in a co-occurrence analysis. It was decided that ten keyword occurrences would be minimal. 108 keywords out of 3284 matched the criterion. The 108 keywords exhibit significant connectedness.

Impact

Soil structure

Xu et al. (24) found that increasing landfill leachate concentrations decreased soil strength, leading to plastic deformation. The dislocation between soil particles and plastic lateral deformation occurred due to leachate pollution and axial load, ultimately destroying the soil structure (25). Giri and Reddy (26) showed that leachate significantly influences pore water pressure and forms numerous pores in the soil. Meanwhile, water adsorption by soil particles increased (27).

At higher landfill leachate concentrations, the maximum pore radius saturated with leachate expanded from 1.03 to 1.18 μm , while the radius of other pores grew from 11.01 to 135.73 μm . Pore sizes in leachate-contaminated soil were primarily between 0.02-1 μm and 3-12 μm (28). Increased leachate concentrations led to




Fig. 1. Method of Bibliometric Analysis




Figure 3. Vosviewer network visualization on recent hotspots

greater soil porosity, forming an unstable honeycomb structure and reducing particle uniformity. The specific surface area rapidly increased, stabilizing between 500 and 650 kg/m². Additionally, higher leachate levels caused a significant decrease in particle size and a sharp rise in pore volume (29).

Compaction

(30) found that soil contaminated with leachate exhibits lower dry density, likely due to chemical interactions between the leachate and soil pore fluid properties. The study suggests that incorporating leachate into the soil could improve compaction efficiency, potentially reducing soil volume in landfill cells. Nayak et al. (31) observed a decrease in maximum dry density is likely due to these chemical reactions between the acidic leachate and the soil. At high leachate concentrations, an excessive amount of leachate in the soil can trigger further chemical reactions between the acidic leachate and the soil particles. The compaction study showed that as the percentage of liquid leachate increased, both the maximum dry unit weight and the optimum moisture content decreased (32). Adding moisture facilitates compaction by making the soil easier to knead and capable of achieving higher dry density. However, the dry density decreases at higher moisture content as the soil becomes more saturated with water. Liquid leachate increases soil saturation, contributing to the observed reductions in maximum dry unit weight and optimum moisture content (33).

Hydraulic conductivity

According to Zheng et al. (34) an increase in leachate concentration results in an increase in the soil's hydraulic

conductivity; high ion content in leachate causes an increase in mass loss due to the dissolving of clay minerals; channels emerge in the soil; and adequate pore space expands. The more significant permeability channel formed by the soil particles and the infiltration of heavy metal ions into the soil causes an increase in hydraulic conductivity. A summary of studies on the impact of heavy metals on hydraulic conductivity found in the literature is furnished in Table 3 (35).

Nayak et al. (31) observed changes in soil structure after leachate contamination. They found that replacing pore water with leachate increased the void ratio of the soil. The increase in pore fluid volume and hydraulic conductivity was attributed to the leachate's capacity to dissolve clay minerals within the soil. Xie et al. (36) studied soil compacted with various concentrations of leachate and observed that hydraulic conductivity to leachate was consistently higher than that to demineralized water across all compacted samples. This difference was primarily attributed to the lower viscosity of leachate than water. Long-term soil exposure to leachate led to a notable decrease in hydraulic conductivity to both leachate and water, especially in samples with more significant voids. This decrease was due to reduced active pore space, influenced by thicker diffuse double layers, clay particle rearrangement, chemical precipitation and biofilm formation within soil pores. Microbial activity significantly reduces soil hydraulic conductivity (37-39). This reduction occurs as biofilms and colonies form on mineral particle surfaces and grow within soil pores, obstructing them and contributing to decreased hydraulic conductivity (40,41).

Table 3. Impact of Heavy metals on Hydraulic conductivity of soil

Contaminants	Hydraulic conductivity (cm/s)	Change in hydraulic conductivity	Reference
Pb	4.7×10^{-9}	Hydraulic conductivity was found to be close to each other because the soil samples were mainly composed of sand.	(34)
Cu	4.8×10^{-9}		
Pb + Cu	4.3×10^{-9}		
Pb + Cu + Cd	2.4×10^{-9} (m/s)		
0-4 mg/L Pb ⁺²	~Increase from 1.25×10^{-9} to 2.2×10^{-9}	As the concentration increased, hydraulic conductivity also increased	(69)
0-40 mg/L Pb ⁺²	Decrease from 2×10^{-5} to 7.8×10^{-7}	As the concentration increased, hydraulic conductivity decreased	(70)
0-10 mM Pb ⁺²	Increase from 10^{-9} to 10^{-7}	As the concentration increased, hydraulic conductivity also increased	(71)
0 mg/kg Pb or Zn ⁺²	2.2×10^{-8}	As the concentration increased, hydraulic conductivity also increased	(34)
1,000 mg/kg Pb ⁺²	5×10^{-8}		
1,000 mg/kg Zn ⁺²	4.8×10^{-8} (m/s)		
0-10 g/L Cu ⁺²	3.54×10^{-6} 42.25×10^{-6}	As the concentration increased, hydraulic conductivity also increased.	(72)
0-100 ppm Cu ⁺²		As the concentration increased, hydraulic conductivity also increased.	(73)
0-1,000 ppm Cu ⁺²	1.7 times increase		
0-100 ppm Pb ⁺²	2.6 times increase one time increase		
0-1,000 ppm Pb ⁺²	1.2 times increase		
0-100 ppm Zn ⁺²	1.4 times increase 2.2 times increase		
0-1,000 ppm Zn ⁺²			

Soil Nutrient Imbalance

Changes in soil pH can limit nutrient availability (42). Rahman et al. (43) reported that copper, zinc and nickel are crucial micronutrients for plants in small amounts but become toxic in excess.

Rao (44) reported that nitrogen levels in the contaminated soils were notably high, ranging from 115 to 262 kg/acre, with the control soil sample showing a lower value of 62 kg/acre. The phosphorus content in the dump yard soils varied between 73 and 91 kg/acre, while the control site had a lower 34 kg/acre value. The elevated nitrogen and phosphorus levels in dump site soil are likely due to the high organic matter (45). Potassium levels in the dump site soils ranged from 157 to 363 kg/acre, compared to a low of 15 kg/acre at the control site. Although potassium is essential for plant growth, anthropogenic activities can elevate its levels, potentially contaminating groundwater. According to Agbeshie et al. (46), the high nutrient content at the dump site, mainly the organic carbon and exchangeable bases, significantly affected soil bulk density, porosity and nutrient availability. High concentrations of calcium, magnesium, sodium, potassium, ammonium, iron, chloride, sulfate, nitrate and hydrogen carbonate ions in leachate and soil increase osmotic pressure, hindering water uptake by plant roots and impairing growth (47). Letsoalo (48) suggests that essential nutrients and chromium affect plants' absorption of calcium (Ca²⁺) and magnesium (Mg²⁺) through soil interactions. Dimethyl arsenic acid in soil reduces concentrations of essential macronutrients (P, K, Ca, Mg) and micronutrients (B, Cu, Fe, Mn) in plants (49).

Soil microbes

(50) reported that bacteria found at waste or leachate dumpsites can include *Arthrobacter*, *Bacillus*, *E. coli*, *Klebsiella*, *Micrococcus*, *Proteus*, *Serratia marcescens*, *Klebsiella aerogenes*, *Staphylococcus aureus*, *Alcaligenes*

sp., *Proteus mirabilis* and *Salmonella*. Fungi isolated from waste dumpsites include *Aspergillus*, *Fusarium*, *Mucor*, *Penicillium*, *Rhizopus* and *Saccharomyces*. *Aspergillus niger*, *Aspergillus flavus*, *Rhizopus* and yeast species were explicitly isolated from dumpsite leachates. Wydro et al. (51) experimented using soil treated with different doses of leachate (50 LL and 100 LL). They found that the highest total number of bacteria was observed in pots treated with 50 LL (1.05×10^7 cfu/g DM, T1), while the lowest number was in the control pots (1.43×10^6 cfu/g DM, T3). According to (52), leachate (LL) contains a mixture of soluble organic matter, heavy metals, PAHs and other toxic substances, which, when introduced into the soil, can affect its activity and reduce the number of microorganisms (Fig 4.) (53). The presence of these toxic substances can interfere with the adaptability of some organisms, resulting in a decrease in their numbers (54). Wydro et al. (51) also reported that leachate alters the structure of the microbial community, as indicated by the T-RFLP approach, affecting microbial richness and relative abundance in the soil. Daniel et al. (55) suggested that heavy metals indirectly impact soil enzymatic activities by altering the microbial community responsible for enzyme synthesis. These heavy metals affect soil microorganisms by modifying their diversity, population size and overall activity within the soil microbial communities. Heavy metals like lead, silver and cadmium penetrate bacterial plasma membranes and generate superoxide ions in the cytosol. These ions, converted by Super Oxide Dismutase (SOD) into hydrogen peroxide or hydroxyl radicals, oxidize lipids, proteins and DNA. Reactive Oxygen Species (ROS) and other oxidative intermediates further damage cellular components. Cells produce antioxidant enzymes such as catalase, SOD and glutathione peroxidase to mitigate ROS. However, the oxidative stress caused by heavy metals can result in apoptosis, necrosis, tissue damage and malignancy (53).




Figure 4. Mechanism of Heavy Metal Toxicity on Bacteria

Heavy metal

Heavy metals are major pollutants in landfill leachate and can remain in landfills for about 150 years if leaching occurs at 400 mm/year (56, 57). Their toxicity disrupts the biological balance and impairs natural purification processes (58). Leachate production and heavy metal mobility are influenced by rainfall (posing risks to soil, groundwater and surface water (59). Non-threshold pollutants like arsenic, chromium (VI), cadmium, mercury and lead are toxic even in small amounts (60,61). Torkashvand et al. (56) reported copper, cadmium, lead, iron, and nickel concentrations in landfill leachate from Iran as 1, 0.45, 0.85, 14 and 1.1 mg/L, respectively. Pasalari et al. (62) found manganese levels in Iranian landfill leachate ranging from 3.2 to 8.1 mg/L. Beinabaj et al. (63) indicated that iron concentrations in Nigeria were the highest among the metals, at 22.94 mg/L. Johar et al. (64) discovered the highest concentrations of cadmium (Cd) and silver (Ag) in soil samples from a landfill in New Delhi, India, highlighting the landfill as a significant source. The soil exhibited a higher Cd and Ag adsorption capacity than iron (Fe) and copper (Cu). The high level of transferable Cd is particularly concerning due to its potential for significant plant uptake and accumulation (65).

Mitigation measures

Landfill leachate significantly threatens soil and water resources, leading to degradation. Without adequate containment measures, leachate can directly contaminate surrounding soil and seep into groundwater, exacerbated by rainfall. Various industrial and scientific initiatives have been implemented to mitigate leachate release, each tailored to specific environmental conditions and with varying biomedical implications. The landfill liner is crucial to preventing leachate from seeping into the subsoil (66). The foundation of a landfill site should be designed to support the weight of the overlying waste and cover

material. The foundation material must have sufficient compressive strength to bear this load.

In some cases, grouting or other techniques may be needed to reinforce the foundation. For a landfill liner to be effective, it must exhibit specific properties such as swelling behaviour, strength and low permeability. Clay with a higher content of Montmorillonite, combined with overburden pressure, needle punching density, and areal density, demonstrates better self-healing properties and low hydraulic conductivity. However, hydraulic conductivity increases with higher water pressure in clayey soil (67). Using nanotechnology, (68) discussed the application of nanoclay and nanofiber filters during the landfill stage for solid waste management to control leachate leakage from landfill liners.

Conclusion

Landfill operations are vital for waste disposal, but landfill leachate, produced by chemical and biological reactions within landfills, can contaminate soil and groundwater, posing environmental health risks. This review explored the effects of landfill leachate on soil structure, hydraulic conductivity and heavy metal impact. Recent innovations, such as advanced landfill liners with nanotechnology, are essential for preventing leachate contamination. Developing new bioinoculants shows promise in reducing heavy metals in landfills. Biochar and Hydrochar are effective for treating landfill leachate due to their customizable adsorption properties, though challenges like limited research and the difficulty of scaling laboratory methods to treat the average 167 million tonnes of leachate produced globally. Further research could enhance their effectiveness, mitigating waste and providing sustainable ecosystem services.

Acknowledgements

I sincerely thank the Department of Soils and Environment, Agricultural College and Research Institute, Madurai and the Department of Environmental Sciences, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India, for their support.

Authors' contributions

J K designed the study and wrote the protocol. S Jagasri and R J wrote the first draft of the manuscript. S S and C P managed the analyses of the study. K P performed the analysis. R M J and R M and S M managed the literature searches. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None.

Did you use generative AI to write this manuscript?

No

Declaration of generative AI and AI-assisted technologies in the writing process

No AI tools are used for manuscript preparation

References

- Kaza S, Yao LC, Bhada-Tata P, Van Woerden F. What a waste 2.0: a global snapshot of solid waste management to 2050. World Bank Publications; 2018. <https://doi.org/10.1596/978-1-4648-1329-0>
- United Nations Environment Programme. Global Waste Management Outlook 2024: Beyond an age of waste - Turning rubbish into a resource. Nairobi; 2024.
- Dutta A, W Jinsart, Waste generation and management status in the fast-expanding Indian cities: a review. Journal of the Air & Waste Management Association, 2020. 70(5): 491-03. <https://doi.org/10.1080/10962247.2020.1738285>
- Prajapati KK, Yadav M, Singh RM, Parikh P, Pareek N, Vivekanand V. An overview of municipal solid waste management in Jaipur city, India-Current status, challenges and recommendations. Renewable and Sustainable Energy Reviews. 2021; 152. <https://doi.org/10.1016/j.rser.2021.111703>
- Cossu R. Groundwater contamination from landfill leachate: when appearances are deceiving! Waste Management. 2013; 33(9):1793-794. <https://doi.org/10.1016/j.wasman.2013.07.002>
- Peng Y. Perspectives on technology for landfill leachate treatment. Arabian Journal of Chemistry. 2017; 10. <https://doi.org/10.1016/j.arabjc.2013.09.031>
- Vinti G, Bauza V, Clasen T, Medlicott K, et al. Municipal solid waste management and adverse health outcomes: A systematic review. International Journal of Environmental Research and Public Health. 2021; 18(8). <https://doi.org/10.3390/ijerph18084331>
- Youcai Z. Pollution control technology for leachate from municipal solid waste: landfills, incineration plants, and transfer stations. Butterworth-Heinemann; 2018.
- Nanda S, Berruti F. A technical review of bioenergy and resource recovery from municipal solid waste. Journal of Hazardous Materials. 2021; 403. <https://doi.org/10.1016/j.jhazmat.2020.123970>
- Hussein M, Yoneda K, Mohd-Zaki Z, Amir A, Othman N. Heavy metals in leachate, impacted soils and natural soils of different landfills in Malaysia: An alarming threat. Chemosphere. 2021; 267. <https://doi.org/10.1016/j.chemosphere.2020.128874>
- Diamadopoulos E, Samaras P, Dabou X, Sakellariopoulos GP. Combined treatment of landfill leachate and domestic sewage in a sequencing batch reactor. Water Science and Technology. 1997; 36(2-3):61-68. <https://doi.org/10.2166/wst.1997.0483>
- Serdarevic A. Landfill leachate management-control and treatment. In: Advanced Technologies, Systems and Applications II: Proceedings of the International Symposium on Innovative and Interdisciplinary Applications of Advanced Technologies (IAT). Springer; 2018. https://doi.org/10.1007/978-3-319-71321-2_54
- Mahmoud H. Trends in the remediation methods for landfill leachate. Nigerian Journal of Engineering Science and Technology Research. 2023; 9(2):149-60.
- Zhou W, Chai J, Xu Z, Qin Y, Cao J, Zhang P. A review of existing methods for predicting leachate production from municipal solid waste landfills. Environmental Science and Pollution Research. 2024; 31(11):16131-149. <https://doi.org/10.1007/s11356-024-32289-y>
- Dabaghian Z, Peyravi M, Jahanshahi M, Rad AS. Potential of advanced nano-structured membranes for landfill leachate treatment: A review. ChemBioEng Reviews. 2018; 5(2):119-38. <https://doi.org/10.1002/cben.201600020>
- Costa AM, Alfaia RGSM, Campos JC. Landfill leachate treatment in Brazil-An overview. Journal of Environmental Management. 2019; 232:110-16. <https://doi.org/10.1016/j.jenvman.2018.11.006>
- Qin ZH, Siddiqui MA, Xin X, Mou JH, et al. Identification of microplastics in raw and treated municipal solid waste landfill leachates in Hong Kong, China. Chemosphere. 2024; 351. <https://doi.org/10.1016/j.chemosphere.2024.141208>
- Jayanthi B. Bioaugmentation and phytoremediation of heavy metal from leachate contaminated soil. University of Malaya; 2018.
- Wuana RA, Okieimen FE. Heavy metals in contaminated soils: A review of sources, chemistry, risks and best available strategies for remediation. International Scholarly Research Notices. 2011. <https://doi.org/10.5402/2011/402647>
- Iskander SM, Zhao R, Pathak A, Gupta A, et al. A review of landfill leachate induced ultraviolet quenching substances: Sources, characteristics and treatment. Water Research. 2018; 145:297-11. <https://doi.org/10.1016/j.watres.2018.08.035>
- Yu X, Sui Q, Lyu S, Zhao W, Liu J, et al. Municipal solid waste landfills: An underestimated source of pharmaceutical and personal care products in the water environment. Environmental Science & Technology. 2020; 54(16):9757-768. <https://doi.org/10.1021/acs.est.0c00565>
- Gao M, Li S, Zou H, Wen F, Cai A, et al. Aged landfill leachate enhances anaerobic digestion of waste activated sludge. Journal of Environmental Management. 2021; 293. <https://doi.org/10.1016/j.jenvman.2021.112853>
- Kurniawan TA, Haider A, Mohyuddin A, Fatima R, Salman M, et al. Tackling microplastics pollution in global environment through integration of applied technology, policy instruments and legislation. Journal of Environmental Management. 2023; 346. <https://doi.org/10.1016/j.jenvman.2023.118971>
- Xu Y, Seshadri B, Bolan N, Sarkar B, Ok YS, et al. Microbial functional diversity and carbon use feedback in soils as affected by heavy metals. Environment International. 2019; 125:478-88.

- <https://doi.org/10.1016/j.envint.2019.01.071>
25. Sunil B, Shrihari S, Nayak S. Shear strength characteristics and chemical characteristics of leachate-contaminated lateritic soil. *Engineering Geology*. 2009; 106(1-2):20-25. <https://doi.org/10.1016/j.enggeo.2008.12.011>
 26. Giri RK, Reddy KR. Slope stability of bioreactor landfills during leachate injection: effects of heterogeneous and anisotropic municipal solid waste conditions. *Waste Management & Research*. 2014; 32(3):186-97. <https://doi.org/10.1177/0734242X14522492>
 27. Francisca FM, Glatstein DA. Long term hydraulic conductivity of compacted soils permeated with landfill leachate. *Applied Clay Science*. 2010; 49(3):187-93. <https://doi.org/10.1016/j.clay.2010.05.003>
 28. Lu H, Xu S, Li D, Li J. An experimental study of mineral and microstructure for undisturbed loess polluted by landfill leachate. *KSCE Journal of Civil Engineering*. 2018; 22:4891-900. <https://doi.org/10.1007/s12205-017-1799-8>
 29. Jayasekera S, Mohajerani A. A study of the effects of municipal landfill leachate on a basaltic clay soil. *Australian Geomechanics*. 2001; 36(4):63-73.
 30. Xie Y, Wang H, Guo Y, Wang C, et al. Effects of biochar-amended soils as intermediate covers on the physical, mechanical and biochemical behaviour of municipal solid wastes. *Waste Management*. 2023; 171:512-21. <https://doi.org/10.1016/j.wasman.2023.10.004>
 31. Nayak S, Sunil B, Shrihari S. Hydraulic and compaction characteristics of leachate-contaminated lateritic soil. *Engineering Geology*. 2007; 94(3-4):137-44. <https://doi.org/10.1016/j.enggeo.2007.05.002>
 32. Ezema NM, Oduma CC, Nwaiwu CM, Mezie EO. Effects of leachate on geotechnical properties of lateritic soil. *UNIZIK Journal of Engineering and Applied Sciences*. 2022; 21(1):859-65.
 33. Harun N, Rahman AZ, Rahim AS, Lihan T, Idris RMW. Effects of leachate on geotechnical characteristics of sandy clay soil. *AIP Conference Proceedings*. 2013; American Institute of Physics. <https://doi.org/10.1063/1.4858709>
 34. Zheng M, Li S, Dong Q, Huang X, Liu Y. Effect of blending landfill leachate with activated sludge on the domestic wastewater treatment process. *Environmental Science: Water Research & Technology*. 2019; 5(2):268-76. <https://doi.org/10.1039/C8EW00799C>
 35. Özçoban MŞ, Acarer S, Tüfekci N. Effect of solid waste landfill leachate contaminants on hydraulic conductivity of landfill liners. *Water Science and Technology*. 2022; 85(5):1581-99. <https://doi.org/10.2166/wst.2022.033>
 36. Xie Y, Xue J, Gnanendran C. Effect of landfill leachate on hydraulic properties of an organic soil. In: *Proceedings of the 7th International Young Geotechnical Engineers Conference*, Sydney, Australia. 2022.
 37. Soon NW, Lee LM, Khun TC, Ling HS. Factors affecting improvement in engineering properties of residual soil through microbial-induced calcite precipitation. *Journal of Geotechnical and Geoenvironmental Engineering*. 2014; 140(5):04014006. [https://doi.org/10.1061/\(ASCE\)GT.1943-5606.0001089](https://doi.org/10.1061/(ASCE)GT.1943-5606.0001089)
 38. Tang S, She D, Wang H. Effect of salinity on soil structure and soil hydraulic characteristics. *Canadian Journal of Soil Science*. 2020; 101(1):62-73. <https://doi.org/10.1139/cjss-2020-0018>
 39. Juwonlo Osinubi K, Gadzama E, AdrianOshioname E. Unsaturated draulic conductivity of compacted bio cemented.
 40. Brovelli A, Malaguerra F, Barry DA. Bioclogging in porous media: Model development and sensitivity to initial conditions. *Environmental Modelling & Software*. 2009; 24(5):611-26. <https://doi.org/10.1016/j.envsoft.2008.10.001>
 41. Glatstein DA, Francisca FM. Hydraulic conductivity of compacted soils controlled by microbial activity. *Environmental Technology*. 2014; 35(15):1886-92. <https://doi.org/10.1080/09593330.2014.885583>
 42. Suganya A, Saravanan A, Manivannan N. Role of zinc nutrition for increasing zinc availability, uptake, yield and quality of maize (*Zea mays* L.) grains: An overview. *Communications in Soil Science and Plant Analysis*. 2020; 51(15):2001-21. <https://doi.org/10.1080/00103624.2020.1820030>
 43. Rahman R, Sofi JA, Javeed I, Malik TH, Nisar S. Role of micronutrients in crop production. *International Journal of Current Microbiology and Applied Sciences*. 2020; 8:2265-87.
 44. Rao GSPPP. Impact of leachate on soil properties in the dumpsite.
 45. Samadder SR, Prabhakar R, Khan D, Kishan D, Chauhan MS. Analysis of the contaminants released from municipal solid waste landfill site: a case study. *Science of the Total Environment*. 2017; 580:593-01. <https://doi.org/10.1016/j.scitotenv.2016.12.003>
 46. Agbeshie AA, Adjei R, Anokye J, Banunle A. Municipal waste dumpsite: Impact on soil properties and heavy metal concentrations, Sunyani, Ghana. *Scientific African*. 2020; 8. <https://doi.org/10.1016/j.sciaf.2020.e00390>
 47. Irvanian A, Ravari SO. Types of contamination in landfills and effects on the environment: a review study. In: *IOP Conference Series: Earth and Environmental Science*. 2020; IOP Publishing. <https://doi.org/10.1088/1755-1315/614/1/012083>
 48. Letsoalo ML. The effect of biochar on immobilization and phytoavailability of chromium, nickel and lead in soils amended with slag. 2020.
 49. Carbonell AA, Aarabi MA, DeLaune RD, Gambrell RP, Patrick Jr WH. Arsenic in wetland vegetation: availability, phytotoxicity, uptake and effects on plant growth and nutrition. *Science of the Total Environment*. 1998; 217(3):189-99. [https://doi.org/10.1016/S0048-9697\(98\)00195-8](https://doi.org/10.1016/S0048-9697(98)00195-8)
 50. Bouaouda S, Souabi S, Bouyakhssas R, Taleb A, et al. Techniques for treating leachate discharges: A critical review. *Euro-Mediterranean Journal for Environmental Integration*. 2023; 8(3):573-99. <https://doi.org/10.1007/s41207-023-00366-2>
 51. Wydro U, Wolejko E, Sokołowska G, Leszczyński J, Jabłońska-Trypuć A. Investigating Landfill Leachate Influence on Soil Microbial Biodiversity and Its Cytotoxicity. *Water*. 2022; 14(22):3634. <https://doi.org/10.3390/w14223634>
 52. Teng C, Zhou K, Peng C, Chen W. Characterization and treatment of landfill leachate: A review. *Water Research*. 2021; 203:117525. <https://doi.org/10.1016/j.watres.2021.117525>
 53. Sahu SK, Behuria HG. Heavy Metal Toxicity and Their Bioremediation Using Microbes, Plants, and Nano biomaterials. In: *Bioresource Utilization and Management*. Apple Academic Press; 2021; 457-70. <https://doi.org/10.1201/9781003057826-24>
 54. Tabatabai M. Soil enzymes. *Methods of soil analysis: Part 2 Microbiological and biochemical properties*. 1994; 5:775-33. <https://doi.org/10.2136/sssabookser5.2.c37>
 55. Daniel AN, Ekeleme IK, Onuigbo CM, Ikpeazu VO, Obiekezie SO. Review on effect of dumpsite leachate to the environmental and public health implication. *GSC Advanced Research and Reviews*. 2021; 7(2):051-060. <https://doi.org/10.30574/gscarr.2021.7.2.0097>
 56. Torkashvand J, Rezaei Kalantary R, Heidari N, Kazemi Z, et al. Application of ultrasound irradiation in landfill leachate treatment. *Environmental Science and Pollution Research*. 2021; 28:47741-751. <https://doi.org/10.1007/s11356-021-15280-9>
 57. Adelopo AO, Haris PI, Alo BI, Huddersman K, Jenkins RO. Multivariate analysis of the effects of age, particle size and

- landfill depth on heavy metals pollution content of closed and active landfill precursors. *Waste Management*. 2018; 78:227-37. <https://doi.org/10.1016/j.wasman.2018.05.040>
58. Gworek B, Dmuchowski W, Koda E, Marecka M, Baczevska AH. Impact of the municipal solid waste Łubna Landfill on environmental pollution by heavy metals. *Water*. 2016; 8(10):470. <https://doi.org/10.3390/w8100470>
 59. Adamcová D, Radziemska M, Ridošková A, Bartoň S, et al. Environmental assessment of the effects of a municipal landfill on the content and distribution of heavy metals in *Tanacetum vulgare* L. *Chemosphere*. 2017; 185:1011-18. <https://doi.org/10.1016/j.chemosphere.2017.07.060>
 60. Jayanthi S, Eswar NK, Singh SA, Chatterjee K, et al. Macroporous three-dimensional graphene oxide foams for dye adsorption and antibacterial applications. *RSC Advances*. 2016; 6(2):1231-42. <https://doi.org/10.1039/C5RA19925E>
 61. Rahman Z, Singh VP. The relative impact of toxic heavy metals (THMs)(arsenic (As), cadmium (Cd), chromium (Cr) (VI), mercury (Hg) and lead (Pb)) on the total environment: an overview. *Environmental Monitoring and Assessment*. 2019; 191:1-21. <https://doi.org/10.1007/s10661-019-7528-7>
 62. Pasalari H, Farzadkia M, Gholami M, Emamjomeh MM. Management of landfill leachate in Iran: valorization, characteristics and environmental approaches. *Environmental Chemistry Letters*. 2019; 17:335-48. <https://doi.org/10.1007/s10311-018-0804-x>
 63. Beinabaj SM, Heydariyan H, Aleii HM, Hosseinzadeh A. Concentration of heavy metals in leachate, soil and plants in Tehran's landfill: Investigation of the effect of landfill age on the intensity of pollution. *Heliyon*. 2023; 9(1). <https://doi.org/10.1016/j.heliyon.2023.e13017>
 64. Johar P, Singh D, Kumar A. Spatial variations of heavy metal contamination and associated risks around an unplanned landfill site in India. *Environmental Monitoring and Assessment*. 2020; 192:1-14. <https://doi.org/10.1007/s10661-020-08315-0>
 65. Li H, Luo N, Li YW, Cai QY, et al. Cadmium in rice: transport mechanisms, influencing factors, and minimizing measures. *Environmental Pollution*. 2017; 224:622-30. <https://doi.org/10.1016/j.envpol.2017.01.087>
 66. Panthee S. Possible methods of preventing groundwater contamination at landfill sites; case studies from Nepal. *Bulletin of the Department of Geology*. 2008; 11:51-60. <https://doi.org/10.3126/bdg.v11i0.1542>
 67. Kumar R, Kumari S. Geotechnical properties of materials used in landfill clay liner: A critical review. *Sādhanā*. 2023; 48(2):64. <https://doi.org/10.1007/s12046-023-02124-0>
 68. Nikbakht M, Behrooz Sarand F, Dabiri R, Bonab MH. Application of nanoclay and nanofiber filters to reduce soil permeability and leachates from landfill liners: A Review. *Geotechnical Geology*. 2022; 18(1):671-80.
 69. Wang B, Xu J, Chen B, Dong X, Dou T. Hydraulic conductivity of geosynthetic clay liners to inorganic waste leachate. *Applied Clay Science*. 2019; 168:244-48. <https://doi.org/10.1016/j.clay.2018.11.021>
 70. Ajitha A, Chandrakaran S, Evangeline YS. Impact of lead on the geotechnical properties and adsorption characteristics of landfill liner. *International Journal of Applied Engineering Research*. 2019; 14(13):3104-10.
 71. Yu F, Wu Z, Wang J, Li Y, et al. Effect of landfill age on the physical and chemical characteristics of waste plastics/microplastics in a waste landfill site. *Environmental Pollution*. 2022; 306. <https://doi.org/10.1016/j.envpol.2022.119366>
 72. Ban R, Chen X, Song Y, Bi P, et al. Study on permeability and electrical resistivity of red clay contaminated By Cu²⁺. *Stavební obzor-Civil Engineering Journal*. 2021; 30(1). <https://doi.org/10.14311/CEJ.2021.01.0023>
 73. Dutta J, Mishra AK. Influence of the presence of heavy metals on the behaviour of bentonites. *Environmental Earth Sciences*. 2016; 75:1-10. <https://doi.org/10.1007/s12665-016-5811-2>