



RESEARCH ARTICLE

# Effect of different packaging methods and storage conditions on postharvest quality and shelf-life of Manila Tamarind (*Pithecellobium dulce* (Roxb.) Benth.) var. PKM2

K. Udhayakumar<sup>1</sup>, S. Muthulakshmi<sup>1\*</sup>, A. Beulah<sup>2</sup>, A. Vijayasamundeeswari<sup>1</sup>, T. Anitha<sup>2</sup>, S. Saraswathy<sup>1</sup>, K.R. Rajadurai<sup>3</sup>, C. Rajamanickam<sup>1</sup> & C. Sankar<sup>1</sup>

<sup>1</sup>Department of Fruit Science, Horticultural College and Research Institute, Periyakulam, Theni, Tamil Nadu Agricultural University- 625 604, Tamil Nadu, India

<sup>2</sup>Department of Postharvest Technology, Horticultural College and Research Institute, Periyakulam, Theni, Tamil Nadu Agricultural University - 625 604, Tamil Nadu, India

<sup>3</sup>Department of Floriculture and Landscape Architecture, Horticultural College and Research Institute, Periyakulam, Theni, Tamil Nadu Agricultural University - 625 604, Tamil Nadu, India

\*Email: [muthulakshmi.s@tnau.ac.in](mailto:muthulakshmi.s@tnau.ac.in)

 OPEN ACCESS

## ARTICLE HISTORY

Received: 30 August 2024  
Accepted: 11 September 2024  
Available online  
Version 1.0 : 04 November 2024



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc  
See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Udhayakumar K, Muthulakshmi S, Beulah A, Vijayasamundeeswari A, Anitha T, Saraswathy S, Rajadurai KR, Rajamanickam C, Sankar C. Effect of different packaging methods and storage conditions on postharvest quality and shelf-life of Manila Tamarind (*Pithecellobium dulce* (Roxb.) Benth.) var. PKM2. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.4880>

## Abstract

Manila tamarind (*Pithecellobium dulce* (Roxb.) Benth.), is a dry land fruit crop that belongs to the Fabaceae family commonly known Camacchile or Jungle jilebi. *Pithecellobium dulce* (Roxb.) has the potential to become productive under harsh climatic and edaphic conditions of dry land areas but its pods have a shorter shelf life. The most important objective was to use proper packaging methods to reduce moisture loss, slow down physiological and biochemical changes and prevent spoilage. The study evaluated 8 packaging methods, including different combinations of CFB (Corrugated Fiberboard) lined with 100-gauge polythene film with 0 %, 5 % and 10 % vents, CFB with 0 %, 5 % and 10 % vents, vacuum packing and a control, under cold storage (13-15 °C) and ambient conditions (26 ± 2 °C). On the 15<sup>th</sup> day of cold storage conditions, vacuum packing showed minimum physiological loss of weight (19.82 %), titratable acidity (0.68 %), ascorbic acid (77.20) and total sugar (8.36 %). Maximum total soluble solids (15.32 °Brix), Anthocyanin (22.72 mg 100 g<sup>-1</sup>m), shelf life (15 days) followed by CFB with 10 % vent. On the 12<sup>th</sup> day of ambient storage condition, vacuum packing recorded minimum physiological loss of weight (16.42 %), titratable acidity (0.42 %), ascorbic acid (76.13) and total sugar (8.53 %). Maximum total soluble solids (14.93 °Brix), Anthocyanin (21.83 mg 100 g<sup>-1</sup>m) and shelf life (11.5 days) followed by CFB with 10 % vent. In this study, the overall results indicated that both vacuum packaging and CFB with 10 % vent at cold storage had synergistic effect in manila tamarind fruit pods, not only in extending the shelf life but also maintained the physiological and biochemical attributes of manila tamarind.

## Keywords

cold storage; Corrugated fibre board (CFB) packaging; Manila Tamarind; shelf life; vacuum packaging

## Introduction

Manila tamarind is one of the major underutilized fruit crops which may be an important fruit for the future due to its high medicinal value, high production per unit area and suitable for wasteland cultivation. Being the second most populous country in the world (1), India holds responsibility of

feeding its 1.3 billion people. However, with increasing population pressure and improving standards of living, human survivorship has become one of the most serious challenges for mankind. There is a potential threat of food and nutritional scarcity due to the apathetic exploitation of resources, leading to their depletion. This emerges the need for alternatives, thus exploring the unexplored and underutilised plant species. *Pithecellobium dulce* (Roxb.) Benth. belongs to the family Fabaceae, native to Mexico, South America and Central America. It is common in India, Malaysia and Thailand. Often planted for living fence or thorny hedge, eventually nearly impenetrable, camachile furnishes food, forage and firewood, while fixing a little nitrogen. In India it is mostly found in states of Tamil Nadu, Andhra Pradesh, Rajasthan, West Bengal and Delhi. The tree bears coiled fruits from end of the April till June. The fruit is a pod having sweet and slightly astringent white and red arils depending upon the tree covering black seeds. The fruit pulp is taken orally to stop blood flow in case of haemoptysis, seeds when grounds are used to cleanse ulcers and also numerous studies have been performed on anti-oxidant, anti-inflammatory, anti-diabetic and anti-cancerous properties (2).

Inadequate postharvest management accounts for 30 to 40 % loss of manila tamarind pods, which are highly perishable in nature. Pods can be stored for 2-3 days under normal conditions. Postharvest management such as packaging and storage technologies are crucial for extending shelf life and retention of nutrients. Fruit pods can be stored at low temperature to slowdown the ripening process. Packaging is also crucial in reducing the risk of pathogens being transported by fresh produce (3). The main objective was to keep the product well-packaged and stored to reduce moisture loss, slow down physical and chemical changes and prevent damage like bruising or spoilage during transport.

## Materials and Methods

### Collection of Manila Tamarind Fruits

The present experiment was conducted at Department of postharvest technology, Horticultural College and Research Institute, Periyakulam during the month of February to May 2024. Manila tamarind fruits were collected from 5-year-old orchard at Theni, Tamil Nadu, India. The fruits were greenish-brown to red-pinkish, indehiscent Pods. The pods were rather thin, 10-15 cm long and 1-2 cm wide. The spirally twisted fruits with clear constrictions, pale yellow pods and red attractive arils were found in clusters of 2-3. The Pods hold 10 seeds, with the fruit arils displaying a reddish colour. The seeds are flat, black and shiny. There are 48 kg of fruits are taken for this experiment.

## Methodology

Manila tamarind harvested at optimum maturity the skin of fruits turns yellow and seeds turn black



Harvested fruits were collected in perforated plastic trays



Within an hour after harvesting, fruits kept in a pre-cooling room at 15 °C to remove the field heat and keep the fruits as cool



After the pre-cooling fruits are packed different packaging treatments



Packed fruits were stored under ambient (26 ± 2 °C) and cold storage (13 ± 1 °C) conditions for further analysis



During experimentation, manila tamarind quality parameters were assessed once in 3 days

Laboratory research was conducted at Horticultural College and Research Institute, Periyakulam, Theni district, Tamil Nadu, India during March- June 2024 to evaluate the influence of CBF and vacuum packaging on storage of freshly harvested Manila Tamarind var. PKM2. The treatments include packing of Manila Tamarind var. PKM 2 pods in 8 different treatments of CFB lined with 100 gauge polythene film with 0 % vent, CFB lined with 100 gauge polythene film with 5 % vent, CFB lined with 100 gauge polythene film with 10 % vent, CFB with 0 % vent, CFB with 5 % vent, CFB with 10 % vent, vacuum packing and control. The 16 treatment combinations in the experiment were set up in a 2-factor Factorial Completely Randomized Block Design (FCRD) with 3 replications. The particulars of the treatment are provided in Table 1.

To study physiological and biochemical changes during storage of the fruits under cold and ambient, observations of the fruit samples were recorded at 3 days interval.

### Physiological loss in weight

The physiological loss in weight of fruits was assessed at 3 days interval and calculated using the formula given below and expressed in % (4).

$$PLW (\%) = \frac{\text{Initial weight (g)} - \text{Final weight (g)}}{\text{Initial weight (g)}} \times 100$$

### Total Soluble Solids (TSS)

TSS With the use of digital refractometer, an LED was allowed to pass light through a prism in contact with the fruit sample. An image sensor determines the critical angle at which the light is no longer refracted through the sample and the TSS of the fruits was calculated in terms of a degree brix.

**Table 1.** Storage conditions and packaging methods Manila Tamarind var. PKM 2.

Storage conditions and packaging methods Manila Tamarind var. PKM 2	
Factor 1 (Storage condition)	Factor 2 (Packaging methods)
S <sub>1</sub> - Cold storage (13-15 °C)	P <sub>1</sub> - CFB lined with 100 gauge polythene film with 0 % vent
S <sub>2</sub> - Ambient condition 26 + 2 °C)	P <sub>2</sub> - CFB lined with 100 gauge polythene film with 5 % vent
	P <sub>3</sub> - CFB lined with 100 gauge polythene film with 10 % vent
	P <sub>4</sub> - CFB with 0 % vent
	P <sub>5</sub> - CFB with 5 % vent
	P <sub>6</sub> - CFB with 10 % vent
	P <sub>7</sub> - Vacuum packing
	P <sub>8</sub> - Control

### Titrateable acidity (%)

Titrateable acidity was determined according to the method described (5). 50 g of manila tamarind flakes was homogenized using a blender. An amount of 10 g of blended flakes was mixed with 25 mL of distilled water and titrated against 0.1 N NaOH using Phenolphthalein indicator. Appearance of faint pink colour indicates the end point and the result was expressed as (mg 100 g<sup>-1</sup>) citric acid equivalent.

$$\text{Titrateable acidity (\%)} = \frac{T \times N \times E \times V_1}{V_2 \times W} \times 100$$

Where,

T=titre value, V<sub>1</sub>=volume made up, N= normality of NaOH, V<sub>2</sub>= volume of extract taken for estimation, E= equivalent weight of citric acid, W= weight of sample taken for estimation.

### Ascorbic acid (mg 100 g<sup>-1</sup>)

Ascorbic acid content was estimated according to method prescribed (6) and formula given below.

$$\text{Amount of ascorbic acid Present in 100 g of sample} = \frac{0.5 \text{ mg}}{V_1 \text{ mL}} \times \frac{V_2}{5 \text{ mL}} \times \frac{0.5 \text{ mg}}{V_1 \text{ mL}} \times 100$$

### Total sugars (%)

Total sugars were calculated using the anthrone method. It was computed using the following formula and expressed as a % (7).

$$\text{Total sugars (\%)} = \frac{\text{Sugar value from graph (mg)} \times \text{Total volume of extract (mL)}}{\text{Aliquot taken (mL)} \times \text{Weight of the sample (mg)}} \times 100$$

### Anthocyanin

Anthocyanin was extracted from the sample by blending 10 g of sample with 10 mL of ethanolic HCl and transferred to 100 mL volumetric flask. The volume was made up and the solution was stored in refrigerator at 4 °C and then filtered through Whatman No.1 filter paper. Optical density of filtrate was recorded at 535 nm (8).

$$\text{Total O.D/100 g} = \frac{\text{O.D X Volume made up}}{\text{Weight of sample}} \times 100$$

$$\text{Total anthocyanin (mg/ 100 g)} = \frac{\text{Total O.D/100 g}}{98.2}$$

### Shelf life (days)

The shelf-life of berries in each treatment was determined by recording number of days, where fruit pods retain their appearance and fit for consumption, without any decay. The shelf-life of fruits is expressed in days.

### Statistical analysis:

The statistical analysis was performed using AGRES software version SVN-1FN. Factorial Completely Randomized Design (FCRD) was used to understand the main effects of with and without ventilation, storage conditions and their interaction effects for the various parameters observed in the packed manila tamarind fruits. The mean comparisons was made after computing analysis of variance (ANOVA), standard deviation [SE (d)] and least significant differences (LSD) with p = 0.05 level of significance (9).

## Results

### Physiological loss in weight (% PLW)

During the storage period, the mean physiological loss in weight gradually increased and ranged from 3.69 to 19.82 % (3<sup>th</sup> day to 15<sup>th</sup> day). Among the various packaging methods, minimum mean PLW of 2.85 % was recorded in P<sub>7</sub> (vacuum packaging) followed by 3.15 % in P<sub>6</sub> whereas, a maximum mean PLW of 4.27 % was noted in P<sub>8</sub> on 3<sup>rd</sup> day of storage. At the end of the storage period (12<sup>th</sup> day), P<sub>7</sub> recorded minimum PLW of 14.99 % whereas, a maximum PLW of 15.52 % was recorded in P<sub>8</sub> (control). Among the storage conditions, manila tamarind fruit pods stored at cold storage (S<sub>1</sub>) recorded minimum mean values of PLW (3.10 %, 6.58 %, 10.37 % and 14.54 %) whereas, maximum mean values of PLW (4.28 %, 7.50 %, 10.45 % and 16.42 %) was recorded in fruits stored at ambient storage (S<sub>2</sub>) on 3<sup>rd</sup>, 3<sup>th</sup>, 6<sup>th</sup>, 9<sup>th</sup> and 12<sup>th</sup> day of storage respectively. The interaction between packaging and storage conditions inferred that the minimum PLW (2.21 %, 4.98 %) was observed in P<sub>7</sub>S<sub>1</sub> (vacuum packaging at cold storage) followed by (2.59 %, 5.36 %) in P<sub>6</sub>S<sub>1</sub> (CFB with 10 % vent at cold storage) on 3<sup>rd</sup> and 6<sup>th</sup> day of storage. Whereas, maximum PLW (4.83 %) was observed in P<sub>8</sub>S<sub>2</sub> (control at ambient storage) on 3<sup>rd</sup> day of storage (Table 2).

**Table 2.** Effect of packaging methods and storage conditions on physiological weight loss (%) of manila tamarind during storage.

Treatments	Storage (days)														
	3 Day			6 Day			9 Day			12 Day			15 Day		
	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean
P <sub>1</sub>	3.68	4.69	<b>4.18</b>	8.53	-	<b>8.53</b>	-	-	-	-	-	-	-	-	-
P <sub>2</sub>	3.38	4.57	<b>3.97</b>	7.28	-	<b>7.28</b>	-	-	-	-	-	-	-	-	-
P <sub>3</sub>	2.64	3.80	<b>3.22</b>	5.97	7.82	<b>6.89</b>	10.53	-	<b>10.53</b>	-	-	-	-	-	-
P <sub>4</sub>	3.52	4.62	<b>4.07</b>	7.81	-	<b>7.81</b>	-	-	-	-	-	-	-	-	-
P <sub>5</sub>	3.11	4.53	<b>3.82</b>	6.14	8.26	<b>7.20</b>	11.82	-	<b>11.82</b>	-	-	-	-	-	-
P <sub>6</sub>	2.59	3.71	<b>3.15</b>	5.36	7.13	<b>6.24</b>	9.85	-	<b>9.85</b>	15.52	-	<b>15.52</b>	-	-	-
P <sub>7</sub>	2.21	3.50	<b>2.85</b>	4.98	6.82	<b>5.90</b>	9.31	10.45	<b>9.88</b>	13.57	16.42	<b>14.99</b>	19.82	-	<b>19.82</b>
P <sub>8</sub>	3.72	4.83	<b>4.27</b>	-	-	-	-	-	-	-	-	-	-	-	-
Mean	<b>3.10</b>	<b>4.28</b>	<b>3.69</b>	<b>6.58</b>	<b>7.50</b>	<b>7.12</b>	<b>10.37</b>	<b>10.45</b>	<b>10.52</b>	<b>14.54</b>	<b>16.42</b>	<b>15.25</b>	<b>19.82</b>	-	<b>19.82</b>
Factors	S	P	S×P	S	P	S×P	S	P	S×P	S	P	S×P	S	P	S×P
SE (d)	0.062	0.123	0.175	0.082	0.164	0.232	0.08	0.161	0.227	0.081	0.162	0.229	0.099	0.198	0.28
C.D (p=0.05)	0.13**	0.26**	0.21*	0.175**	0.35**	0.49**	0.17**	0.34**	0.48**	0.17**	0.34**	0.48**	0.21**	0.42**	0.59**

(\* - significant at 5 % level; \*\* - significant at 1 % level; NS - Non Significant).

### Total Soluble Solids (TSS °Brix)

During storage, mean values of TSS increased from 13.55 °Brix to 14.44 °Brix on 3<sup>rd</sup> to 9<sup>th</sup> day. From 12<sup>th</sup> to 15<sup>th</sup> day, TSS decreased from 14.31 °Brix to 13.51 °Brix. Among the various packaging methods, minimum increase in TSS (12.45 °Brix and 12.40 °Brix) was recorded in P<sub>7</sub> vacuum followed by 12.50 °Brix, 13.50 °Brix and 13.79 °Brix in P<sub>6</sub> (CFB with 10 % vent) whereas, maximum increase in TSS 13.6 °Brix was noted in P<sub>8</sub> on 3<sup>rd</sup> day of storage. On ninth day of storage, all the packed fruit pods at ambient storage were decayed, except vacuum packaging and cold stored pods came up to 15 days P<sub>7</sub>. At end of the Storage 12<sup>th</sup> day, TSS started decreasing and it was minimum TSS content (14.2 °Brix) in P<sub>7</sub> followed by (14.42 °Brix) in P<sub>6</sub>. The fruits pods stored at cold storage (S<sub>1</sub>) recorded minimum increase in TSS (12.52 °Brix to 13.51 °Brix) from 3<sup>rd</sup> to 15<sup>th</sup>

day of storage whereas, maximum increase in TSS (14.58 °Brix to 14.93 °Brix) was observed in fruits stored at ambient storage (S<sub>2</sub>) from 3<sup>rd</sup> to 12<sup>th</sup> day of storage. Fruits stored in ambient condition were discarded after 12<sup>th</sup> days due to decay. The interaction effect on 3<sup>rd</sup> day but the lower TSS was observed (10.80 °Brix) in P<sub>7</sub>S<sub>1</sub> (vacuum packaging at cold storage) and the higher TSS was recorded in P<sub>8</sub>S<sub>2</sub> (control at ambient storage) is 15.70 °Brix. On 3<sup>rd</sup> day of storage, all the packed fruits pods at ambient storage were decayed on ninth day, except vacuum packaging fruits pods and cold stored fruits came up to 15 days (P<sub>7</sub> - vacuum packaging). At the end of storage period minimum TSS was observed (13.48 °Brix) in P<sub>7</sub>S<sub>1</sub> (vacuum packaging at cold storage) and the maximum TSS was recorded in P<sub>7</sub>S<sub>2</sub> (vacuum packaging at ambient storage) is 14.93 °Brix on 12<sup>th</sup> day of storage (Table 3).

**Table 3.** Effect of packaging methods and storage conditions on total soluble solids (°Brix) of manila tamarind during storage.

Treatments	Storage (days)														
	3 Day			6 Day			9 Day			12 Day			15 Day		
	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean
P <sub>1</sub>	12.69	13.64	<b>13.16</b>	14.84	-	<b>14.84</b>	-	-	-	-	-	-	-	-	-
P <sub>2</sub>	13.40	14.67	<b>14.03</b>	14.20	-	<b>14.2</b>	-	-	-	-	-	-	-	-	-
P <sub>3</sub>	11.90	14.70	<b>13.3</b>	13.20	15.80	<b>14.5</b>	14.78	-	<b>14.78</b>	-	-	-	-	-	-
P <sub>4</sub>	13.20	15.30	<b>14.25</b>	14.50	-	<b>14.5</b>	-	-	-	-	-	-	-	-	-
P <sub>5</sub>	12.70	14.80	<b>13.75</b>	13.90	15.90	<b>14.9</b>	15.08	-	<b>15.08</b>	-	-	-	-	-	-
P <sub>6</sub>	11.20	13.80	<b>12.5</b>	12.30	14.70	<b>13.5</b>	13.79	-	<b>13.79</b>	14.42	-	<b>14.42</b>	-	-	-
P <sub>7</sub>	10.80	14.10	<b>12.45</b>	11.70	15.10	<b>13.4</b>	13.45	14.79	<b>14.12</b>	13.48	14.93	<b>14.20</b>	13.51	-	<b>13.51</b>
P <sub>8</sub>	14.32	15.70	<b>15.01</b>	-	-	-	-	-	-	-	-	-	-	-	-
Mean	<b>12.52</b>	<b>14.58</b>	<b>13.55</b>	<b>13.52</b>	<b>15.37</b>	<b>14.26</b>	<b>14.27</b>	<b>14.79</b>	<b>14.44</b>	<b>13.95</b>	<b>14.93</b>	<b>14.31</b>	<b>13.51</b>	-	<b>13.51</b>
Factors	S	P	S×P	S	P	S×P	S	P	S×P	S	P	S×P	S	P	S×P
SE (d)	0.221	0.442	0.625	0.172	0.344	0.486	0.111	0.222	0.314	0.079	0.157	0.223	0.068	0.135	0.191
C.D (p=0.05)	0.472*	0.945*	NS	0.367*	0.735*	1.039*	0.237*	0.475*	0.671*	0.168*	0.336*	0.476*	0.144*	0.289*	0.409*

(\* - significant at 5 % level; \*\* - significant at 1 % level; NS - Non Significant).

### Titratable acidity (%)

The maximum acidity (1.87 %) was recorded in P<sub>7</sub> followed by 1.66 % in P<sub>6</sub>, minimum acidity (1.45 %) was noted in P<sub>8</sub> (Control) on 3<sup>rd</sup> day of storage. On ninth day of storage, all the manila tamarind fruits pods at ambient storage were decayed, except vacuum packaging manila tamarind pods and cold stored fruits came up to 15 days P<sub>7</sub>. The mean acidity of 1.01 % was recorded in vacuum packaging (P<sub>7</sub>) on ninth day. On the 12<sup>th</sup> and 15<sup>th</sup> day of storage, maximum acidity (1.05 % and 0.68 %) was noted in P<sub>7</sub> followed by 3.45 % in P<sub>6</sub> (CFB with 10 % vent), at 12<sup>th</sup> day of storage. Among the various storage conditions, cold storage (S<sub>1</sub>) recorded the higher values of acidity (4.64 %, 4.17 %, 4.08 %, 3.75 % and 3.68 %) on 3<sup>rd</sup>, 6<sup>th</sup>, 9<sup>th</sup>, 12<sup>th</sup> and 15<sup>th</sup> day of storage whereas, lower values of acidity (4.46 %, 3.88 % and 3.76 %) were observed in manila tamarind fruits pods stored at ambient storage (S<sub>2</sub>) on 3<sup>rd</sup>, 6<sup>th</sup>, 9<sup>th</sup> and 12<sup>th</sup> day of storage, respectively. Manila tamarind fruit pods stored at ambient storage (S<sub>1</sub>) were discarded after 12<sup>th</sup> days of storage due to decay. On Day 3, there is no significant difference between packaging and storage conditions. The results showed that the acidity was higher (4.91 % and 4.65 %) in P<sub>7</sub>S<sub>1</sub> (vacuum packaging at cold storage) Followed by (4.70 % and 4.56 %) in P<sub>6</sub>S<sub>1</sub> (CFB with 10 % vent at cold storage) on 3<sup>rd</sup> and 6<sup>th</sup> day of storage respectively, whereas, acidity was lower in P<sub>8</sub>S<sub>2</sub> (control at ambient storage) with 4.39 % on 3<sup>rd</sup> day of storage. At the end of 12<sup>th</sup> day of storage, lowest acidity (3.45 %) was recorded in P<sub>6</sub>S<sub>1</sub> (CFB with 10 % vent at cold storage) and highest acidity (4.05 %) was recorded in T<sub>6</sub>S<sub>2</sub> (vacuum packaging at cold storage) (Table 4).

### Ascorbic acid (mg 100 g<sup>-1</sup>)

The mean value of ascorbic acid content decreased from 83.09 mg 100 g<sup>-1</sup> on 3<sup>rd</sup> day to 77.20 mg 100 g<sup>-1</sup> on 15<sup>th</sup> day of storage. Results showed that the maximum ascorbic acid content (86.20 mg 100 g<sup>-1</sup>) was recorded in P<sub>7</sub> which was on

par with 85.65 mg 100 g<sup>-1</sup> in P<sub>6</sub> followed by 85.30 mg 100 g<sup>-1</sup> in P<sub>3</sub> (CFB lined with 100 gauge polythene film with 10 % vent) whereas, minimum ascorbic acid content (78.58 mg 100 g<sup>-1</sup>) was noted in P<sub>8</sub> on 3<sup>rd</sup> day of storage. On ninth day of storage, all the packed fruits pods at ambient storage were decayed, except vacuum packed fruit pods and cold stored fruits came up to 15<sup>th</sup> days (vacuum packaging). The mean ascorbic acid of 80.32 mg 100 g<sup>-1</sup> was recorded in vacuum packaging (P<sub>7</sub>) on ninth day. At the end of storage period, maximum ascorbic acid content (77.20 mg 100 g<sup>-1</sup>) was recorded in P<sub>7</sub>. Among the various storage conditions, manila tamarind fruit pods stored at cold storage (S<sub>1</sub>) recorded maximum ascorbic acid content (84.79 to 77.20 mg 100 g<sup>-1</sup>) on 3<sup>rd</sup> to 15<sup>th</sup> day of storage whereas, minimum ascorbic acid content (81.38 to 76.13 mg 100 g<sup>-1</sup>) was recorded in fruits stored at ambient storage (S<sub>2</sub>) on 3<sup>rd</sup> to 12<sup>th</sup> day of storage. Fruits stored at ambient storage were discarded after 12<sup>th</sup> days of storage due to decay. The interaction between edible coatings and storage conditions inferred that the higher values of ascorbic acid content (88.10 and 85.25 mg 100 g<sup>-1</sup>) was registered in P<sub>7</sub>S<sub>1</sub> (vacuum packaging at cold storage) followed by 87.90 and 82.90 mg 100 g<sup>-1</sup> in P<sub>6</sub>S<sub>1</sub> (CFB with 10% vent at cold storage) on 3<sup>rd</sup> and 6<sup>th</sup> day of storage, respectively. Whereas, lower values were noted in P<sub>8</sub>S<sub>2</sub> (control at ambient storage) with 77.56 mg 100 g<sup>-1</sup> on 3<sup>rd</sup> day of storage (Table 5).

### Total Sugar (%)

The mean value of total sugar content starts increased up to ninth day (8.68 to 8.71 %) and then starts decreasing from 12<sup>th</sup> to 15<sup>th</sup> day (8.47 to 8.36 %). Results showed that the minimum total sugar content of 8.22 % was recorded in P<sub>7</sub> followed by 8.55 % in P<sub>6</sub> whereas, maximum total sugar content of 8.91 % was noted in P<sub>8</sub> on 3<sup>rd</sup> day of storage. On ninth day of storage, all the packed fruits pods at ambient storage were decayed, except vacuum packaging fruits pods and cold stored fruits came up to 15

**Table 4.** Effect of packaging methods and storage conditions on titratable acidity (%) of manila tamarind during storage.

Treat-ments	Storage (days)														
	3 Day			6 day			9 day			12 day			15 Day		
	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean
P <sub>1</sub>	1.56	1.39	<b>1.47</b>	0.88	-	<b>0.88</b>	-	-	-	-	-	-	-	-	-
P <sub>2</sub>	1.46	1.27	<b>1.36</b>	0.92	-	<b>0.92</b>	-	-	-	-	-	-	-	-	-
P <sub>3</sub>	1.75	1.34	<b>1.54</b>	1.13	0.45	<b>0.79</b>	0.74	-	<b>0.74</b>	-	-	-	-	-	-
P <sub>4</sub>	1.59	1.34	<b>1.46</b>	1.91	-	<b>0.91</b>	-	-	-	-	-	-	-	-	-
P <sub>5</sub>	1.65	1.56	<b>1.60</b>	1.15	0.83	<b>0.99</b>	0.13	-	<b>0.13</b>	-	-	-	-	-	-
P <sub>6</sub>	1.70	1.63	<b>1.66</b>	1.56	1.02	<b>1.29</b>	1.21	-	<b>1.21</b>	0.45	-	<b>0.45</b>	-	-	-
P <sub>7</sub>	1.91	1.83	<b>1.87</b>	1.65	1.24	<b>1.44</b>	1.26	0.76	<b>1.01</b>	1.05	0.42	<b>0.73</b>	0.68	-	<b>0.68</b>
P <sub>8</sub>	1.52	1.39	<b>1.45</b>	-	-	-	-	-	-	-	-	-	-	-	-
<b>Mean</b>	<b>1.64</b>	<b>1.46</b>	<b>1.55</b>	<b>1.17</b>	<b>0.88</b>	<b>1.03</b>	<b>1.08</b>	<b>0.76</b>	<b>1.02</b>	<b>0.75</b>	<b>0.42</b>	<b>0.59</b>	<b>0.68</b>	-	<b>0.68</b>
<b>Factors</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>
<b>SE (d)</b>	0.071	0.143	0.202	0.049	0.099	0.14	0.032	0.064	0.09	0.022	0.045	0.063	0.018	0.037	0.052
<b>C.D (p=0.05)</b>	0.15	0.28	NS	0.10	0.21	0.29	0.06	0.13	0.19	0.04	0.09	0.13	0.03	0.07	0.11**
	**	**		**	**	**	**	**	**	**	**	**	**	**	**

(\* - significant at 5 % level; \*\* - significant at 1 % level; NS - Non Significant).

**Table 5.** Effect of packaging methods and storage conditions on ascorbic acid (mg 100 g<sup>-1</sup>) of manila tamarind during storage.

Treatments	Storage (days)														
	3 day			6 day			9 day			12 day			15 day		
	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean
P <sub>1</sub>	82.70	80.01	<b>81.35</b>	77.21	-	<b>77.21</b>	-	-	-	-	-	-	-	-	-
P <sub>2</sub>	82.40	79.74	<b>81.07</b>	76.27	-	<b>76.27</b>	-	-	-	-	-	-	-	-	-
P <sub>3</sub>	87.80	82.80	<b>85.3</b>	82.93	77.43	<b>80.18</b>	78.98	-	<b>78.98</b>	-	-	-	-	-	-
P <sub>4</sub>	83.66	81.03	<b>82.34</b>	77.61	-	<b>77.61</b>	-	-	-	-	-	-	-	-	-
P <sub>5</sub>	86.20	82.24	<b>84.22</b>	80.97	76.42	<b>78.69</b>	77.78	-	<b>77.78</b>	-	-	-	-	-	-
P <sub>6</sub>	87.90	83.40	<b>85.65</b>	82.90	78.05	<b>80.47</b>	79.15	-	<b>79.15</b>	77.37	-	<b>77.37</b>	-	-	-
P <sub>7</sub>	88.10	84.30	<b>86.2</b>	85.25	82.87	<b>84.06</b>	80.32	78.86	<b>79.59</b>	79.57	76.13	<b>77.85</b>	77.20	-	<b>77.20</b>
P <sub>8</sub>	79.60	77.56	<b>78.58</b>	-	-	-	-	-	-	-	-	-	-	-	-
<b>Mean</b>	<b>84.79</b>	<b>81.38</b>	<b>83.09</b>	<b>80.44</b>	<b>78.69</b>	<b>79.21</b>	<b>79.05</b>	<b>78.86</b>	<b>78.87</b>	<b>78.47</b>	<b>76.13</b>	<b>77.61</b>	<b>77.20</b>	-	<b>77.20</b>
<b>Factors</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>
<b>SE (d)</b>	1.312	2.623	3.710	0.966	1.931	2.731	0.683	1.366	1.932	0.453	0.905	1.280	0.386	0.772	1.092
<b>C.D (p=0.05)</b>	2.80**	5.32**	6.13*	2.06**	4.12**	5.84*	1.46**	2.92**	4.13**	0.96**	1.93**	2.73**	0.82**	1.65**	2.334**

(\*- significant at 5 % level; \*\*-significant at 1 % level; NS- Non Significant).

days P<sub>7</sub>. The mean total sugars of 8.88 % were observed in vacuum packaging (P<sub>7</sub>) on ninth day. At the end of storage period, minimum total sugar content of 8.36 % was recorded in P<sub>7</sub>. Among the various storage conditions, manila tamarind fruit pods stored at cold storage (S<sub>2</sub>) recorded minimum total sugar content (8.20 % to 8.36 %) from 3<sup>rd</sup> to 15<sup>th</sup> day of storage whereas, maximum total sugar content (9.16 % to 9.26 %) was observed in fruits stored at ambient storage (S<sub>2</sub>) from 3<sup>rd</sup> to 6<sup>th</sup> day of storage. Fruits pods stored at ambient storage were discarded after fifteen days of storage due to decay. There is no significant difference between packaging and storage

conditions on 3<sup>rd</sup> day and it was inferred that the lower values of total sugar content (7.92 % and 8.26 %) was registered in P<sub>7</sub>S<sub>1</sub> (vacuum packaging at cold storage) followed by 8.06 and 8.45 % in P<sub>6</sub>S<sub>1</sub> (CFB with 10 % vent at cold storage) whereas, higher values were noted (9.21 %) in P<sub>7</sub>S<sub>2</sub> (control at ambient storage) on 3<sup>rd</sup> day of storage. During 12<sup>th</sup> day of storage, minimum total sugar content of 8.46 % was recorded in P<sub>7</sub>S<sub>1</sub> (vacuum packaging at cold storage) whereas, maximum total sugar content was recorded in P<sub>6</sub>S<sub>1</sub> - CFB with 10 % vent at cold storage with 8.49 % (Table 6).

**Table 6.** Effect of packaging methods and storage conditions on total sugar (%) of manila tamarind during storage.

Treatments	Storage (days)														
	3 day			6 day			9 day			12 day			15 Day		
	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean
P <sub>1</sub>	8.25	9.40	<b>8.82</b>	8.79	-	<b>8.79</b>	-	-	-	-	-	-	-	-	-
P <sub>2</sub>	8.47	9.50	<b>8.98</b>	8.82	-	<b>8.82</b>	-	-	-	-	-	-	-	-	-
P <sub>3</sub>	8.06	9.11	<b>8.58</b>	8.54	9.30	<b>8.92</b>	8.74	-	<b>8.74</b>	-	-	-	-	-	-
P <sub>4</sub>	8.14	9.35	<b>8.74</b>	8.78	-	<b>8.78</b>	-	-	-	-	-	-	-	-	-
P <sub>5</sub>	8.10	9.19	<b>8.64</b>	8.85	9.34	<b>9.09</b>	8.74	-	<b>8.74</b>	-	-	-	-	-	-
P <sub>6</sub>	8.06	9.05	<b>8.55</b>	8.45	9.27	<b>8.86</b>	8.51	-	<b>8.51</b>	8.49	-	<b>8.49</b>	-	-	-
P <sub>7</sub>	7.92	8.52	<b>8.22</b>	8.26	9.14	<b>8.7</b>	8.85	8.91	<b>8.88</b>	8.40	8.53	<b>8.46</b>	8.36	-	<b>8.36</b>
P <sub>8</sub>	8.61	9.21	<b>8.91</b>	-	-	-	-	-	-	-	-	-	-	-	-
<b>Mean</b>	<b>8.20</b>	<b>9.16</b>	<b>8.68</b>	<b>8.64</b>	<b>9.26</b>	<b>8.85</b>	<b>8.71</b>	<b>8.91</b>	<b>8.71</b>	<b>8.44</b>	<b>8.53</b>	<b>8.47</b>	<b>8.36</b>	-	<b>8.36</b>
<b>Factors</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>
<b>SE (d)</b>	0.139	0.278	0.394	0.106	0.212	0.299	0.146	0.293	0.414	0.047	0.094	0.133	0.042	0.084	0.118
<b>C.D (p=0.05)</b>	0.29**	0.54**	NS	0.22**	0.45**	0.64***	0.14**	0.29**	0.41**	0.10**	0.20**	0.28**	0.08**	0.17**	0.25**

(\*- significant at 5 % level; \*\*-significant at 1 % level; NS- Non Significant).

## Anthocyanin (mg 100 g<sup>-1</sup>)

The anthocyanin content of manila tamarind fruit pods was initially increased up to 12<sup>th</sup> day and decreased at the end of storage period. Results showed that the maximum anthocyanin content (18.23 mg 100 g<sup>-1</sup>) was recorded in P<sub>7</sub> followed by (18.00 mg 100 g<sup>-1</sup>) in P<sub>6</sub> whereas, minimum anthocyanin content (15.53 mg 100 g<sup>-1</sup>) was observed in P<sub>8</sub> on 3<sup>rd</sup> day of storage. On 9<sup>th</sup> day of storage, all the packed fruits pods at ambient storage were decayed, except vacuum packaging fruits pods cold stored fruits came up to 15 days P<sub>7</sub>. The mean total anthocyanin content of 21.67 mg 100 g<sup>-1</sup> was recorded in vacuum packaging (P<sub>7</sub>) on 9<sup>th</sup> day. At the end of storage period (12<sup>th</sup> day), total carotenoid content (23.25 mg 100 g<sup>-1</sup>) was higher in P<sub>7</sub> followed by 24.43 mg 100 g<sup>-1</sup> in P<sub>6</sub>. Among the various storage conditions, fruits stored at cold storage (S<sub>1</sub>) recorded the highest anthocyanin content (17.94 mg 100 g<sup>-1</sup>) whereas, lowest anthocyanin content (16.71 mg 100 g<sup>-1</sup>) was observed in fruits stored at ambient storage (S<sub>2</sub>) on 3<sup>rd</sup> day of storage. During 12<sup>th</sup> day of storage, fruits stored at cold storage (S<sub>1</sub>) recorded the anthocyanin content of about 24.55 mg 100 g<sup>-1</sup> whereas, fruits stored at ambient storage (S<sub>2</sub>) recorded the anthocyanin content of (21.83 mg 100 g<sup>-1</sup>) on 12<sup>th</sup> day of storage and the fruits were discarded after 12<sup>th</sup> day of storage due to decay. The interaction between packaging and storage conditions showed that the anthocyanin content was higher (18.91 mg 100 g<sup>-1</sup>) in P<sub>7</sub>S<sub>1</sub> (vacuum packaging at cold storage) followed by (18.76 mg 100 g<sup>-1</sup>) P<sub>6</sub>S<sub>1</sub> (CFB with 10 % vent at cold storage) whereas, lower (15.03 mg 100 g<sup>-1</sup>) in P<sub>8</sub>S<sub>2</sub> (control at ambient storage) on 3<sup>rd</sup> day of storage. At the end of storage period 12<sup>th</sup> day, highest anthocyanin content (24.67 mg 100 g<sup>-1</sup>) was recorded in P<sub>7</sub>S<sub>1</sub> (vacuum packaging at cold storage) followed by 24.43 mg 100 g<sup>-1</sup> in P<sub>6</sub>S<sub>1</sub> (CFB with 10 % vent at cold storage) (Table 7).

## Shelf life

Among the packaging, maximum shelf life 12.00 days was recorded in P<sub>7</sub> followed by 10.00 days in P<sub>6</sub>, 7.5 days in P<sub>3</sub> (CFB lined with 100-gauge polythene film with 10 % vent) and 6.75 days in P<sub>5</sub> (CFB with 5 % vent) whereas, minimum shelf life of 2.75 days was recorded in P<sub>8</sub>. Among the storage conditions, cold storage (S<sub>1</sub>) extended the shelf life of about 8.06 days whereas, fruits stored at ambient storage (S<sub>2</sub>) had the shelf life of about 4.87 days only. The interaction between packaging and storage conditions indicated that the maximum shelf life of 15.00 days was observed in P<sub>7</sub>S<sub>1</sub> (vacuum packaging at cold storage) (Fig. 1) followed by 11.5 days in P<sub>6</sub>S<sub>1</sub> (CFB with 10 % vent at cold storage), 9.0 days in P<sub>3</sub>S<sub>1</sub> (CFB lined with 100 gauge polythene film with 10 % vent at cold storage) and 8.5 days in P<sub>5</sub>S<sub>1</sub> (CFB with 5 % vent at cold storage) whereas, minimum shelf life of 2.5 days was observed in P<sub>8</sub>S<sub>2</sub> (control at ambient storage) on par with 2.5 days in P<sub>1</sub>S<sub>2</sub> (CFB lined with 100 gauge polythene film with 0 % vent at ambient storage) (Table 8).


## Microbial count (log CFU g<sup>-1</sup>)

One of the important factors in preservation and commercialization of fresh fruits is its shelf life and microbial activity (10). The manila tamarind fruits pods packed with vacuum packaging had shown bacteria, fungi, yeast and mould of 6.97, 5.03 and 4.07 log CFU g<sup>-1</sup> (ambient storage) and 1.02, 5.82 and 3.80 log CFU g<sup>-1</sup> (cold storage) whereas, the control fruits had a bacteria, fungi, yeast and mould of 6.97, 9.52 and 5.01 log CFU g<sup>-1</sup> (ambient storage) and 6.62, 9.02 and 4.95 log CFU g<sup>-1</sup> (cold storage) during 15<sup>th</sup> day of storage respectively (Fig. 2) (Table 9).


**Table 7.** Effect of packaging methods and storage conditions on anthocyanin (mg 100 g<sup>-1</sup>) of manila tamarind during storage.

Treatments	Storage (days)														
	3 day			6 day			9 day			12 day			15 day		
	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean
P <sub>1</sub>	17.64	16.65	<b>17.14</b>	20.18	-	<b>20.18</b>	-	-	-	-	-	-	-	-	-
P <sub>2</sub>	17.42	16.21	<b>16.81</b>	20.21	-	<b>20.21</b>	-	-	-	-	-	-	-	-	-
P <sub>3</sub>	18.31	17.04	<b>17.67</b>	21.56	19.24	<b>20.40</b>	21.97	-	<b>21.97</b>	-	-	-	-	-	-
P <sub>4</sub>	17.87	16.86	<b>17.36</b>	20.34	-	<b>20.34</b>	-	-	-	-	-	-	-	-	-
P <sub>5</sub>	18.64	17.14	<b>17.89</b>	21.21	19.58	<b>20.39</b>	22.23	-	<b>22.23</b>	-	-	-	-	-	-
P <sub>6</sub>	18.76	17.24	<b>18.00</b>	21.56	19.23	<b>20.39</b>	22.74	-	<b>22.74</b>	24.43	-	<b>24.43</b>	-	-	-
P <sub>7</sub>	18.91	17.56	<b>18.23</b>	21.67	19.58	<b>20.62</b>	22.94	20.40	<b>21.67</b>	24.67	21.83	<b>23.25</b>	22.72	-	<b>22.72</b>
P <sub>8</sub>	16.03	15.03	<b>15.53</b>	-	-	-	-	-	-	-	-	-	-	-	-
<b>Mean</b>	<b>17.94</b>	<b>16.71</b>	<b>17.33</b>	<b>20.96</b>	<b>19.40</b>	<b>20.36</b>	<b>22.47</b>	<b>20.40</b>	<b>22.15</b>	<b>24.55</b>	<b>21.83</b>	<b>23.84</b>	<b>22.72</b>	-	<b>22.72</b>
<b>Factors</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>
<b>SE (d)</b>	0.272	0.543	0.768	0.293	0.586	0.829	0.2	0.4	0.566	0.174	0.348	0.492	0.114	0.227	0.321
<b>C.D (p=0.05)</b>	0.58**	1.16**	1.67*	0.62**	1.25**	1.77**	0.42**	0.85**	1.21**	0.37**	0.74**	1.05**	0.24**	0.48**	0.68**

(\* - significant at 5 % level; \*\* - significant at 1 % level; NS - Non Significant).



**Fig. 1.** Effect of vacuum packaging under cold storage observed on 3<sup>rd</sup>, 6<sup>th</sup>, 9<sup>th</sup>, 12<sup>th</sup> and 15<sup>th</sup> days.



**Fig. 2.** Effect of packaging methods and storage conditions on microbial count observed in Manilla tamarind.



**Table 8.** Effect of packaging methods and storage conditions on shelf life (days) of manila tamarind during storage.

Treatments	Shelf life (days)		Mean
	S <sub>1</sub>	S <sub>2</sub>	
P <sub>1</sub>	5.50	2.50	<b>4.00</b>
P <sub>2</sub>	6.00	3.00	<b>4.50</b>
P <sub>3</sub>	9.00	6.00	<b>7.50</b>
P <sub>4</sub>	6.00	2.50	<b>4.25</b>
P <sub>5</sub>	8.50	5.00	<b>6.75</b>
P <sub>6</sub>	11.50	8.50	<b>10.00</b>
P <sub>7</sub>	15.00	11.5	<b>12.75</b>
P <sub>8</sub>	3.00	2.50	<b>2.75</b>
<b>Mean</b>	<b>8.06</b>	<b>4.87</b>	<b>6.46</b>
	<b>SE (d)</b>	<b>C.D (p=0.05)</b>	
<b>Storage (S)</b>	0.113	0.243**	
<b>Packaging (P)</b>	0.227	0.485**	
<b>S×P</b>	0.321	0.686*	

(\*- significant at 5 % level; \*\*-significant at 1 % level; NS- Non Significant).

## Discussion

### Physiological loss in weight (% PLW)

Loss of weight on prolonging the storage period mainly was attributed to the loss of moisture caused by evaporation and respiration (11) and may also be due to changes in peel stomatal density as well as texture (12). The PLW of manila tamarind occurs due to the reduction in moisture content. This slower rates of weight loss in vacuum packed fruits can be attributed to the barrier properties for gas diffusion of stomata (13), the organelles that regulate the transpiration process and gas exchange between the fruit and the environment (14). The less weight loss during cold storage was due to maintenance of the maximum moisture content around the surface of fruit along with storage having high humidity and cold storage conditions (15).

### Total Soluble Solids (TSS °Brix)

It is well documented that the filmogenic property of ventilated results in an excellent semi-permeable film around the fruit, modifying the internal atmosphere by reducing O<sub>2</sub> and/or elevating CO<sub>2</sub> and suppressing ethylene evolution. A suppressed respiration rate also

slows down the synthesis and the use of metabolites, resulting in lower TSS due to the slower hydrolysis of carbohydrates to sugar. The rate of increase in TSS was found to be faster in fruits stored at room temperature as compared to cold stored fruits (4).

### Titrateable acidity (%)

These results are corroborated with the findings decreased due to the apparent stability observed as an indicator of metabolic stability, because the same serves as a substrate for such reactions as respiration and production of volatiles during ripening (16, 17).

### Ascorbic acid (mg 100 g<sup>-1</sup>)

The loss in ascorbic acid content with the progress of the storage period could be attributed to the rapid conversion of L-ascorbic acid into dehydroascorbic acid in the presence of the L-ascorbic acid oxidase enzyme (18). Similarly, higher vitamin C content in ventilated packaging papaya fruit has also been reported previously by strawberry (19) and mango (20).

### Total Sugar (%)

Storage condition influenced the effectiveness of treatments on the total sugars content of manila tamarind fruits pods during storage. It was found to be higher in manila tamarind fruits pods stored at ambient temperature on the 9th day of storage, compared to the fruits stored in cold storage. According to a study, it might be due to the fact that high temperature and low relative humidity at room temperature resulted in conversion of starch and other insoluble carbohydrates in soluble sugars (4). The sugars were further utilized for respiration thus showing the lower content of sugars at the later period of storage. The low temperature in cold storage reduces fruit metabolism, particularly respiratory activity, delaying the ripening process and increasing fruit shelf life up to 2 weeks (21).

### Anthocyanin (mg 100 g<sup>-1</sup>)

In the present study, anthocyanin content was higher in vacuum packed fruits in cold storage conditions compared to fruits in ambient storage. The vacuum packaging was much more effective than other corrugated fibre board

**Table 9.** Effect of packaging methods and storage conditions on microbial count (log CFU g<sup>-1</sup>) of manila tamarind during storage.

Treatments	Storage (at 15 <sup>th</sup> day)								
	Bacteria			Fungi			Yeast and Mould		
	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean	S <sub>1</sub>	S <sub>2</sub>	Mean
P <sub>1</sub>	6.54	6.82	<b>6.68</b>	8.71	8.96	<b>8.83</b>	4.62	4.68	<b>4.65</b>
P <sub>2</sub>	5.37	5.48	<b>5.42</b>	8.42	8.57	<b>8.49</b>	4.32	4.46	<b>4.39</b>
P <sub>3</sub>	4.62	5.02	<b>4.82</b>	7.91	8.09	<b>8.00</b>	4.22	4.32	<b>4.27</b>
P <sub>4</sub>	6.29	6.37	<b>6.33</b>	8.62	8.68	<b>8.65</b>	4.52	4.61	<b>4.56</b>
P <sub>5</sub>	5.83	5.91	<b>5.87</b>	8.05	8.27	<b>8.16</b>	4.30	4.41	<b>4.35</b>
P <sub>6</sub>	3.42	4.87	<b>4.14</b>	6.21	6.92	<b>6.56</b>	4.19	4.28	<b>4.23</b>
P <sub>7</sub>	1.02	1.13	<b>1.07</b>	5.82	5.03	<b>5.42</b>	3.80	4.07	<b>3.93</b>
P <sub>8</sub>	6.62	6.97	<b>6.79</b>	9.02	9.52	<b>9.27</b>	4.95	5.01	<b>4.98</b>
<b>Mean</b>	<b>4.96</b>	<b>5.32</b>	<b>5.14</b>	<b>7.84</b>	<b>8.00</b>	<b>7.92</b>	<b>4.36</b>	<b>4.48</b>	<b>4.42</b>
<b>Factors</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>	<b>S</b>	<b>P</b>	<b>S×P</b>
<b>SE (d)</b>	0.087	0.174	0.245	0.128	0.256	0.363	0.070	0.139	0.197
<b>C.D (p=0.05)</b>	0.186**	0.371**	0.525**	0.247**	0.548**	0.724*	0.138**	0.297**	0.372*

(\*- significant at 5 % level; \*\*-significant at 1 % level; NS- Non Significant).

packaging in reducing the anthocyanin losses. Compared with control, the loss of total anthocyanin in vacuum packaging fruits pods occurred more slowly during storage. It's also observed concurrent biosynthesis of anthocyanin in the pulp along with the development of external purple colour (anthocyanin pigments) during ripening of manila tamarind fruit pods and the decline of anthocyanin content in manila tamarind fruit during storage period (22). The decrease in anthocyanin towards the later part of the storage period may be due to the degradation of anthocyanin pigments (23). Changes in anthocyanin profile due to storage temperature regimes during the storage of manila tamarind fruit pods. In general, the development of colour during postharvest storage of fruit might be due to degradation of chlorophyll content in fruit and increase synthesis of anthocyanin pigments (24).

### Shelf life

The shelf life of manila tamarind fruit pods is interconnected with broader physiological and biochemical parameters. Each parameter has its impact on fruit quality and storage life. This extension of the storage period was due to the delay in the ripening process caused by respiration, transpiration and antimicrobial properties of vacuum packaging which also delayed the respiration rate during storage. Vacuum packed fruit pods with cold storage helps improve the shelf-life and postharvest quality of bananas (25). Similar results have been reported in papaya (26), jujube (27), fig (28) and pomegranate (29).

### Microbial count (log CFU g<sup>-1</sup>)

In the present study, manila tamarind fruit pads packed with vacuum packaging were found to reduce the microbial growth under cold storage condition (Table 9). Compared to with and without ventilated corrugated fibre board packaging with ventilated packaging on the lid had lowered the microbial count during storage. When fruits stored under cold storage reduces the growth of many microorganisms (30). The phytosanitary measures adopted along with postharvest treatment and MAP ensures absence of Coliforms and Salmonella which were toxic to consumers. In addition to this, use of antimicrobials as postharvest treatment makes the fresh fruit produce to tolerate temperature fluctuations during storage (31). It was stated that the minimum microbial count in vacuum packed apple fruits might be due to its inhibitory effect on microbial growth by strengthening the cell wall compared to control which allows spore spreading, resulting in increased microbial contamination; in addition to this vacuum packaging and corrugated fibre board packaging lower the intracellular pH level, this provides a protective barrier against food borne pathogens (32). It was also reported that the main function of vacuum packaging is to maintain cell wall and firmness which also found to inhibit microbial growth in fresh cut produce (33). Inferred that ventilated CFB Packaging pomegranate arils had lower microbial count (bacteria, fungi, yeast and mould count) than control and also stated that modified atmospheric packaging was advantageous in minimizing the bacterial count only (34).

## Conclusion

In this study, the overall results indicated that both vacuum packaging and CFB with 10 % vent at cold storage had synergistic effect in manila tamarind fruit pods, not only in extending the shelf life but also maintained the physiological and biochemical attributes of manila tamarind. Since post-harvest loss is a major problem in fruit crops the packaging will be capable of protecting the product from the transport hazards; minimising the physiological and biochemical changes and losses in weight and storage helps in preventing from microbial damage. Considering the long-term needs of eco-systems other alternatives available like corrugated fibre board boxes and vacuum packaging are used. The CFB with 10 % vent is effective than vacuum packaging due to its cost and easy availability in all areas. So, instead of using vacuum packaging, the CFB with 10 % vent is effective packaging method for manila tamarind fruit to extend the shelf-life.

## Acknowledgements

The support and guidance of all the peer-reviewed manuscript by reviewers are very much appreciated.

## Funding information

This work was supported providing assistance by Lovely offset Printers Private Limited, Siva Kasi, Tamil Nadu, India.

## Authors' contributions

KSA: Writing of original draft and conceptualization. ATSKRR: Revision of draft, inclusion of tables and figures, proof reading. KSAATSKRRCC: Revision, formatting and Supervision. All the authors read and approved the final version of the manuscript.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interests to declare.

**Ethical issues:** There is no ethical issues.

## Declaration of generative AI and AI-assisted technologies in the writing process

For paraphrasing few sentences, I have used only the Chatgpt AI tool. For data running and analysis, I have used AGRES software version SVN-1FN.

## References

1. Arthi L, Kumar S, Shakila A, Prabudoss V. Standardization of integrated nutrient management for enhancing the yield of Manila tamarind (*Pithecellobium dulce*) cv. Local. *Crop Research*. 2023; 58(5and6):248-50. <https://doi.org/10.1007/s00608-023-01151-8>
2. Ashok A, Kayalvizhi K, Ravivarman J. Advanced production techniques in Manila Tamarind (*Pithecellobium dulce*). *Agriculture & Food e-newsletter*. 2020.

3. Kumar S, Mukherjee A, Dutta J. Chitosan based nanocomposite films and coatings: Emerging antimicrobial food packaging alternatives. Trends in Food Science & Technology. 2020;97:196-209. <https://doi.org/10.1016/j.tifs.2020.01.002>
4. Gupta N, Jain S. Storage behavior of mango as affected by post-harvest application of plant extracts and storage conditions. Journal of Food Science and Technology. 2014;51:2499-507. <https://doi.org/10.1007/s13197-012-0774-0>
5. Ranganna S. Handbook of analysis and quality control for fruit and vegetable products: Tata McGraw-Hill Education; 1986.
6. Sadasivam S, Balasubramanian. TJTNAU, Coimbatore, India. Practical Manual in Biochemistry. 1987;14.
7. Hedge J, Hofreiter B, Whistler R. Carbohydrate chemistry. Academic Press, New York. 1962; 17:371-80.
8. Srivastava A, Kohli D, Vishnoi S, Kumar S, Badola R. Quality evaluation of prepared guava-orange fruit bar. International Journal of Chemical Studies. 2019;7(4):1574-81.
9. Singh R, Singh M. Character association trend among yield attributing traits in pigeonpea [*Cajanus cajan* (L.) Millsp.]. Indian Journal of Science and Technology. 2016;9(6):1-4. <https://doi.org/10.17485/ijst/2008/v1i6.9>
10. Kalia A, Gupta RP. Microbiology of fresh and processed fruits. Handbook of Fruits and Fruit Processing. 2012;51-72. <https://doi.org/10.1002/9781118352533.ch4>
11. Lufu R, Ambaw A, Opara UL. The contribution of transpiration and respiration processes in the mass loss of pomegranate fruit (cv. Wonderful). Postharvest Biology and Technology. 2019;157. <https://doi.org/10.1016/j.postharvbio.2019.110982>
12. Brat P, Lechaudel M, Segret L, Morillon R, Hubert O, Gros O, et al. Post-harvest banana peel splitting as a function of relative humidity storage conditions. Acta Physiologiae Plantarum. 2016;38:1-14. <https://doi.org/10.1007/s11738-016-2253-0>
13. Madhana Supriya R, Sudheer K. Development and evaluation of modified atmosphere packed passion fruit (*Passiflora edulis*): Department of Post-Harvest Technology and Agricultural Processing; 2012.
14. Rezaiyan Attar F, Sedaghat N, Pasban A, Yeganehzad S, Hesarinejad MA. Modified atmosphere packaging with chitosan coating to prevent deterioration of fresh in-hull Badami's pistachio fruit. Chemical and Biological Technologies in Agriculture. 2023;10(1):16. <https://doi.org/10.1186/s40538-023-00393-9>
15. Mohammed M, Alqahtani N, El-Shafie H. Development and evaluation of an ultrasonic humidifier to control humidity in a cold storage room for postharvest quality management of dates. Foods. 2021;10(5):949. <https://doi.org/10.3390/foods10050949>
16. Gol NB, Chaudhari ML, Rao TR. Effect of edible coatings on quality and shelf life of carambola (*Averrhoa carambola* L.) fruit during storage. Journal of Food Science and Technology. 2015;52:78-91. <https://doi.org/10.1007/s13197-013-0988-9>
17. Sun Q, Zhang N, Wang J, Zhang H, Li D, Shi J, et al. Melatonin promotes ripening and improves quality of tomato fruit during postharvest life. Journal of Experimental Botany. 2015;66(3):657-68. <https://doi.org/10.1093/jxb/eru332>
18. Ashitha G, Preetha P, Varadharaju N. Effect of post-harvest hexanal treatment on shelf life of guava. Int J Curr Microbiol App Sci. 2019;8(7):1057-67. <https://doi.org/10.20546/ijcmas.2019.807.127>
19. Buvé C, Kebede BT, De Batselier C, Carrillo C, Pham HT, Hendrickx M, et al. Kinetics of colour changes in pasteurised strawberry juice during storage. Journal of Food Engineering. 2018;216:42-51. <https://doi.org/10.1016/j.jfoodeng.2017.08.002>
20. Hajebi Seyed R, Rastegar S, Faramarzi S. Impact of edible coating derived from a combination of *Aloe vera* gel, chitosan and calcium chloride on maintain the quality of mango fruit at ambient temperature. Journal of Food Measurement and Characterization. 2021;15:2932-42. <https://doi.org/10.1007/s11694-021-00861-6>
21. Brizzolara S, Manganaris GA, Fotopoulos V, Watkins CB, Tonutti P. Primary metabolism in fresh fruits during storage. Frontiers in Plant Science. 2020;11:80. <https://doi.org/10.3389/fpls.2020.00080>
22. Hari V. Plant foods for nutritional good health: Notion Press; 2019.
23. Enaru B, Dreţcanu G, Pop TD, Stănilă A, Diaconeasa Z. Anthocyanins: Factors affecting their stability and degradation. Antioxidants. 2021;10(12):1967. <https://doi.org/10.3390/antiox10121967>
24. Kaur S, Bal J. Effect of chemicals and pedicel retention on storage life and quality of ber fruits (*Zizyphus mauritiana* Lamk.). 2014.
25. Yadav A, Kumar N, Upadhyay A, Fawole OA, Mahawar MK, Jalgaonkar K, et al. Recent advances in novel packaging technologies for shelf-life extension of guava fruits for retaining health benefits for longer duration. Plants. 2022;11(4):547. <https://doi.org/10.3390/plants11040547>
26. Ali A, Hei GK, Keat YW. Efficacy of ginger oil and extract combined with gum arabic on anthracnose and quality of papaya fruit during cold storage. Journal of Food Science and Technology. 2016;53:1435-44. <https://doi.org/10.1007/s13197-015-2124-5>
27. Meighani H, Sadat-Hosseini M. Assessment of quality and biochemical changes in Indian jujube fruits (*Zizyphus mauritiana* Lamk.): Effect of chitosan coating and putrescine treatments. International Journal of Horticultural Science and Technology. 2024;11(4):481-90.
28. Saki M, ValizadehKaji B, Abbasifar A, Shahrjerdi I. Effect of chitosan coating combined with thymol essential oil on physicochemical and qualitative properties of fresh fig (*Ficus carica* L.) fruit during cold storage. Journal of Food Measurement and Characterization. 2019;13:1147-58. <https://doi.org/10.1007/s11694-019-00030-w>
29. Candir E, Özdemir AE, Aksoy MC. Effects of modified atmosphere packaging on the storage and shelf life of Hicaznar pomegranate fruits. Turkish Journal of Agriculture and Forestry. 2019;43(2):241-53. <https://doi.org/10.3906/tar-1801-119>
30. Carlin F. Fruits and vegetables. Food Microbiology: Fundamentals and Frontiers. 2012;187-201. <https://doi.org/10.1128/9781555818463.ch8>
31. Dukare AS, Singh RK, Jangra RK, Bhushan B. Non-fungicides-based promising technologies for managing post-production penicillium induced spoilage in horticultural commodities: a comprehensive review. Food Reviews International. 2022;38(3):227-67. <https://doi.org/10.1080/87559129.2020.1727497>
32. Saleh I. Effects of active phytochemicals of *Prosopis juliflora* as anti-spoiling agents of postharvest fruits. 2022
33. Huynh NK, Wilson MD, Eyles A, Stanley RA. Recent advances in postharvest technologies to extend the shelf life of blueberries (*Vaccinium* sp.), raspberries (*Rubus idaeus* L.) and blackberries (*Rubus* sp.). Journal of Berry Research. 2019;9(4):687-707. <https://doi.org/10.3233/JBR-190421>
34. Moradinezhad F, Heydari A, Ansarifard E. Enhancement of bioactive compounds and nutritional quality of pomegranate arils during storage by ultrasound pretreatment. Scientia Horticulturae. 2024;337. <https://doi.org/10.1016/j.scienta.2024.113499>