



ISSN: 2348-1900

**Plant Science Today**

<http://www.plantsciencetoday.online>



## Research Communication

# A report on new chromosome number of three *Dioscorea* species

Chiranjit Paul\* & Bimal Debnath

Plant Diversity and Forest Biotechnology Laboratory, Department of Forestry and Biodiversity, Tripura University, Suryamaninagar 799022, India

### Article history

Received: 25 January 2019

Accepted: 04 April 2019

Published: 27 April 2019

### Abstract

Chromosomal study conducted in nine species of *Dioscorea* from different forest belts of Tripura revealed that their somatic chromosome number ranged from  $2n=40$  to  $2n=60$ . The record of  $2n=40$  chromosome in the sexual phenotypes of *Dioscorea hamiltonii*, *Dioscorea glabra* and *Dioscorea pubera* are the first time report from Tripura, North East India. Moreover the somatic chromosome counts of  $2n=60$  in *Dioscorea pentaphylla* would be attributed as a new cytotype. However at the respective ploidy level no difference in somatic chromosome count was observed between their sexes.

**Keywords:** First report; chromosome number; ploidy level; *Dioscorea*

### Publisher

Horizon e-Publishing Group

**Citation:** Paul C, Debnath B. A report on new chromosome number of three *Dioscorea* species. Plant Science Today 2019;6(2):147-150. <https://doi.org/10.14719/pst.2019.6.2.490>

**Copyright:** © Paul & Debnath (2019). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>).

### \*Correspondence

Chiranjit Paul

✉ [chiranjit.paul869@gmail.com](mailto:chiranjit.paul869@gmail.com)

**Indexing:** Plant Science Today is covered by Scopus, CAS, AGRIS, CABI, Google Scholar, etc. Full list at <http://www.plantsciencetoday.online>

## Introduction

The genus *Dioscorea* commonly known as yam, a perennial climber belongs to the family Dioscoreaceae is highly economically important tuberous crop (1). *Dioscorea* tuber secure fourth rank as a tuber crop after potato, cassava and sweet potato (2). The taxon is mostly dioecious in nature although few monoecious species were also recorded (3-7). Interestingly, the male and female plants of *Dioscorea* species produced morphologically distinct tuber. Because of small chromosome sizes *Dioscorea* is considered as one of the most challenging genera with respects to cytotaxonomic and cytogenetic perspectives (8-11). However, the occurrence of polyploidy in cultivated *Dioscorea* species is common possibly due to the usage of tubers or bulbils for multiplication.

Previous cytological investigation expounded *D. alata* somatic chromosome number ranged from  $2n=30$  to  $2n=80$  (12-15). Lack of sex specific chromosomal information prompted us to evaluate the chromosome number of two different sexual phenotypes of nine *Dioscorea* species collected from different forest belts of Tripura.


## Materials and Methods

For chromosomal study, healthy root tips were collected from tuber of male and female plants of all the *Dioscorea* species from the experimental garden of Tripura University campus. The collected healthy fresh root tips were carefully washed and pre-treated in saturated solution of Para-dichloro benzene (PDB) at 4°C for 5 min and finally at 12°C

**Table 1:** Chromosomal profile of the male and female plant of nine *Dioscorea* species found in Tripura

Name of plant species	Sex specific chromosome number		2n	Established Reports
	2n	2n		
	Male	Female	References	
<i>Dioscorea alata</i>	60	60	30, 40, 50, 60, 70, 80	Sundara Raghavan 1958, Essad 1984
<i>Dioscorea bulbifera</i> var. <i>bulbifera</i>	40	40	36, 40, 54, 60	Miege 1954
<i>Dioscorea glabra</i>	40	40		***
<i>Dioscorea hamiltonii</i>	40	40		***
<i>Dioscorea hispida</i>	40	40	40	Sundara Raghavan 1958, Essad 1984
<i>Dioscorea oppositifolia</i>	40	40	40	Smith 1937, Essad 1984
<i>Dioscorea pubera</i>	40	40		***
<i>Dioscorea pentaphylla</i>	60	60	40, 80, 70	Sundara Raghavan 1958, 1959, Essad 1984
<i>Dioscorea wallichii</i>	40	40	40	Sundara Raghavan 1959, Essad 1984

\*\*\* First report of chromosome number



**Fig. 1:** Microphotographs of chromosomes of *Dioscorea* species 1-*D. alata* (male), 2n=60; 2. *D. alata* (female), 2n=60; 3. *D. bulbifera* var. *bulbifera* (male), 2n=40; 4. *D. bulbifera* var. *bulbifera* (female), 2n=40; 5. *D. glabra* (male), 2n=40; 6. *D. glabra* (female), 2n=40 7. *D. hispida* (male), 2n=40; 8. *D. hispida* (female), 2n=40; 9. *D. pubera* (male), 2n=40; 10. *D. pubera* (female), 2n=40; 11 *D. pentaphylla* (male), 2n=60; 12. *D. pentaphylla* (female), 2n=60; Scale bars=(1, 3, 10 & 12)=200µm; (4,6&8)=250 µm; (2,5,7, 9&11) =300 µm.




Fig. 2: Microphotographs of chromosomes of *Dioscorea* species. 13. *D. oppositifolia* (male),  $2n=40$ ; 14. *D. oppositifolia* (female),  $2n=40$ ; 15. *D. wallichii* (male),  $2n=40$ ; 16. *D. wallichii* (female),  $2n=40$ ; 17. *D. hamiltonii* (male),  $2n=40$ ; 18. *D. hamiltonii* (female),  $2n=40$ , Scale bars= (13 & 17) =300 $\mu$ m; (14) = 250  $\mu$ m; (15, 16 & 18) =250  $\mu$ m.

for 3 h followed by fixation in acetic acid: ethyl alcohol (1: 3) for 24 h. Then root tips were treated with 45% acetic acid for 10 min and stained in 2% aceto-orcein and 1(N) HCl (9:1) mixture for overnight. After staining the root tips were squashed in 45% acetic acid and studied under a compound microscope (16).

## Results and Discussion

In the present study the somatic chromosome count of both the sex forms of nine *Dioscorea* species was determined and it was found that their somatic chromosome count ranged from  $2n=40$  to  $2n=60$  (Table 1, Fig. 1 & 2). It was also observed that the male and female plant of *Dioscorea bulbifera* var. *bulbifera*, *D. glabra*, *D. hamiltonii*, *D. hispida*, *D. oppositifolia*, *D. pubera* and *D. wallichii* had a chromosome count of  $2n=40$ . In comparison to these the diploid chromosome number of the sex forms of *D. alata*, and *D. pentaphylla* were recorded as  $2n=60$  chromosome. Curiously the diploid chromosome number of each sex form of three *Dioscorea* species viz. *D. hamiltonii*, *D. pubera* and *D. glabra* also had a chromosome count of  $2n=40$  and thus no difference was observed between the sexes at chromosome number level. This study also revealed that the most of the *Dioscorea* species found in Tripura have diploid chromosome number  $2n=40$  and two species viz. *D. alata* and *D. pentaphylla* have somatic chromosome count  $2n=60$  in their respective phenotypes as was reported by earlier researchers (8,17-20). The present study also depicts at the basic chromosome number, is indeed  $X=10$  in all the dioecious species studied in this exploration. Similar such results were also recorded in previous findings (9,22). However, basic number  $X=8$ , 9 and 12 (8) reported by the earlier worker was not exposed in this

investigation. The Ploidy level of *Dioscorea* species is an important trait for the utilization of these species in breeding programme as well as in the enhancement of ploidy manipulation in inter- and intra-specific crosses.

## Conclusion

The present study embodies the first time chromosomal report ( $2n=40$ ) in three dioecious species of *Dioscorea* viz. *D. hamiltonii*, *D. glabra* and *D. pubera* carried out from the forest belts of Tripura. In general no difference was observed in somatic chromosome number in relation to sex in all the species studied. The detail analysis of their karyotype is essential for further exploitation of germplasm conservation.

## Authors' contributions

CP carried out the experiment and wrote the manuscript. BD conceived of the presented idea. Both the authors contributed equally to the final manuscript.

## Acknowledgement

Authors are highly gratitude to Prof. Sangram Sinha, Department of Botany, Tripura University, for his valuable advice to improve the manuscript. We are also indebted to the CSIR, New Delhi for their financial support [ref. no. 38(1371)/13/EMR-II].

## References

1. Coursey DG. Yam: An Account of the Nature, Origins, Cultivation and Utilization of the useful Member of the Dioscoreaceae. Longmans, London; 1967.



2. Lev LS, Shriver AL. A trend analysis of yam production, area, yield, and trade (1961-1996), in Ligname, Plante Seculaire et Culture Davenir. Actes du Seminaire International cirad-inra-orstom-coraf (Montpellier: CIRAD); 1998.
3. Botanical Research and Herbarium Management System, 1985. University of Oxford <https://herbaria.plants.ox.ac.uk/bol/plants400/Profiles/CD/Dioscorea>
4. Zhengyi W, Peter H, Raven, De Yuan Hong. Flora of China. Beijing: Science Press, St. Louis: Missouri Botanical Garden Press; 1994.
5. Edeoga HO, Okoli BE, Agba I. Observations on the floral morphology of some *Dioscorea* species (Dioscoreaceae) in Nigeria. Folia Geobotanica 1998; 33: 201-205. <https://doi.org/10.1007/BF02913346>
6. Flora of North America Editorial Committee, eds. Flora of North America. Volume 26, New York and Oxford; 2002.
7. Kumar S, Das G, Shin HS, Patra JK. *Dioscorea* spp. (A Wild Edible Tuber): A Study on Its Ethnopharmacological Potential and Traditional Use by the Local People of Similipal Biosphere Reserve, India. Front. Pharmacol 2017;8:52. <https://doi.org/10.3389/fphar.2017.00052>
8. Essad S. Variation géographique des nombres chromosomiques de base et polyploidie dans le genre *Dioscorea* a propos du denombrement des especes transversa Brown. pilosiuscula Bert et trifida Agron 1984;4:611-617. <https://doi.org/10.1051/agro:19840704>
9. Zoundjihepon J, Essad S, Toure B. Denombrement chromosomique dans dix groupes varietaux du complexe *Dioscorea cayenensis-rotundata*. Cytologia 1990;55:115-120. <https://doi.org/10.1508/cytologia.55.115>
10. Gamiette F, Bakry F, Ano G. Ploidy determination of some yam species (*Dioscorea* spp.) by flow cytometry and conventional chromosomes counting. Genet Resour Crop Evol. 1999;46:19-27. <https://doi.org/10.1023/A:1008649130567>
11. Dansi A, Pillay M, Mignouna HD, Dainou O, Mondeil F, Moutairou K. Ploidy level of the cultivated yams (*Dioscorea cayenensis*/*D. rotundata* complex) from Benin Republic as determined by chromosome counting and flow cytometry. ACSJ 2000;8:355-364. <https://doi.org/10.4314/acsj.v8i4.27676>
12. Sharma AK, Deepesh ND. Polyploidy in *Dioscorea*. Genetica 1956;28:112-120. <https://doi.org/10.1007/BF01694314>
13. Franklin WM, Ortiz S. Chromosome numbers and behavior in some species of *Dioscorea*. Cytologia 1963;28:96-101. <https://doi.org/10.1508/cytologia.28.96>
14. Abraham K, Nair PG. Polyploidy and sterility in relation to sex in *Dioscorea alata* L. (Dioscoreaceae). Genetica 1991;83:93-97. <https://doi.org/10.1007/BF00058525>
15. Egesi CN, Pillay M, Asiedu R, Egunjobi JK. Ploidy analysis in water yam, *Dioscorea alata* L. germplasm. Euphytica 2002;128:225-230. <https://doi.org/10.1023/A:1020868218902>
16. Sharma AK, Sharma A. Chromosome Techniques. Theory and Practice. Butterworth-Heinemann Ltd., 1980.
17. Sundara Raghavan R. A chromosome survey of Indian Dioscoreas. Proc Indian Acad Sci Sect B. 1958; 48:59-63. <https://doi.org/10.1007/BF03050251>
18. Miegé J. Nombres chromosomiques et repartition géographique de quelque plantes tropicales et equatoriales. Rev Cytol Paris 1954;15:312-348. <http://www.documentation.ird.fr/hor/fdi:010056353>
19. Cox HE, Pearson D. The Chemical Analysis of Foods Chemical. Publishing Co Inc., New York; 1962.
20. Smith BW. Notes on the cytology and distribution of the Dioscoreaceae. Bull Torr Bot Club. 1937; 64: 189-197. <https://doi.org/10.2307/2481142>
21. Ramachandran K. Cytological studies in Dioscoreaceae. Cytologia 1968;33:27-33. <https://doi.org/10.1508/cytologia.33.401>
22. Dansi A, Pillay M, Mignouna HD, Dainou O, Mondeil F, Moutairou K. Ploidy level of the cultivated yams (*Dioscorea cayenensis*/*D. rotundata* complex) from Benin Republic as determined by chromosome counting and flow cytometry. ACSJ 2000a; 8:355-364. <https://doi.org/10.4314/acsj.v8i4.27676>

