



REVIEW ARTICLE

Exploring issues and solution in biodiversity management at Ramsar sites

Athira A Nair¹, KP Ragunath^{1*}, S Pazhanivelan¹, D Muthumanickam¹, PC Prabu²

¹Department of Remote Sensing and GIS, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

²Department of Environmental Science, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

*Email: ragunathkp@tnau.ac.in



ARTICLE HISTORY

Received: 03 September 2024

Accepted: 02 November 2024

Available online

Version 1.0 : 29 December 2024

Version 2.0 : 24 February 2025



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Athira AN, Ragunath KP, Pazhanivelan S, Muthumanickam D, Prabu PC. Exploring issues and solution in biodiversity management at Ramsar sites. Plant Science Today. 2024;11(sp4):01-20.
<https://doi.org/10.14719/pst.4938>

Abstract

Ramsar sites are designated wetlands of international importance under the Ramsar convention, an intergovernmental treaty established in 1971. These sites play a crucial role in conserving biodiversity and providing ecosystem services. The Ramsar convention encourages countries to designate and manage Ramsar sites sustainably, aiming to maintain the ecological character of these wetlands. Wetlands are areas that include marsh, fen, peat land or water and it is well known that these wetlands are important ecosystems that have a major positive impact on productivity, biodiversity and the well-being of residents. Though preserved, Ramsar sites face numerous threats, including habitat loss, pollution, invasive species and climate change, warranting management, conservation efforts and international cooperation to safeguard these valuable wetland ecosystems for future generations. Pollution from agriculture, industry and urban runoff threatens water quality. Climate change exacerbates these issues, affecting wetland ecosystems and altering migration patterns. Due to the importance of Ramsar sites and to preserve for better livelihood, this paper explores the issues and challenges pertaining to the biodiversity of Ramsar sites in Tamil Nadu for better management through appropriate solutions.

Keywords

biodiversity; conservation; Ramsar site; wetland

Introduction

Biodiversity, or biological diversity, refers to the variety of life on earth, including species diversity, genetic variation and ecosystems. It is critical in ecosystem services like pollination, water purification and climate regulation (1). It also contains a vast spectrum of plants, animals, fungi, microorganisms and diverse ecosystems ranging from oceans to mountain peaks (2). For instance, coral reefs support numerous marine species and provide resources for coastal communities. Biodiversity also enhances resilience to environmental changes and diseases, making it essential for sustainable development and human well-being. Understanding its connections highlights its importance in maintaining ecological balance and supporting life. Biodiversity mapping is necessary to identify the flora and fauna in the region. It involves creating spatial representations that show the distribution, composition, and richness of species, ecosystems and genetic variation across landscapes or regions. It integrates the abundance, variability and diversity of life, spanning ecosystems, species and genetic variation within the species (3). It also contains a vast spectrum of plants, animals, fungi, microorganisms and diverse ecosystems

ranging from oceans to mountain peaks and everything in between. Biodiversity is fundamental in stabilizing ecosystems, regulating the water cycle and ensuring resilience against climate change. It provides numerous ecosystem services that are essential for human well-being. These services include clean air and water, fertile soil and climate regulation.

Additionally, biodiversity supports the pollination of crops, pest control and the provision of food, medicine and raw materials. Mapping biodiversity is essential for identifying the distribution of flora and fauna, understanding ecosystem functions and monitoring changes over time. It involves creating spatial representations that show the distribution, composition and richness of species, ecosystems and genetic variation across landscapes or regions. It is crucial for conservation, ecological research, land management, and policy-making. It is also essential for conservation, environmental research, land management and policy-making (4).

From a biodiversity perspective, areas with continuous water sources are highly productive and biologically diverse, *i.e.*, wetlands, river basins and floodplains. (5). Wetlands lie between vegetative and aquatic systems, with shallow water overlaying the ground and the water table, often at or near the surface (6). Wetlands, as defined by the Ramsar convention on Wetlands (established in 1971), are "areas of marsh, fen, peatland or water" and are broadly categorized into three types: inland wetlands, marine or coastal wetlands and man-made wetlands (7). It encompasses marine and inland marshes, covering over 12.1 million km², with 54% permanently submerged and 46% seasonally inundated. A Ramsar site is a wetland recognized for its international importance, protected under the Ramsar convention—an international treaty dedicated to the conservation and sustainable use of wetlands worldwide. As of 2024, there are over 2453 Ramsar sites globally, covering a total area of more than 255 million ha, with participation from 171 national governments. Coastal marine and inland marshes encompass over 12.1 million km², of which 54% are continuously underwater and 46% are seasonally inundated. Ramsar sites are crucial in mitigating these effects by protecting vital wetland ecosystems. These sites provide essential services, such as flood regulation, water purification and carbon sequestration, essential for climate adaptation and mitigation. (8). Ramsar sites in Tamil Nadu face several challenges, including habitat degradation, pollution and lack of awareness that necessitate effective management strategies to ensure long-term conservation and sustainable use.

Current status of Ramsar sites

The convention on wetlands, established in 1960, is the oldest global intergovernmental agreement. It was formulated through dialogues among states and non-governmental organizations in reaction to the escalating loss and deterioration of wetland ecosystems, particularly those vital for migrating waterbirds. The treaty was officially adopted in Ramsar, Iran, in 1971 and came into effect in 1975 (9). Today, the Ramsar sites are the most extensive network of protected areas globally, containing over 2400 Ramsar sites within the

territories of 172 contracting parties worldwide, spanning more than 2.5 million km² (10). The world's first site was the Cobourg Peninsula in Australia, which was designated in 1974. Among the largest sites are Rio Negro in Brazil, covering 120000 km², Ngiri-Tumba-Maindombe in the Democratic Republic of Congo and Queen Maud Gulf in Canada, both spanning over 60000 km² each. Conversely, some sites are as modest as 1 ha in size. Table.1. describes Ramsar sites located in specific regions of countries.

Ramsar sites in India

India is home to diverse wetlands that provide vital ecosystem services, including water supply, flood control and biodiversity conservation (11) (Fig. 1). Many of these wetlands have been designated as Ramsar sites, recognized under the Ramsar convention for their International Importance in conserving wetland biodiversity. India has a total of 75 Ramsar sites (Table 2). Tamil Nadu has a maximum number of Ramsar sites (14), followed by Uttar Pradesh (12). The mission established by the convention is: "the conservation and wise use of all wetlands through local and national actions and international cooperation, as a contribution towards achieving sustainable development throughout the world." (convention on Wetlands, Ramsar, 1971).

India has increased its Ramsar sites (Wetlands of International Importance) from 75 to 80 by designating five additional wetlands. The newly recognized sites include Karaivetti Bird Sanctuary and Longwood Shola Reserve Forest in Tamil Nadu as well as Magadi Kere Conservation Reserve, Ankasamudra Bird Conservation Reserve and Aghanashini Estuary in Karnataka.

Ramsar sites in Tamil Nadu

In 2023, the Union Ministry of Environment approved 2 additional Ramsar sites: Karaivetti Bird Sanctuary and Longwood Shola Reserve Forest. These additions reflect India's dedication to conserving wetlands of international significance. With this addition, Tamil Nadu now has a total of 16 Ramsar sites, the highest number in the country (Table 3 and Fig. 2) (The Hindu, February 1, 2024) With this addition, Tamil Nadu now has a total 16 Ramsar sites, the highest in the country (The Hindu, February 1, 2024).

The Karikili Bird Sanctuary

It borders the small village of Karikili, located 8 km north of Vedanthangal, partially extending into Maduranthangam Taluk, Chengalpattu District, Tamil Nadu (Fig. 3). The sanctuary includes 2 rain-fed, non-perennial irrigation tanks spanning 58.442 ha, characterized by generally flat terrain with intermittent undulations and gentle slopes. It consists of rich diversity of flora and fauna, including mangroves like *Avicennia* and *Rhizophora*, herbaceous plants, and wetland vegetation. It is a vital habitat for numerous bird species, such as the Painted Stork, Black-headed Ibis and Little Egret as well as small mammals like the Indian Fox and various reptiles, especially during the migratory season. Situated at an elevation of 100 m above mean sea level (MSL), the sanctuary experiences variable rainfall, ranging from 400 mm to nearly 1700 mm annually. This variability leads to periodic flooding during the monsoon months (October to December). It is situated at an elevation of 100 m above mean sea level

Table 1. Ramsar sites located in specific regions across different countries

Sl.No.	Continent	Country	Notable Ramsar site	Total area (ha)	Reference
1.	Europe	United Kingdom	The Wash, Solway Firth, Loch Lomond, Lough Neagh and Lough Beg, Dyfi Estuary	154143	(75)
		France	Camargue, Bassin d'Arcachon, Marais Poitevin, Baie de Somme, Etang de Biguglia	225450	(76)
		Spain	Donana National Park, Ebro Delta , Tablas de Daimiel, Albufera de Valencia, Lagunas de Ruidera, Marismas de Santoña	155480	(77)
		Germany	Lake Constance, Wadden Sea, Müritz National Park, Oder Delta , Schorfheide-Chorin Biosphere Reserve, Donau-Auen National Park, Lake Neusiedl	5426100	(78)
		Netherlands	Wadden Sea, Oostvaardersplasp Nature Reserve, Lake IJsselmeer, Biesbosch National Park, Deurnsche Peel and Mariapeel, Lauwersmeer	159470	(79)
		Sweeden	Lake, Vaddövikén, Kristianstads Vattenrike, Ottenby, Lake Stora Le, Vombsjön,	19750	(80)
		Greece	Lake Mikri Prespa, Lake Orestiada, Lagoon of Gialov, Kopais Lake	15700	(81)
		Italy	Venezia Lagoon, Mincio River, Orbetello Lagoon, Marismas de Coto Doñana, Cabrera Archipelago	4400	(82)
		Poland	Biebrza Wetlands	5900000	(83)
2	North America	United States	Everglades National Park	607000	(84)
		Canada	Queen Maud Gulf in Canada	6800000	(85)
		Mexico	Wetland Complex	2500000	(86)
3	Latin America	Brazil	Pantanal , Rio Negro	500000	(87)
		Argentina	Laguna Mar Chiquita	350000	(88)
		chile	Chiloe Archipelago	12000	(89)
		colombia	Santander Wetlands Complex	1400	(90)
		peru	Lake Titicaca	83000	(91)
4	Asia	china	Poyang Lake	2500000	(92)
		India	Sundarban, Keoladeo National Park	1002873	(93)
		Indonesia	Kepulauan Seribu	107000	(94)
		Iran	Lake Urmia	520000	(95)
		Bangladesh	Sundarban	601700	(96)
		Japan	Kushiro Shitsugen Wetland	26000	(97)
		Nepal	Sundarjal Wetlands, Rara Lake, Koshi Tappu Wildlife Reserve, Barandabhar Forest	20583	(98)
		Pakistan	Keenjhar Lake, Hingol National Park, Indus Delta, Rann of Kutch, Tarnab Wetland	15151600	(99)
5	Africa	South Africa	St. Lucia Wetland Park, Wakkerstroom, Barberspan, The Berg River Wetlands	335700	(100)
		Kenya	Lake Victoria, Lake Nakuru	6898800	(101)
		Uganda	Lake Mburo Wetland , Murchison Falls Wetland	40000	(102)
		Egypt	Lake Burullus, Lake Burullus, Wadi El Rayan Wetlands, Lake Qaraoun, Gulf of Aqaba	443000	(103)
		Nigeria	Ogun River Estuary, Kainji Lake National Park, Niger Delta Wetlands	2735800	(104)
		Madagascar	Tsimanampetsotsa Wetlands, Lake Alaotra, Belo Sur Mer	24000	(105)
		Ethiopia	Lake Tana	367300	(106)
6	Oceania	Australia	Cobourg Peninsula, Kakadu National Park	2130400	(107)
		New zealand	Firth of Thames	25000	(108)
		Papua	Tonda Wildlife, Varirata National Park	1003000	(109)
		New Guinea	Simeulue Island, Lorentz National Park, Mamberamo River Basin, Tonda Wildlife	7550000	(110)

Table 2. Ramsar sites in India

Sl.No	State	Name of wetlands	Total area (ha)	Citation
1	Andhrapradesh	Kolleru lake	1447133	(111)
2	Assam	Deepor beel	764372	(112)
3	Bihar	Kabartal wetland	403209	(113)
4	Goa	Nanda lake	21337	(114)
5	Gujarat	Wadhvana wetland Nalsarovar bird sanctuary Khijadia wildlife sanctuary Thol lake wildlife sanctuary	3474950	(115)
6	Haryana	Suitanpur National park Bhindawas wildlife sanctuary	42478	(116)
7	Himachal pradesh	Pong dam lake Renuka wetland Chandertal wetland	98496	(117)
8	Jammu & Kashmir	Wular lake Hokera wetland Surinsar –mansar lakes Hygam wetland Conservation reserve Shallbugh wetland conservation reserve	391501	(118)
9	Karnataka	Ranganathittu bird sanctuary	643576	(119)
10	Kerala	Asthamudi wetland Sasthamkotta lake Vembanad kol wetland	1609590	(120)
11	Ladakh	Tso kar wetland complex Tsomorin wetland		(121)
12	Madhyapradesh	Sakhya sagar Bhoj wetlands Sirpur wetlands Yashwant sagar	818166	(122)
13	Maharashtra	Lonar lake Thane creek Nandur madhameshwar	1014522	(123)
14	Manipur	Loktak lake	63616	(124)
15	Mizoram	Pala wetland	13968	(125)
16	Odisha	Chilka lake Satkosia gorge Bitarkanika mangroves Tampara lake Hirakund reservoir Anuspa lake	690904	(126)
17	Punjab	Harike lake Kanjili lake Ropar lake Nangal wildlife sanctuary Beas conservation reserve Keshopur – miani community reserve	86283	(127)
18	Rajasthan	Sambhar lake Keoladeo Ghana	782314	(128)
19	Tamil Nadu	Pichavaram mangrove Karilili bird sanctuary Vellodde bird sanctuary Vadavur bird sanctuary Chithrangudi bird sanctuary Vedanthangal bird sanctuary Vembanur wetland complex Point calimere wildlife and bird sanctuary Gulf of mannar Marine Biosphere Reserve Udayamarthandapuram Bird Sanctuary Koonthankulam Bird Sanctuary Pallikaranai Marsh Reserve Forest Kanjirankulam Bird Sanctuary Karivetti Bird Sanctuary Nanjarayan Bird Sanctuary Longwood Shola Reserve Forest	902534	(129)
20	Tripura	Rudrasagar lake	17542	(130)
21	Uttarakhad	Asan conservation reserve	103882	(131)
22	Uttarpradesh	Sur sarovar Haiderpur Sarsai nawar jheel Sandi bird sanctuary Saman bird sanctuary Bakhira wildlife sanctuary Parvati agra bird sanctuary Samaspur bird sanctuary	1242530	
23	West Bengal	Sunderbans wetland East Kolkata wetlands	1107907	(132)

Table 3. Details of Ramsar sites in Tamil Nadu (Source: Tamil Nadu Wetland Mission Authority)

Sl.No.	Name of wetland	District	Area	Declaration year
1.	Karikili Bird Sanctuary	Chengalpattu	2480	2022
2.	Point Calimere	Nagapattinam and Tiruvarur	1210	2022
3.	Pallikaranai Marshland	Chennai	2481	2022
4.	Pichavaram Mangrove	Cuddalore	2482	2022
5.	Vedanthangal Bird Sanctuary	Chengalpattu	2477	2022
6.	Vellode Bird Sanctuary	Erode	2475	2022
7.	Vembannur Wetland complex	Kanyakumari	2474	2022
8.	Gulf of Mannar Marine Biosphere Reserve	Ramanathapuram	2472	2022
9.	Chitrangudi Bird Sanctuary	Ramanathapuram	2491	2022
10.	Suchindram – Theroor wetland complex	Kanyakumari	2492	2022
11.	Koonthankulam Bird Sanctuary	Tirunelveli	2479	2022
12.	Kanjirankulam Bird Sanctuary	Ramanathapuram	2486	2022
13.	Udayamarthandapuram Bird Sanctuary	Tiruvarur	2476	2022
14.	Vaduvur Bird Sanctuary	Tiruvarur	2493	2022
15.	Karaivetti Bird Sanctuary	Ariyalur	4.53	2024
16.	Longwood Shola Reserve Forest	Nilgiris	116	2024

(MSL).

Point Calimere

Situated near the southern extremity of Nagappattinam District, Tamil Nadu, the Point Calimere region is acknowledged for its significant contribution to avian conservation (Fig. 4). The area features diverse flora, including mangroves like *Avicennia* and *Rhizophora*, halophytic plants in salt marshes and grasses in grasslands. It is a vital habitat for numerous migratory birds, such as the Black-headed Ibis, Greater Flamingo and Painted Stork. Additionally, it supports mammals like the Indian Fox and Blackbuck as well as reptiles, including the Olive Ridley Turtle. Overall, Point Calimere plays a significant role in preserving biodiversity, particularly for migratory bird populations. During the monsoon season, water levels gradually diminish. By the end of January, most of the water in the forested and low-lying regions becomes saline due to evaporation and saltwater intrusion. By the end of January, the remaining water evaporates, leaving compacted mud that gradually transforms into fine powdery dust as summer progresses.

The Pallikaranai Marshland

Located at coordinates 12.949371 N latitude and 80.218184 E longitude, it is one of the last remaining natural wetlands in Chennai city (Fig. 5). Locally referred to as 'kazhuveli' in Tamil, signifying a floodplain or waterlogged area, it is bordered on the east by the Buckingham Canal and the Old Mahabalipuram Road, home to the Information Technology (IT) Corridor. Its flora includes wetland vegetation, such as various grasses and sedges, some mangrove species and dominant reeds like *Phragmites*. The marshland is a vital stopover for migratory birds, including the Painted Stork,

Black-winged Stilt, Common Teal and Indian Pond Heron. Additionally, it is home to small mammals like the Indian Fox, reptiles such as the Indian Cobra and monitor lizards and various frog species. This marshland is crucial for supporting both resident and migratory wildlife. Mixed residential and institutional land uses characterize its southern and western perimeters. To the north lie dense human settlements and public infrastructure like the Mass Rapid Transit System. Covering an area of 250 km² in South Chennai, the Pallikaranai Marsh drains through 2 outlets, Okkiyam Madavu and Kovalam Creek, eventually flowing into the Bay of Bengal. Its topography ensures a consistent storage capacity, establishing a quintessential wetland ecosystem.

The Pichavaram mangrove

Covering 1478 ha, it is situated between 2 significant estuaries: the Vellar estuary to the north and the Coleroon estuary to the south in the Cuddalore district (Fig. 6). This area forms the Killai backwater and Pichavaram mangroves. Interconnected by the Vellar and Coleroon river systems, these backwaters offer ample opportunities for water sports such as rowing, kayaking and canoeing. What makes the Pichavaram forest unique is the sight of mangrove trees rooted in just a few feet of water, creating a striking waterscape ideal for backwater cruises. Key flora includes mangrove species like *Avicennia marina* (Grey Mangrove), *Rhizophora apiculata* (Red Mangrove) and *Bruguiera gymnorhiza* (Black Mangrove), along with other vegetation such as *Casuarina equisetifolia* and *Phragmites karka*. The area is home to various fauna, including migratory and resident birds like the Spot-billed Duck and Painted Stork and mammals like the Fishing Cat. Reptiles such as the Saltwater Crocodile and diverse fish species, including Barramundi, thrive in this unique ecosystem. Pichavaram is vital for




Fig. 1. Location of Ramsar sites in India. (Source: Tamil Nadu Wetland Mission Authority)

biodiversity conservation and supports a wide range of species.

Vedanthangal Bird Sanctuary

The sanctuary, one of the oldest avian protected areas in the nation and Tamil Nadu, is in Maduranthagam Taluk of Chengalpattu District (formerly Kancheepuram). It includes a small irrigation tank spanning 40 ha within Vedanthangal village and the surrounding 5 km radius around the lake (Fig. 7). Key flora includes wetland vegetation such as grasses and sedges, along with Tamarind trees (*Tamarindus indica*) and Neem trees (*Azadirachta indica*), which provide essential habitats. The sanctuary is home to diverse bird species,

including the Painted Stork, Black-headed Ibis, Little Egret, Common Teal and Grey Heron. Small mammals like the Indian Fox, reptiles such as the Indian Cobra and various fish species also thrive in the area. Internationally recognized as an Important Bird and Biodiversity Area (IBA) under the code IN284, Vedanthangal meets Criteria A1 (species of global conservation concern) and A4iii (significant congregations of birds). It is categorized within the Coromandel Coast Biotic Province, known for its rich coastal wetlands and diverse bird species.

The Vellode Bird Sanctuary

It is in Vadamugam Vellode village within the Perundurai

RAMSAR SITES - TAMILNADU




Fig. 2. Location of Ramsar sites in Tamil Nadu. (Source: Tamil Nadu Wetland Mission Authority)

taluk, Erode district, Tamil Nadu, India (Fig. 8). Situated about 12 km from Erode along the Chinnamalai main road, it spans approximately 77 ha. It showcases diverse flora like water hyacinth and lotus and supporting fauna such as painted storks, jungle cats and various reptiles, all of which are crucial for local ecosystems. The sanctuary experiences temperatures ranging from a maximum of 38 °C in summer to a minimum of 19 °C in winter, creating a seasonal environment for the birds that visit the area. Rainfall occurs predominantly during the Northeast monsoon season (September to December), replenishing the sanctuary's water bodies and supporting migratory bird species.

The Vembanur wetland

Situated near Vembanur, a small hamlet in Rajakkamangalam block, Kanniyakumari district, is an

artificial inland tank covering approximately 20 ha (Fig. 9). This wetland is designated as an important bird and biodiversity area due to its critical role in supporting diverse bird species, including migratory populations. It also provides essential habitats such as marshes and reed beds, supports threatened species and offers vital ecological services like water purification and flood control. Additionally, the wetland sustains local livelihoods through fishing and ecotourism, emphasizing the importance of conservation efforts to protect this rich ecosystem. It is known for its rich biodiversity, supporting diverse flora such as Water Hyacinth and Lotus and fauna including migratory birds like the Lesser Flamingo, small mammals like Otters and various freshwater fish, all contributing to essential ecological services and biodiversity conservation. Historical records suggest that the




Fig. 3. Map of Karikili Bird Sanctuary. (Source: Tamil Nadu Wetland Mission Authority)








Fig. 4. Map of Point Calimere. (Source: Tamil Nadu Wetland Mission Authority)



This biosphere reserve stretches across Toothukudi and



<https://plantsciencetoday.online>




Fig. 9. Map of Vembanur wetland. (Source: Tamil Nadu Wetland Mission Authority)




Fig. 10. Map of the Gulf of Mannar. (Source: Tamil Nadu Wetland Mission Authority)

Chitrangudi Bird Sanctuary

This place is located 5 km from Mudukulathur and 25 km from Paramakudi in Ramanathapuram and was declared a Bird Sanctuary and protected area in 1989 (Fig. 11). The Sanctuary is enclosed by 15 m (49 ft) high embankments surrounding a community irrigation tank. The total embankment stretches for 4.01 km (2.492 miles) and encloses an area of 260 ha. It is renowned for its rich avian biodiversity, supporting key flora such as water hyacinth and lotus and fauna including migratory species like the lesser flamingo, resident birds like the painted stork and a variety of mammals, reptiles and fish, all of which underscore its critical role in conserving wildlife and their habitats. Bottom of the Fluvial landforms around the Chitrangudi wetland include floodplains from Vaigai, Varshalei, Pambar, Kottakkarai and Gundar rivers. In contrast, marine landforms feature sand mounds (Teri dunes) and barrier dunes along the coast.

The Suchindrum Theroor Wetland

This is a constituent of the Suchindrum-Theroor Manakudi Conservation Reserve, covering an area of 94.229 ha. Commonly referred to as Suchindrum Eri and Theroor Kulam, it is situated in Kanniyakumari district, Tamil Nadu (Fig. 12). It is known for its rich biodiversity, featuring key flora such as

Water Hyacinth, Lotus and various grasses, alongside limited mangrove species like *Avicennia* and *Rhizophora*. It supports diverse fauna, including migratory birds like the Lesser Flamingo and Black-tailed Godwit, resident species such as the painted stork and little egret and small mammals like jungle cats. Additionally, it is home to various reptiles and freshwater fish, making it vital for ecological balance and biodiversity conservation in the region. The Theroor tank receives water from the Thovalai channel, while the Pazhaiyar river channel feeds the Suchindram tank. Positioned at an intermittent point of the Kodaiyar River drainage system, the wetland complex serves as a collection point for water. This water is released from the Perunchanidam, Pechipparai and Kodaiyar dams and flows into the channel, draining numerous ponds.

Koonthankulam Bird Sanctuary

It is also known as Kunthankulam and is a significant artificial wetland in Tamil Nadu (Fig. 13). Encompassing an area of 72 ha, this wetland is situated near Koonthankulam village in Nanguneri Taluk of Tirunelveli district. It supports a rich diversity of flora and fauna, including key species such as various aquatic plants, migratory and resident birds, Spot-billed Duck (*Anas poecilorhyncha*), Painted Stork (*Mycteria leucocephala*), Common Coot (*Fulica atra*) small mammals, and reptiles, highlighting its significance for biodiversity conservation. Designated as a bird sanctuary in 1994, it falls under the purview of the Tamil Nadu Water Resources



Fig. 11. Map of Chitrangudi Bird Sanctuary. (Source: Tamil Nadu Wetland Mission Authority)




Fig. 12. Map of Suchindrum Theroor wetland. (Source: Tamil Nadu Wetland Mission Authority)

Department (PWD) and the Tamil Nadu Forest Department, Tirunelveli district. Constructed several centuries ago, the inland tank is interconnected by canals drawing water from rivers originating in the Western Ghats. These rivers and tanks hold immense social and cultural significance and play a pivotal role in supporting agriculture by irrigating. The interaction between the rivers and tanks surrounding Koonthankulam Bird Sanctuary is vital for both local agriculture and conservation, as these water bodies sustain wetland ecosystems, support biodiversity, engage communities in stewardship, promote ecotourism and enhance water quality, illustrating the potential for harmonious coexistence between sustainable resource management and ecological preservation.

The Kanjirankulam Bird Sanctuary

It is a designated protected area near Mudukulathur in Ramanathapuram District, Tamil Nadu, India, established in 1989 (Fig. 14). It is located within the village of Kanjirankulam in southern Tamil Nadu. The sanctuary spans an estimated area divided between Keela (lower) Kanjirankulam, covering 66 hectares, and Mela (upper) Kanjirankulam, spanning 30.231 hectares. It is a vital wetland ecosystem that supports a diverse array of flora and fauna, including key aquatic plants, *Prosopis juliflora* and *Acacia nilotica*, *Nelumbo nucifera* and a variety of migratory and resident birds, Spot-billed Duck (*Anas poecilorhyncha*), Painted Stork (*Mycteria leucocephala*), Black-winged Stilt small mammals, and reptiles, underscoring its significance for biodiversity conservation.

Udayamarthandapuram

Udayamarthandapuram is a significant bird sanctuary in Tamil Nadu, located within the administrative jurisdiction of Thiruthuraipoondi taluk, Muthupet block, Tiruvarur district (Fig. 15). Established in 1998, the sanctuary covers approximately 44 ha and is near the confluence of the Baminiyar and Kannanaar Rivers. It is a vital wetland habitat that supports a rich diversity of flora and fauna, including key aquatic plants, numerous migratory and resident birds, small mammals and various reptiles, highlighting its significance for biodiversity conservation.

The Vaduvur bird sanctuary

A large human-made irrigation tank, spanning roughly 113 hectares, is situated in Needamangalam taluk, Thiruvavur district (Fig. 16). It lies partly within the Cauvery delta, relying on water releases from the Mettur dam to sustain its ecosystem. It is a vital wetland ecosystem that supports a rich diversity of flora, species such as *Typha angustifolia* and *Phragmites australis* and fauna, Spot-billed Duck (*Anas poecilorhyncha*) Painted Stork (*Mycteria leucocephala*) Black-winged Stilt (*Himantopus himantopus*) Common Coot (*Fulica atra*) including key aquatic plants, a variety of migratory and resident birds, small mammals and reptiles, underscoring its importance for biodiversity conservation. Encircled by a substantial bund to the south and a shorter one to the north, the natural elevation acts as a barrier, retaining water up to a depth of 2.5 m. With 4 outlets and discharge canals, potable groundwater is accessible at depths ranging from 20 to 50 m.




Fig. 13. Map of Koonthankulam Bird Sanctuary. (Source: Tamil Nadu Wetland Mission Authority)




Fig. 14. Map of Kanjirankulam Bird Sanctuary. (Source: Tamil Nadu Wetland Mission Authority)




Fig. 15. Map of Udayamarthandapuram. (Source: Tamil Nadu Wetland Mission Authority)

The Karaivetti bird sanctuary:

It is situated in the Ariyalur district and covers roughly 454 ha (Fig. 17). Positioned within the central Asian Flyway, this sanctuary is a crucial breeding and foraging habitat for resident and migratory birds. It is a vital wetland ecosystem that supports a rich diversity of flora species such as *Typha angustifolia* and *Phragmites australis* and fauna, Spot-Billed Duck (*Anas poecilorhyncha*), Painted Stork (*Mycteria leucocephala*), Black-winged Stilt (*Himantopus Himantopus*). Recognized as a Ramsar site with site number 2537, the Karaivetti Bird Sanctuary holds significance as one of the

Table 4. Criteria for designating Ramsar sites

Group A: Sites containing representative, rare or unique wetland types	Group B: Sites of international importance for conserving biological diversity. Criteria based on species and ecological communities
<p>Criterion 1: If a wetland includes a representative, rare or singular illustration of a natural or near-natural wetland type located within the suitable biogeographic area, it should be deemed internationally significant.</p>	<p>Criterion 2: If a wetland sustains vulnerable, endangered, or critically endangered species or imperiled ecological communities, it merits recognition as internationally significant.</p> <p>Criterion 3: If a wetland harbors populations of plant and/or animal species crucial for preserving the biological diversity of a specific biogeographic region, it should be acknowledged as internationally significant.</p> <p>Criterion 4: A wetland should attain international importance if it aids plant and/or animal species during pivotal stages in their life cycles or offers sanctuary during unfavorable conditions, with specific criteria centered on water birds.</p> <p>Criterion 5: A wetland should be recognized as internationally significant if it consistently sustains a population of 20000 or more water birds.</p> <p>Criterion 6: A wetland's international importance should be acknowledged if it consistently hosts 1% of the total individuals in a population of a particular species or subspecies of water bird with specific criteria focused on fish.</p> <p>Criterion 7: A wetland merits international recognition if it sustains a noteworthy portion of native fish subspecies, species or families, along with diverse life-history stages, species interactions and/or populations that epitomize the advantages and values of wetlands, thus enhancing global biological diversity.</p> <p>Criterion 8: A wetland's international significance should be acknowledged if it serves as a vital food source for fishes, a spawning ground, nursery, and/or migration route crucial for fish stocks, whether within the wetland or beyond. Specific criteria are based on other taxa.</p> <p>Criterion 9: A wetland warrants international recognition if it consistently sustains 1 %the individuals within a population of one species or subspecies of non-avian animals reliant on wetlands.</p>

information among individuals or populations of the same species, including plants, animals and microorganisms (15). On the other hand, species diversity is the variation of different species within a given ecosystem or region (16). Thirdly, the ecosystem diversity includes the variety of habitats, ecosystems and physical structures, ranging from forests and wetlands to coral reefs and grasslands (17).

Biodiversity provides a vast source of genetic variation within species, which is crucial for the survival and adaptability of organisms. This genetic diversity is essential for breeding programs in agriculture, forestry and aquaculture, helping to develop crops and livestock with improved productivity, disease resistance and adaptability to changing environmental conditions. Many pharmaceuticals and traditional medicines are derived from plants, animals and microorganisms. Biodiversity provides many natural compounds used to develop drugs to treat various diseases and health conditions. Apart from these, Ramsar sites support industries such as agriculture, forestry, fisheries and tourism. These industries are vital sources of income and livelihoods for millions of people worldwide. Additionally, they enrich human cultures and societies, inspiring art, literature, spirituality and recreation. Many indigenous cultures have deep spiritual and cultural connections to the natural world, reflected in their traditions, beliefs and practices.

Biodiverse ecosystems, such as forests, wetlands and mangroves, play a critical role in regulating the earth's climate by sequestering carbon dioxide, a major greenhouse gas responsible for global warming. Biodiversity is essential for global food security. A diverse range of plant and animal species provides the genetic resources to develop new crop varieties and livestock breeds. This adaptability is crucial for coping with changing environmental conditions and pests,

ensuring a stable and nutritious food supply for growing populations. It also increases the resilience of ecosystems and societies to environmental change and disturbances, including climate change, habitat loss, pollution and invasive species. We enhance our capacity to adapt and thrive in a rapidly changing world by preserving biodiversity.

Mapping biodiversity

This mapping facilitates evidence-based conservation planning by identifying habitats, tracking species and evaluating environmental variables such as water levels and vegetation cover. This process, facilitated by Geographic Information System (GIS) and remote sensing technologies, provides a detailed understanding of the wetland's ecological diversity. Such mapping enables evidence-based conservation planning by identifying habitats, tracking species and assessing environmental variables. It helps prioritize areas for protection, monitor changes over time and engage local communities in data collection and decision-making processes. Ultimately, mapping biodiversity in Ramsar sites is essential for preserving these valuable ecosystems and ensuring their long-term health and resilience.

Biodiversity e-infrastructures encourage education and successful public policies in addition to promoting data accessibility and usability for science. This is especially true for sustainable development, as important biodiversity metrics are hampered and skewed by a lack of accurate information about species' prosperity distribution and status (18).

Mapping habitats

A GIS is used to map and evaluate diverse habitats, which aids in understanding species distribution. This information is crucial for restoring and conserving habitat.

Species distribution modeling (SDM)

A GIS is used to build models that forecast species distributions in response to environmental conditions. Species Distribution Modeling aids in the identification of appropriate habitats and prospective sites for conservation by combining data on species occurrences with environmental characteristics.

The identification of biodiversity hotspots

Areas with high levels of endemism and species richness are made easier using GIS. Conservation activities might be targeted in these areas to protect the greatest number of species.

Corridor planning

Wildlife corridors must be identified and planned using GIS to preserve connectivity between fragmented habitats. By facilitating species mobility, these corridors lower the danger of isolation and increase genetic variety.

Threat mapping

The Threats to biodiversity are mapped and analyzed using GIS. It includes habitat destruction, deforestation and climate change. This knowledge is essential for creating conservation plans that lessen these risks.

Community-based conservation

Local knowledge is incorporated into conservation efforts by including local communities through GIS. The programs to conserve biodiversity can be more successful if the resources are mapped and local communities are involved in data collection.

Monitoring and surveillance

Using satellite imagery and remote sensing, GIS makes monitoring biodiversity in real-time possible. This facilitates monitoring alterations in land use, deforestation and other environmental elements impacting biodiversity.

Management of protected areas

Geographic Information System makes planning, organizing and creating protected areas easier. It assists in planning new reserves, evaluating the efficacy of current ones and keeping track of changes in land use in the surrounding areas.

Challenges in Ramsar sites

Wetlands are among the world's most threatened ecosystems, and the wildlife that depends on them is increasingly endangered due to habitat destruction, pollution and other human activities. Wetlands face numerous and tremendous threats such as natural and anthropogenic threats and are disappearing 3 times faster than forests. Some common threats are habitat destruction, wastewater discharge, weed infestation, pollution, invasive species, over-harvesting, unregulated tourism and climate change (19).

In developing countries, key issues include inadequate knowledge and data, population and development impacts, poor regulatory and planning processes and socio-economic inequities and conflict (20-22). This is compounded by the uneven growth of Ramsar sites globally, with a need for increased site and area representation, improved management and reporting (20). As the Ramsar convention

requires, integrating wetlands into river basin management is also a challenge, with barriers including mismatched priorities and a lack of recognition of ecosystem services (21). These challenges underscore the need for comprehensive, integrative approaches that address local needs and perspectives and for increased commitment, resourcing and stakeholder engagement (22).

The management of Ramsar sites in Tamil Nadu, India, faces various challenges, including encroachments, pollution and inadequate governance (20, 23-25). Biodiversity-related problems in Ramsar sites are critical issues that threaten the rich and diverse array of plant and animal species found in these wetland ecosystems. Some problems related to biodiversity in Ramsar sites includes:

Habitat Loss and Fragmentation

Urbanization, agriculture and infrastructure development can result in the loss and fragmentation of wetland habitats, directly impacting the biodiversity of Ramsar sites. Habitat loss can lead to declines in populations of various species, affecting their ability to find suitable breeding, feeding and nesting grounds (23). Habitat loss and fragmentation are significant threats to biodiversity, often exacerbated by other factors such as hunting and fire (24). Human activities, particularly deforestation and changes in land use, are key drivers of habitat fragmentation. This process can reduce continuous habitats into smaller, distinct patches, impacting the quality and connectivity of these habitats (25). The consequences of habitat fragmentation include habitat loss, increased edge habitat and isolation effects, with variable responses from species (26). These findings underscore the need for further research and conservation efforts to address the challenges posed by habitat loss and fragmentation.

Invasive Species

The introduction of invasive plant and animal species can outcompete native species, disrupt the ecological balance and pose a threat to the diversity of Ramsar site ecosystems. The water hyacinth has significantly disrupted nutrient cycling and outcompeted native plants in many wetland ecosystems (27). Common Reed, water hyacinths, particularly invasive plants, can drastically alter ecosystems by monopolizing resources such as sunlight and water, leading to the decline of native species. The spread of these species is facilitated by various vectors, including human activities and effective management strategies are crucial in preventing new invasions (28). The term "invasive species" refers to non-native species that cause harm in their new environments (29). The increasing globalization of commerce, through the movement of goods and people, facilitated the spread of invasive species, necessitating the development of strategies to address this growing problem (30).

Pollution

Agricultural runoff, industrial discharges and untreated sewage can introduce pollutants into wetland ecosystems, affecting water quality and directly threatening aquatic biodiversity. Pollution can lead to declines in sensitive species, especially those affected by environmental pollution, defined as introducing harmful substances or products into the environment, has become a significant global issue (31). This

pollution can take various forms, including air, noise and water (32). Air pollution, in particular, is a major concern, with contaminants such as dust, fumes and gas threatening human health and the environment (33). The sources of pollution are diverse, ranging from industrial plants and garbage dumps to military facilities (34). These findings underscore the urgent need for effective pollution control measures to safeguard the environment, public health and specific water quality conditions.

Climate Change

Changes in temperature, precipitation patterns and sea levels associated with climate change can alter the habitats and distribution of species within Ramsar sites. Climate-induced stress can affect the reproductive success and survival of various plant and animal species. Climate change is a complex and multifaceted issue driven by various factors, including carbon emissions, deforestation and changes in atmospheric composition. While natural processes have historically influenced the earth's climate, human activities have significantly accelerated these changes, mainly by releasing greenhouse gases (35). The impacts of climate change are far-reaching, affecting water resources, ecosystems, food production and human health. These impacts are expected to intensify in the future, with potential consequences including extreme weather events and rising global temperatures (36).

Over-Exploitation of Resources

Unsustainable fishing, hunting and harvesting of plants can lead to declines in populations of economically and ecologically important species, affecting the overall biodiversity. Over-extraction of water can alter wetland hydrology, impacting the conditions necessary for diverse species to thrive. The over-exploitation of renewable resources is often driven by misperceptions and inappropriate mental models (37). This is evident in the case of Jordan, where overuse of groundwater resources has led to a drop in levels and deterioration in quality, with significant socio-economic implications. However, the under-exploitation of natural resources can also occur due to market failures and the double role of resource stocks in economies with overlapping generations (38). Despite these challenges, there is hope for the future of food systems, with the potential for new, more sustainable production methods (39).

Disease Spread

Increased human activity, including tourism, can contribute to the spread of diseases affecting wildlife in Ramsar sites. Disease outbreaks can have cascading effects on biodiversity, leading to declines in susceptible species. For example, the chytrid fungus (*Batrachochytrium dendrobatidis*) has caused significant declines in amphibian populations worldwide, resulting in the extinction of several frog species (40). The spread of diseases in Ramsar, Iran, is a significant public health concern, including leishmaniasis, malaria, dengue fever, Zika virus and schistosomiasis, all of which are influenced by the region's unique ecological conditions and require effective monitoring and control measures (41). Rabbit Haemorrhagic Disease, a highly fatal viral disease, has

been spreading in the Arabian Peninsula, posing potential risks to local wildlife and ecosystems. This is particularly relevant to Ramsar sites in the region, given their ecological importance and proximity to the affected areas. Similarly, there are reports on the occurrence of Rift Valley Fever, a zoonotic disease affecting livestock and humans, in the Arabian Peninsula (42). The spread of these diseases underscores the need for effective surveillance and control measures in Ramsar.

Altered Hydrology

Changes in water flow and hydrological patterns due to human activities, such as dam construction or land drainage, can impact aquatic species' reproductive and feeding behaviors (43). The alteration of hydrology in Ramsar wetlands has been a significant concern, with studies highlighting the impact on vegetation and ecosystems. The Bakhau Wetland has experienced habitat reduction for migrating birds and fish due to upstream dam building, resulting in disturbances to their nesting and feeding patterns (44). The reduced freshwater inflow in the Sundarbans has caused declines in native mangrove species like Sundari (*Heritiera fomes*), which is vital for shoreline stability and biodiversity (45). In Vembanad Lake, changes in water flow have facilitated the spread of invasive water hyacinth (*Eichhornia crassipes*), which outcompetes native flora and disrupts local fish populations (46). One study demonstrated the successful restoration of a coastal salt marsh in an Australian Ramsar site through controlled tidal inundation (47). In Greece, rising water levels at the Kerkini Reservoir led to the disappearance of reedbeds and a decline in forested areas, negatively impacting local biodiversity (48). Another study emphasized the need to consider historical inundation patterns in the restoration and management of the Doñana marshes, which have been significantly altered by human transformations (49). A study highlighted the critical need for ecosystem management and restoration in the Sambhar wetland in India, noting a substantial decrease in waterbody size and a loss of saline characteristics (50). These studies highlight the complex and varied impacts of altered hydrology in Ramsar wetlands, underscoring the need for site-specific restoration and management strategies.

Lack of Connectivity

Infrastructure development and habitat fragmentation can disrupt ecological connectivity within Ramsar sites. Lack of connectivity can isolate populations, reduce genetic diversity, and increase the vulnerability of species to environmental changes (51). The issue of connectivity in various contexts has been addressed in several studies. A study proposed an obstacle-avoiding connectivity restoration strategy for mobile robotic sensor networks (52), while a multi-hop network for Earth-Mars communication was developed in another study (53). A separate study detailed the ONERA RAMSES system, a synthetic aperture radar system and its applications in urban environments. Finally, a connectivity protocol for star topology using wireless sensor networks was also introduced (54). These studies collectively highlight the importance of connectivity in different fields and the need for innovative solutions to address connectivity challenges.

Inadequate Conservation Measures

Insufficient protection and enforcement of conservation measures can leave Ramsar sites vulnerable to various threats, jeopardizing the biodiversity within these ecosystems. Lack of effective management plans may fail to address the specific needs of different species (55). The Ramsar convention, while successful in expanding its network of wetlands, faces challenges in ensuring effective conservation measures (56). These challenges include inadequate management planning and reporting and a lack of representativeness in the network. The convention's effectiveness in preserving wintering waterbirds in the Mediterranean is also questioned, with regional disparities in its impact (57). In the case of the Deepor Beel Ramsar site in India, the lack of a single authority for wetland management and the dependence of local communities on its resources further highlight the need for improved conservation measures (58). Public awareness and education are also crucial in enhancing conservation efforts, as demonstrated in the case of Ramsar wetlands in Slovenia (59).

Management strategies available

The management strategies of Ramsar sites in Tamil Nadu can draw insights from various studies from regions like the Sundarbans and Kerala for wetland ecology, the Godavari Delta for water resource management, Odisha for community involvement, the Gulf of Mannar for climate change adaptation, Vembanad Lake for pollution control and Chilika Lake for sustainable ecotourism (60). "The need for sustainable ecotourism development is emphasized, with its application to Ramsar sites in the region being particularly significant. It highlights the importance of involving local communities in conservation efforts, a strategy that could greatly benefit Ramsar sites, including those in Tamil Nadu (61). Research highlights the significance of catchment-scale management for the sustainability of Ramsar sites, which could be a key consideration for the region's sites.

Integrated management plans

Developing comprehensive management plans these are essential as they enable a holistic approach to health challenges by optimizing resource allocation, facilitating stakeholder coordination, ensuring preparedness for emergencies, establishing evaluation metrics for continuous improvement and fostering public trust through organized and proactive management for each Ramsar site is essential (62). These plans should involve stakeholders such as government agencies, local communities, NGOs and researchers. The plans should outline strategies for conservation, restoration and sustainable use of wetland resources. A number of research has examined the development and implementation of Integrated Management Plans (IMPs) in Ramsar sites. The adoption of systems-thinking methodology, utilizing the viable system model and partial least squares path modelling, has been conducted to assess the sustainable management of Ramsar sites. A study also emphasized the importance of stakeholder participation and accountability in the planning, implementing, and reviewing of an IMP for the Okavango Delta Ramsar Site in Botswana (63). Another study focused on

the adaptive management of the Macquarie Marshes, a Ramsar-listed wetland in Australia and incorporating climate change adaptation into the IMP (64). Lastly, an evaluation of Turkey's Uluabat Integrated Wetland Management Plan was also performed, highlighting the need for plan revision, additional activities and integration with a regional rural development project (65). These studies collectively underscore the significance of stakeholder engagement, adaptive management and continuous plan evaluation in the development and implementation of IMPs in Ramsar sites.

Biodiversity conservation

Conservation of biodiversity within Ramsar sites is a priority. This involves identifying and protecting key habitats, species and ecological processes within the wetlands. Monitoring programs should be established to track changes in biodiversity over time. Established in 1975, the Ramsar convention is a key international agreement for the conservation and sustainable use of wetlands. In Jammu and Kashmir, India, Ramsar sites play a crucial role in biodiversity conservation, with 15.6% of the state-designated protected areas (66). Similarly, in Nepal, Ramsar sites are vital for conserving high-altitude, mid-hill and terai wetlands, home to endemic and endangered species (67). In Assam, India, the Deepor Beel Ramsar site is a significant wetland with rich biodiversity, including rare, vulnerable and endemic species (68). These studies collectively highlight the importance of Ramsar sites in biodiversity conservation in various regions.

Water Quality Management

Ensuring good water quality is crucial for the health of Ramsar sites. Measures such as controlling pollution from agricultural runoff, industrial discharge and sewage should be implemented. Wetland restoration projects can help improve water quality by filtering pollutants and reducing nutrient loading. Afenourir Lake in Morocco, the Indus Delta in Pakistan and the Ropar Wetland in India, all Ramsar sites, face water quality challenges due to both natural and anthropogenic factors (68-70). These challenges include the influence of climatic hazards, pollution and contaminants such as heavy metals and coliform bacteria. The need for urgent management and monitoring of these sites is emphasized, with a focus on protecting water-chemistry characteristics and developing guidance for wetland managers (69).

Sustainable use

Ramsar sites often support local communities that depend on wetland resources. Sustainable use of wetland resources should be promoted through activities such as regulated fishing, eco-tourism and traditional agriculture practices. A range of studies have explored sustainable use in RAMSAR sites. A systems thinking approach, using the Viable System Model and Partial Least Squares Path Modeling, was introduced to assess sustainable management in RAMSAR sites. Identification and assessment of developmental options were done on Ghana's coastal RAMSAR sites, focusing on environmental compatibility, technical and economic viability, and social acceptability (70). The Songor Ramsar site in Ghana, covering approximately 85000 ha, is a vital wetland ecosystem providing habitat for migratory birds, supporting

biodiversity and offering opportunities for sustainable fishing and ecotourism, while facing threats from industrialization, pollution and overfishing (71).

Community engagement

Engaging local communities in managing Ramsar sites is vital for their long-term sustainability. Community participation can include involvement in decision-making processes, environmental education programs and capacity-building initiatives.

Research and monitoring

Regular monitoring and research activities are essential for assessing the health of Ramsar sites and understanding ecosystem dynamics. This includes monitoring water levels, biodiversity indicators and human impacts on the wetlands.

International collaboration

Collaboration with international organizations and other Ramsar sites worldwide can provide valuable insights and resources for managing wetlands in Tamil Nadu. Sharing best practices and lessons learned can help improve conservation efforts. The management of Ramsar sites in Tamil Nadu can draw valuable insights from international experiences. Integrated management plans, as seen in the Great Barrier Reef catchments (72), are crucial for conservation and restoration. Biodiversity conservation, a priority in Ramsar sites, can be achieved by protecting key habitats and species. Water quality management, as demonstrated in the Mekong Delta wetlands (73), is essential for the health of these sites. Sustainable use, a key aspect of Ramsar site management, can be promoted through regulated activities such as fishing and eco-tourism (74). Community engagement, as seen in the Hikkaduwa Marine Sanctuary, is vital for the long-term sustainability of these sites. Research and monitoring, as emphasized in all the studies, are crucial for assessing the health of Ramsar sites and understanding ecosystem dynamics. Lastly, international collaboration, as highlighted in the Mekong Delta wetlands, can provide valuable insights and resources for managing wetlands in Tamil Nadu.

Conclusion

Exploring issues and solutions in biodiversity management at Ramsar sites underscores the critical importance of preserving these wetland ecosystems. Challenges such as habitat degradation, pollution, climate change impacts and unsustainable resource use demand immediate attention and innovative solutions. Implementing effective conservation measures, including habitat restoration, sustainable practices and international cooperation, is paramount. Engaging local communities, incorporating advanced technologies for mapping and monitoring and promoting adaptive management strategies are integral components of a comprehensive solution. The commitment to biodiversity management in Ramsar sites is responsible for safeguarding diverse ecosystems and investing in the well-being of nature and humanity. Through collaborative efforts, informed decision-making and a dedication to sustainable practices, we can ensure the resilience and vitality of Ramsar sites for current and future generations.

Acknowledgments

We thank our entire Department of Remote sensing and GIS, Tamil Nadu Agricultural University, Coimbatore, for their support during my research.

Authors' contributions

All the authors contributed valuable insights to this manuscript. AAN carried out the writing process (Methodology, Conceptualization) and drafted the manuscript. KPR supervised and corrected. SP aided in Data Analysis. DM participated in visualization. PC participated in coordination and helped in gathering data.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None

Declaration of generative AI and AI-assisted technologies in the writing process

While preparing this manuscript, the authors used Grammarly to correct grammar and improve language fluency.

References

- Magurran AE, Dornelas M. Biological diversity in a changing world. *The Royal Soc*; 2010. p. 3593-97. <https://doi.org/10.1098/rstb.2010.0296>
- Chivian E. Biodiversity: its importance to human health. Center for Health and the Global Environment, Harvard Medical School, Cambridge, MA. 2002;23. <https://doi.org/10.1093/oso/9780195150933.003.0014>
- Mittermeier RA, Turner WR, Larsen FW, Brooks TM, Gascon C. Global biodiversity conservation: the critical role of hotspots. *Biodiversity hotspots: distribution and protection of conservation priority areas*. Springer; 2011. p. 3-22. https://doi.org/10.1007/978-3-642-20992-5_1
- Behera M, Kushwaha S, Roy P. High plant endemism in an Indian hotspot-eastern Himalaya. *Biodiversity and Conservation*. 2002;11:669-82. https://doi.org/10.1007/978-1-4020-6444-9_18
- Ghermandi A, Van den Bergh JC, Brander LM, De Groot HL, Nunes PA. The economic value of wetland conservation and creation: A meta-analysis; 2008. <https://doi.org/10.2139/ssrn.1273002>
- Tiner RW. *Wetland indicators: A guide to wetland formation, identification, delineation, classification and mapping*; CRC press; 2016.
- Ausseil AG, Gerbeaux P, Chadderton WL, Stephens T, Brown D, Leathwick J. Wetland ecosystems of national importance for biodiversity: criteria, methods and candidate list of nationally important inland wetlands. Landcare Research Contract Report LC0708/158; 2008. <https://doi.org/10.1111/j.1365-2427.2010.02412.x>
- Gardner RC, Finlayson C, editors. In: *Global wetland outlook: state of the world's wetlands and their services to people*. Ramsar convention secretariat; 2018.
- Omondi P, Nonuw AA. Assessing effects of human activities Kipranyie wetland, Sondu Miriu river basin, Kericho County, Kenya. *East African J Environ Nat Res*. 2021;4(1):25-38. <https://doi.org/10.37284/eajenr.4.1.471>

10. López-Calatayud NC, Márquez-Romance AM, Guevara-Pérez E, Buroz-Castillo E. An approach for management modeling of a tropical wetland. *Environ Quality Manage.* 2022;31(3):423-39.<https://doi.org/10.1002/tqem.21798>
11. Bassi N, Kumar MD, Sharma A, Pardha-Saradhi P. Status of wetlands in India: A review of extent, ecosystem benefits, threats and management strategies. *J of Hydro: Regional Stud.* 2014;2:1-19.<https://doi.org/10.1016/j.ejrh.2014.07.001>
12. Asha C, Retina CI, Nandan SB, Suson P. Community structure and ecology of soft bottom polychaetes in Vembanad estuary, a Ramsar site on the South West coast of India: A time scale evaluation. *Regional Stud in Marine Sci.* 2023;67:103213.<https://doi.org/10.1016/j.rsma.2023.103213>
13. Singh Y, Singh G, Khattar JS, Barinova S, Kaur J, Kumar S, et al. Assessment of water quality condition and spatiotemporal patterns in selected wetlands of Punjab, India. *Environ Sci and Pollu Res.* 2022;29(2):2493-509.<https://doi.org/10.1007/s11356-021-15590-y>
14. Mitsch W, Cronk J. Creation and restoration of wetlands: some design consideration for ecological engineering. *Soil Restor: Soil Restor.* 1992;(17):217-59.https://doi.org/10.1007/978-1-4612-2820-2_8
15. Brown S, Swingland I, Hanbury-Tension R, Prance G, Myers N. Carbon sinks for abating climate change: can they work. Center for Environ and Soc, University of Essex; 2001.
16. Gould WR, Ray AM, Bailey LL, Thoma D, Daley R, Legg K. Multistate occupancy modeling improves understanding of amphibian breeding dynamics in the Greater Yellowstone Area. *Ecol Appl.* 2019;29(1):e01825.<https://doi.org/10.1002/eap.1825>
17. Llanos-Lizcano A, Barraza E, Narvaez A, Varela L, Caselles-Osorio A. Efficiency of pilot-scale horizontal subsurface flow constructed wetlands and microbial community composition operating under tropical conditions. *Inter J of Phytoremed.* 2019;21(1):34-42.<https://doi.org/10.1080/15226514.2018.1523874>
18. Kandas P, Minotti PG, Morandeira NS, Grimson R, González TG, González EB, et al. Remote sensing of wetlands in South America: status and challenges. *Inter J of Remote Sens.* 2018;39(4):993-1016.<https://doi.org/10.1080/01431161.2017.1395971>
19. Waly MM, Mickovski SB, Thomson C, Amadi K. Impact of implementing constructed wetlands on supporting the sustainable development goals. *Land.* 2022;11(11):1963.<https://doi.org/10.3390/land11111963>
20. Wohl E, Castro J, Cluer B, Merritts D, Powers P, Staab B, et al. Rediscovering, reevaluating and restoring lost river-wetland corridors. *Front in Earth Sci.* 2021;9:653623.<https://doi.org/10.3389/feart.2021.653623>
21. Stojanović V, Mijatov LM, Dragin AS, Cimbalević M, Obradović S, Dolinaj D, et al. Tourists' motivation in wetland destinations: Gornje Podunavlje special nature reserve case study (Mura-Drava-Danube Transboundary Biosphere Reserve). *Sustain.* 2023;15(12):9598.<https://doi.org/10.3390/su15129598>
22. Boyero L, Datri L, Lopez M, Rodríguez MC, Robertazzi M, Lopez H, et al. Urban planning in arid northern Patagonia cities to maximize local ecosystem services provision. *Ecosystem Services in Patagonia: A Multi-Criteria App for an Integr Assess: Springer;* 2021. p. 349-77.https://doi.org/10.1007/978-3-030-69166-0_17
23. Baiamonte G, Domina G, Raimondo F, Bazan G. Agricultural landscapes and biodiversity conservation: a case study in Sicily (Italy). *Biodivers and Conserv.* 2015;24:3201-16.<https://doi.org/10.1007/s10531-015-0950-4>
24. Nobuoka H, Mimura N. Assessments of early adaptations due to sea-level rises and storm surges in Asian and oceanic coastal zones. *Coastal Structures 2011: (In 2 Volumes): World Scientific;* 2013. p. 120-29.https://doi.org/10.1142/9789814412216_0011
25. Rooney R, Foote L, Krogman N, Pattison J, Wilson M, Bayley S. Replacing natural wetlands with stormwater management facilities: Biophysical and perceived social values. *Water Res.* 2015;73:17-28.<https://doi.org/10.1016/j.watres.2014.12.035>
26. Posa MRC. Peat swamp forest avifauna of Central Kalimantan, Indonesia: Effects of habitat loss and degradation. *Biol Conserv.* 2011;144(10):2548-56.<https://doi.org/10.1016/j.biocon.2011.07.015>
27. Pathak HN, Bhuju DR, Shrestha BB, Ranjitkar S. Impacts of invasive alien plants on ecosystem services of Ramsar lake cluster in middle mountain Nepal. *Global Ecology and Conserv.* 2021;27:e01597.<https://doi.org/10.1016/j.gecco.2021.e01597>
28. Claus S, Imgraben S, Brennan K, Carthey A, Daly B, Blakey R, et al. Assessing the extent and condition of wetlands in NSW. Supporting report a–conceptual framework. Office of Environ and Heritage: Sydney, NSW, Australia; 2011.
29. Claus S, Imgraben S, Brennan K, Carthey A, Daly B, Blakey R, et al. Assessing the extent and condition of wetlands in NSW: Project report. Office of Environ and Heritage, Sydney, AU; 2011.
30. Rodgers JH, Lehman RW, Gladden JB, Bell JF, Mooney FD, editors. In: Wetlands for industrial wastewater treatment at the Savannah river site: a case study. WEFTEC; 2001.<https://doi.org/10.2175/193864701790861271>
31. Xi Y, Peng S, Ciais P, Chen Y. Future impacts of climate change on inland Ramsar wetlands. *Nature Climate Change.* 2021;11(1):45-51.<https://doi.org/10.1038/s41558-020-00942-2>
32. Omotoriogun TC, Onoja JD, Tende T, Manu S, Ottosson U. Density and diversity of birds in the wetlands of Yankari Game Reserve, Bauchi, Nigeria; 2011.<https://doi.org/10.3126/jowe.v5i0.4778>
33. Ding W, Cai Z, Wang D. Preliminary budget of methane emissions from natural wetlands in China. *Atmospheric Environ.* 2004;38(5):751-59.<https://doi.org/10.1016/j.atmosenv.2003.10.016>
34. Berkowitz JF, Pietroski JP, Currie SJ. Evaluation of wetland hydrology in formerly irrigated areas. US Army Engineer Research and Development Center, Environ Laboratory; 2017.<https://doi.org/10.21079/11681/22785>
35. Harrington R, Carroll P, Carty A, Keohane J, Ryder C. Integrated constructed wetlands: concept, design, site evaluation and performance. *Intern J of Water.* 2007;3(3):243-56.<https://doi.org/10.1504/ijw.2007.015217>
36. Kolcsár LP, Soos A, Török E, Graf W, Rákossy L, Keresztes L. New faunistic records of the genus Erioptera meigen (Limoniidae, Diptera, Insecta) from Europe. *Entomologica Romanica.* 2017;21:23-44.<https://doi.org/10.24193/entomolrom.21.4>
37. Milon JW. Pastures, fences, tragedies and marine reserves. *Bull of Marine Sci.* 2000;66(3):901-16.
38. White WA, Morton RA. Wetland losses related to fault movement and hydrocarbon production, Southeastern Texas coast. *J of Coastal Res.* 1997:1305-20.<https://doi.org/10.3133/fs09101>
39. Taran M, Deb S. Plant diversity and ecosystem services of wetland-based agroforestry system in Tripura, Northeast India. *Ind J of Agroforestry.* 2019;21(2):13-17.<https://doi.org/10.1007/s10457-023-00823-5>
40. Iglesias I, Muñoz MJ, Martínez M, De la Torre A. Environmental risk factors associated with H5N1 HPAI in Ramsar wetlands of Europe. *Avian Dis.* 2010;54(2):814-20.<https://doi.org/10.1637/8970-062609-reg.1>
41. El Zein H, Stephan J, Khater C, Al-Zein MS, Dagher-Kharrat MB. Aligning terrestrial habitat typology of Lebanon with EUNIS habitat classification. *Phytocoenologia.* 2022;51(3).<https://doi.org/10.1127/phyto/2022/0390>
42. Verma AK, Prakash S. Zooplankton diversity in Guthia Taal, wetland of Bahraich (UP), India. *Inter J Zool Res.* 2020;10(2):9-18.
43. Benediktsson JA, Chanussot J, Moon WM. Advances in very-high-resolution remote sensing. *Proceedings of IEEE.* 2013;101(3):566-69.<https://doi.org/10.1109/jproc.2012.2237076>

44. Leta MK, Demissie TA, Tränckner J. Optimal operation of Nashe hydropower reservoir under land use and land cover change in Blue Nile River basin. *Water*. 2022;14(10):1606.<https://doi.org/10.3390/w14101606>
45. Dasgupta S, Sobhan I, Wheeler D. The impact of climate change and aquatic salinization on mangrove species in the Bangladesh Sundarbans. *Ambio*. 2017;46:680-94.<https://doi.org/10.1596/1813-9450-7736>
46. Glamore W, Rayner D, Ruprecht J, Sadat-Noori M, Khojasteh D. Eco-hydrology as a driver for tidal restoration: Observations from a Ramsar wetland in eastern Australia. *Plos One*. 2021;16(8):e0254701.<https://doi.org/10.1371/journal.pone.0254701>
47. Kingsford RT, Walker KF, Lester RE, Young WJ, Fairweather PG, Sammut J, et al. A Ramsar wetland in crisis—the Coorong, lower lakes and Murray mouth, Australia. *Marine and Freshwater Res*. 2011;62(3):255-65.<https://doi.org/10.1071/mf09315>
48. Cowardin LM, Golet FC. US fish and wildlife service 1979 wetland classification: A review. *Classification and Inventory of the World's Wetlands*. 1995:139-52.https://doi.org/10.1007/978-94-011-0427-2_12
49. Santamaria L, Green A, Diaz-Delgado R, Bravo M, Castellanos E, Caracoles A. A new laboratory for science and wetland restoration. Doñana, water and biosphere; Confederación Hidrográfica del Guadalquivir Ministerio de Medio Ambiente: Madrid, España; 2006:313-35.
50. Naik R, Sharma LK. Monitoring migratory birds of India's largest shallow saline Ramsar site (Sambhar Lake) using geospatial data for wetland restoration. *Wetlands Ecology and Manag*. 2022;30(3):477-96.<https://doi.org/10.1007/s11273-022-09875-3>
51. Popoff N, Gaget E, Béchet A, Dami L, Du Rau PD, Geijzendorffer I, et al. Gap analysis of the Ramsar site network at 50: over 150 important mediterranean sites for wintering waterbirds omitted. *Biodiversity and Conserv*. 2021;30:3067-85.<https://doi.org/10.1007/s10531-021-02236-1>
52. Parmentier FJW, Zhang W, Mi Y, Zhu X, van Huissteden J, Hayes DJ, et al. Rising methane emissions from northern wetlands associated with sea ice decline. *Geophysical Res Lett*. 2015;42(17):7214-22.<https://doi.org/10.1002/2015gl065013>
53. Haque A, Ali G, Badiou P. Hydrological dynamics of prairie pothole wetlands: Dominant processes and landscape controls under contrasted conditions. *Hydrological Processes*. 2018;32(15):2405-22.<https://doi.org/10.1002/hyp.13173>
54. Gqalaqha Z. Identifying wetland soil properties aiding the dormancy of Rift Valley fever vectors in Central South Africa. *University of the Free State*; 2020.<https://doi.org/10.1371/journal.pone.0232481>
55. Munishi P, Kilungu H, Jackson H, Shirima D, Bulenga G, Seki H. Wetland related livelihoods, institutions and incentives for conservation in the Great Ruaha River wetland system. *Tanzania Journal of Forestry and Nature Conserv*. 2012;81(2):34-44.
56. Kingsford RT, Bino G, Finlayson CM, Falster D, Fitzsimons JA, Gawlik DE, et al. Ramsar wetlands of international importance—improving conservation outcomes. *Front in Environ Sci*. 2021;9:643367.<https://doi.org/10.3389/fenvs.2021.643367>
57. Perennou C, Gaget E, Galewski T, Geijzendorffer I, Guelmami A. Evolution of wetlands in mediterranean region. *Water Resources in the Mediterranean Region*. 2020:297-320.<https://doi.org/10.1016/b978-0-12-818086-0.00011-x>
58. Saikia MK, Saikia PK, Bhatta R. Management perspectives for avian population conservation and enrichment in deepor beel Ramsar site, North-East India. *J of Global Biosci*. 2014;3(2):428-51.
59. Polajnar K. Public awareness of wetlands and their conservation. *Acta Geographica Slovenica*. 2008;48(1):121-46.<https://doi.org/10.3986/ags48105>
60. Hussan A, Sundaray J, Ghosal R, Mallick S. Lovesome chum of aquarium are wreaking havoc in East Kolkata wetlands, India. *Aquaculture Asia*. 2020;24(3):9-15.
61. Everard M, Harrington R, McInnes RJ. Facilitating implementation of landscape-scale water management: The integrated constructed wetland concept. *Ecosystem Services*. 2012;2:27-37.<https://doi.org/10.1016/j.ecoser.2012.08.001>
62. Sánchez-García JY, Ramírez-Gutiérrez AG, Núñez-Ríos JE, Cardoso-Castro PP, Rojas OG. Systems thinking approach to sustainable performance in Ramsar sites. *Sustain*. 2019;11(22):6469.<https://doi.org/10.3390/su11226469>
63. Mfundisi KB. Overview of an integrated management plan for the Okavango Delta Ramsar site, Botswana. *Wetlands*. 2008;28:538-43.<https://doi.org/10.1672/07-55.1>
64. Bino G, Jenkins K, Kingsford R. Adaptive management of Ramsar wetlands; 2013.
65. Gürlük S, Rehber E. Evaluation of an integrated wetland management plan: case of Uluabat (Apollonia) lake, Turkey. *Wetlands*. 2006;26(1):258-64.[https://doi.org/10.1672/0277-5212\(2006\)26\[258:eoiwjm\]2.0.co;2](https://doi.org/10.1672/0277-5212(2006)26[258:eoiwjm]2.0.co;2)
66. Khuroo AA, Mehraj G, Muzafar I, Rashid I, Dar GH. Biodiversity conservation in Jammu and Kashmir state: current status and future challenges. *Biodiversity of the Himalaya: Jammu and Kashmir State*. 2020:1049-76.https://doi.org/10.1007/978-981-32-9174-4_41
67. Shrestha B, Shrestha S, Shrestha A, Khadka UR. Ramsar sites in Nepal: Conservation, present scenario, biodiversity value and threats. *J of Wetlands Ecol*. 2020;15.<https://doi.org/10.3126/jowe.v20i0.24782>
68. Saikia S, Das D. Diversity and productivity (Chlorophyll-a and Biomass) of periphyton on natural and artificial substrates from wetland ecosystem. *J of Wetlands Ecol*. 2011;5:1-9.<https://doi.org/10.3126/jowe.v5i0.4624>
69. Verhoeven JT. Wetlands in Europe: perspectives for restoration of a lost paradise. *Ecol Engineer*. 2014;66:6-9.<https://doi.org/10.1016/j.ecoleng.2013.03.006>
70. Willoughby N, Grimble R, Ellenbroek W, Danso E, Amatekpor J. The wise use of wetlands: identifying development options for Ghana's coastal Ramsar sites. *Hydrobiologia*. 2001;458:221-34.
71. Fianko JR, Dodd HS. Sustainable management of wetlands: a case study of the Sogor Ramsar and Unesco man and biosphere reserve in Ghana. *J of Wetlands Environ Management*. 2018;6(1):45-53.<https://doi.org/10.20527/jwem.v6i1.173>
72. Adame MF, Arthington AH, Waltham N, Hasan S, Selles A, Ronan M. Managing threats and restoring wetlands within catchments of the Great Barrier Reef, Australia. *Aquatic Conservation: Marine and Freshwater Ecosystems*. 2019;29(5):829-39.<https://doi.org/10.1002/aqc.3096>
73. Liu S, Du Q, Tong X, Samat A, Pan H, Ma X. Band selection-based dimensionality reduction for change detection in multi-temporal hyperspectral images. *Remote Sensing*. 2017;9(10):1008.<https://doi.org/10.3390/rs9101008>
74. Lagacherie P. Digital soil mapping: a state of the art. *Digital Soil Mapping with Limited Data*. 2008:3-14.
75. Frost T, Austin G, Calbrade N, Mellan H, Hearn R, Robinson A, et al. Waterbirds in the UK 2017/18: The wetland bird survey. *Thetford: BTO, RSPB and JNCC, in association with WWT*; 2019.
76. Guillemain M, Fritz H, Duncan P. The importance of protected areas as nocturnal feeding grounds for dabbling ducks wintering in western France. *Biol Conserv*. 2002;103(2):183-98.[https://doi.org/10.1016/s0006-3207\(01\)00120-3](https://doi.org/10.1016/s0006-3207(01)00120-3)
77. Novo FG, Santillana JT, Granado-Lorencio C. The state of water ecosystems. *Water Policy in Spain: CRC Press*; 2009. p. 41-48.<https://doi.org/10.1201/9780203866023-11>
78. Müller G, Stelzer K, Smollich S, Gade M, Adolph W, Melchionna S, et al. Remotely sensing the German Wadden sea—A new

- approach to address national and international environmental legislation. *Environ Monitoring and Assess.* 2016;188:1-17.<https://doi.org/10.1007/s10661-016-5591-x>
79. Kleijn D, Cherkakoui I, Goedhart PW, van der Hout J, Lammertsma D. Waterbirds increase more rapidly in Ramsar-designated wetlands than in unprotected wetlands. *J of Applied Ecol.* 2014;51(2):289-98.<https://doi.org/10.1111/1365-2664.12193>
 80. Grip K, Blomqvist S. Establishing marine protected areas in Sweden: Internal resistance versus global influence. *Ambio.* 2018;47:1-14.<https://doi.org/10.1007/s13280-017-0932-8>
 81. Gerakis A, Kalburtji K. Agricultural activities affecting the functions and values of Ramsar wetland sites of Greece. *Agri Ecosystems and Environ.* 1998;70(2-3):119-28.[https://doi.org/10.1016/s0167-8809\(98\)00119-4](https://doi.org/10.1016/s0167-8809(98)00119-4)
 82. Wang J, Chen J, Wen Y, Fan W, Liu Q, Tarolli P. Monitoring the coastal wetlands dynamics in Northeast Italy from 1984 to 2016. *Ecological Indicators.* 2021;129:107906.<https://doi.org/10.1016/j.ecolind.2021.107906>
 83. Pijanowska J. The state of nature in Poland: values, protection, challenges and threats. Renata Siuda-Ambroziak Preface 5 Waldemar Koziol, Renata Siuda-Ambroziak, Mariola Zalewska Poland-Brazil: A general comparative statistical overview of SDG achievements.
 84. Bonells M, Zavagli M. National Ramsar/wetlands committees across the six Ramsar regions: diversity and benefits. *Journal of International Wildlife Law and Policy.* 2011;14(3-4):261-92.<https://doi.org/10.1080/13880292.2011.626718>
 85. Poston B, Hyslop C. Canada's commitment to the "Ramsar" convention. *Blue Jay.* 1987;45(4).<https://doi.org/10.29173/bluejay4750>
 86. Gardner RC, Connolly KD. The Ramsar convention on wetlands: assessment of international designations within the United States. *Envtl L Rep News and Analysis.* 2007;37:10089.
 87. Wittmann F, Householder E, de Oliveira Wittmann A, Lopes A, Junk WJ, Piedade MT. Implementation of the Ramsar convention on South American wetlands: an update. *Res and Reports in Biodivers Stud.* 2015;47-58.<https://doi.org/10.2147/rrebs.s64502>
 88. Seco Pon JP, Hernandez MM, Zumpano F, Castano MV, Favero M, Garcia GO. Marine debris accumulation in the Mar Chiquita coastal Lagoon (Biosphere Reserve, Mab-Unesco), a unique wetland in northern Argentina.<https://doi.org/10.2139/ssrn.4268963>
 89. Marín V, Delgado L, Tironi-Silva A, Finlayson C. Exploring social-ecological complexities of wetlands of international importance (Ramsar sites): the Carlos Anwandter Sanctuary (Valdivia, Chile) as a case study. *Wetlands.* 2018;38:1171-82.<https://doi.org/10.1007/s13157-017-0935-z>
 90. Caballero-Gallardo K, Olivero-Verbel J, Corada-Fernández C, Lara-Martín PA, Juan-García A. Emerging contaminants and priority substances in marine sediments from Cartagena Bay and the Grand Marsh of Santa Marta (Ramsar site), Colombia. *Environ Monitoring and Assess.* 2021;193:1-18.<https://doi.org/10.1007/s10661-021-09392-5>
 91. Romero-Mariscal G, Garcia-Chevesich PA, Morales-Paredes L, Arenazas-Rodriguez A, Ticona-Quea J, Vanzin G, et al. Peruvian wetlands: national survey, diagnosis and further steps toward their protection. *Sustain.* 2023;15(10):8255.<https://doi.org/10.3390/su15108255>
 92. Mao D, Wang Z, Wang Y, Choi CY, Jia M, Jackson M, et al. Remote observations in China's Ramsar sites: Wetland dynamics, anthropogenic threats and implications for sustainable development goals. *J of Remote Sens.* 2021.<https://doi.org/10.34133/2021/9849343>
 93. Biswas RM, Nag S, Halder S, Kumar RP. Assessment of wetland potential and bibliometric review: a critical analysis of the Ramsar sites of India. *Bull of the Nat Res Centre.* 2022;46(1):59.<https://doi.org/10.1186/s42269-022-00740-0>
 94. Halls A, editor. Wetlands, biodiversity and the Ramsar convention: the role of the convention on wetlands in the conservation and wise use of biodiversity. Ramsar convention Bureau, Gland, Switzerland; 1997.
 95. Brochet AL, Jbour S, Sheldon RD, Porter R, Jones VR, Al Fazari W, et al. A preliminary assessment of the scope and scale of illegal killing and taking of wild birds in the Arabian peninsula, Iran and Iraq. *Sandgrouse.* 2019;41:154-75.
 96. Mijan US, Rafiqul HA, Abdullah SA. The changing landscape of mangroves in Bangladesh compared to four other countries in tropical regions. *J of Forestry Res.* 2014;25:605-11.<https://doi.org/10.1007/s11676-014-0448-z>
 97. Fletcher S, Kawabe M, Rewhorn S. Wetland conservation and sustainable coastal governance in Japan and England. *Marine Poll Bull.* 2011;62(5):956-62.<https://doi.org/10.1016/j.marpolbul.2011.02.048>
 98. Paudel N, Adhikari S, Paudel G. Ramsar lakes in the foothills of Himalaya, Pokhara-Lekhnath, Nepal: An overview. *Janapriya J of Interdis Stud.* 2017;6:134-47.<https://doi.org/10.3126/jjis.v6i0.19315>
 99. Aslam RW, Shu H, Naz I, Qudoods A, Yaseen A, Gulshad K, et al. Machine learning-based wetland vulnerability assessment in the Sindh Province Ramsar site using remote sensing data. *Remote Sens.* 2024;16(5):928.<https://doi.org/10.3390/rs16050928>
 100. Mucina L, Rutherford MC, Powrie LW, Gerber J, Bezuidenhout H, Sieben E, et al. Inland azonal vegetation. The vegetation of South Africa, Lesotho and Swaziland. 2006;19:630-31.<https://doi.org/10.4102/abc.v47i1.2223>
 101. Morara G, Omondi R, Obegi B, Getabu A, Njiru J, Rindoria N. Water level fluctuations and fish yield variations in Lake Naivasha, Kenya: The trends and relationship. *J of Fisheries and Environ.* 2022;46(1).
 102. Behangana M, Magala R, Katumba R, Ochanda D, Kigoolo S, Mutebi S, et al. Herpetofaunal diversity and community structure in the Murchison Falls-Albert Delta Ramsar site, Uganda: Herpetofaunal diversity. *European J of Ecol.* 2020;6(2).<https://doi.org/10.17161/eurojcol.v6i2.13792>
 103. Abd El-Hamid HT, Toubar MM, Zarzoura F, El-Alfy MA. Ecosystem services based on land use/cover and socio-economic factors in Lake Burullus, a Ramsar site, Egypt. *Remote Sens Appli Soc Environ.* 2023;30:100979.<https://doi.org/10.1016/j.rsase.2023.100979>
 104. Ayanlade A, Proske U. Assessing wetland degradation and loss of ecosystem services in the Niger Delta, Nigeria. *Marine and Freshwater Res.* 2015;67(6):828-36.<https://doi.org/10.1071/mf15066>
 105. Watson RT, René de Roland L, Rabearivony J, Thorstrom R. Community-based wetland conservation protects endangered species in Madagascar: Lessons from science and conservation. *Banwa.* 2007;4(1):83-97.
 106. Wondefrash M, Abebe Y, Geheb K. Wetlands, birds and important bird areas in Ethiopia. *Wetlands of Ethiopia.* 2003:25.
 107. Asbridge E, Lucas RM. Mangrove response to environmental change in Kakadu National Park. *IEEE J of Selected Topics in App Earth Observ and Remote Sens.* 2016;9(12):5612-20.
 108. Denyer K, Robertson H. National guidelines for the assessment of potential Ramsar wetlands in New Zealand. Department of Conservation, Wellington; 2016. p. 58.
 109. Lynch AJJ, Kalumanga E, Ospina GA. Socio-ecological aspects of sustaining Ramsar wetlands in three biodiverse developing countries. *Marine and Freshwater Res* 2016;67(6):850-68.<https://doi.org/10.1071/mf15419>
 110. Long KE, Schneider L, Connor SE, Shulmeister N, Finn J, Roberts GL, et al. Human impacts and anthropocene environmental change at Lake Kutubu, a Ramsar wetland in Papua New

- Guinea. *Proceed of the Nat Acad of Sci.* 2021;118(40):e2022216118. <https://doi.org/10.1073/pnas.2022216118>
111. Sabesh R. Kolleru lake–Andhra Pradesh. *Eco News.* 2010;15(4):6-9.
 112. Bhattacharjya B, Saud B, Borah S, Saikia P, Das B. Status of biodiversity and limno-chemistry of Deepor Beel, a Ramsar site of international importance: Conservation needs and the way forward. *Aquatic Ecosystem Health and Manage.* 2021;24(4):64-74.
 113. Aslam A, Parthasarathy P, Ranjan RK. Ecological and societal importance of wetlands: a case study of North Bihar (India). *Wetlands Conserv: Curr Challen and Fut Strat.* 2021:55-86.
 114. Bisht S, Bhattacharya TR. Comparative analysis of potential Ramsar wetlands in coastal India based on biodiversity and ecological parameters. *Environ Resour Res.* 2024;12(1):147-66.
 115. Mehta P, Chandravanshi S, Yadav SR, Singh A, Jatav SK. Conservation and management of Ramsar sites in India: challenges and solutions.
 116. Solanki V, Joshi A. Disappearing wetland: a study of Basai wetlands, Haryana (India). *Inter J Eco Res.* 2017;14(21):159-69.
 117. Solanki V, Joshi A. Disappearing wetland: a study of Basai wetlands, Haryana (India). *Inter J Eco Res.* 2017;14(21):159-69.
 118. Bera B. 15 Ramsar wetlands in India: Its type and importance.
 119. Shivappa M, Murthy GP. Diversity, density and conservation strategies of bd5 in Ranganathittu, India: a historical Sanctuary of India.
 120. Nandakrishnan, Prasad PRC. Geoscape characterization of Ashtamudi, Sasthamkotta and Vembanad Ramsar sites in Kerala, India. *Water Resour.* 2024;51(4):462-74. <https://doi.org/10.1134/s009780782360273x>
 121. Bhat H, Mahapatra DM, Boominathan M, Rao S, Ramachandra T. Avian diversity of Ladakh wetlands. *Energy and Wetlands Research Group, Centre for Ecological Sciences, Indian Institute of Science: Bangalore, India.* 2010:1-5.
 122. Banerjee D, Bharti D, Kumar S, Mitra A, Joshi R, Gupta D. Faunal composition of Ramsar wetlands from India: An analysis. *Records of the Zoological Survey of India.* 2023:01-15. <https://doi.org/10.26515/rzsi/v123/i1s/2023/172458>
 123. Samant J. Wetland conservation in Maharashtra: need, threats and potential. *Development Research Awareness and Action Institute, Raai, Kolhapur;* 2002.
 124. Roy R, Majumder M. Assessment of water quality trends in Loktak Lake, Manipur, India. *Environ Earth Sci.* 2019;78(13):383. <https://doi.org/10.1007/s12665-019-8383-0>
 125. Nohro S, Jayakumar S. Tree species diversity and composition of the Pala Wetland Reserve forest, Mizoram, Indo-Burma hotspot, India. *Biocatand Agri Biotech.* 2020;23:101474. <https://doi.org/10.1016/j.bcab.2019.101474>
 126. Jana S, Mohanty WK, Gupta S, Kumar P. An integrated geomorphological and geophysical study of neotectonic activity: Analysis of heavy siltation in the Chilka Lake of Odisha, India. *J of Earth System Sci.* 2021;130:1-20. <https://doi.org/10.1007/s12040-021-01702-2>
 127. Solanki V. Ramsar wetlands of India-importance and conservation.
 128. Kulshreshtha S, Sharma B, Sharma S. The Ramsar sites of Rajasthan: Ecology and conservation of Sambhar Salt Lake, Jaipur and Keoladeo National Park, Bharatpur. *Faunal Heritage of Rajasthan, India: Conservation and Management of Vertebrates: Springer;* 2013. p. 173-219. https://doi.org/10.1007/978-3-319-01345-9_8
 129. Nagarajan VM, Yuvar M, Srinivasan R, Satagopan NR, Asokan A, Anooja A. Status of important coastal habitats of North Tamil Nadu: Diversity, current threats and approaches for conservation. *Reg Stud in Marine Sci.* 2022;49:102106. <https://doi.org/10.1016/j.rsma.2021.102106>
 130. Bharati H, Deshmukhe G, Das SK, Kandpal BK, Sahoo L, Bhusan S, et al. Phytoplankton communities in Rudrasagar Lake, Tripura (North-East India)–A Ramsar Site. *Inter J Bio-res Stress Manage.* 2020;11(Feb, 1):001-7.
 131. Festival DVSB. Asan conservation reserve; 2016.
 132. Chakraborty SK, Sanyal P, Ray R. Conclusion on eco-biological uniqueness of wetland ecosystem with special reference with East Kolkata wetlands, India. *Wetlands Ecology: Eco-biological uniqueness of a Ramsar site (East Kolkata Wetlands, India): Springer;* 2023. p. 679-705.