



RESEARCH ARTICLE

Effectiveness of PlantbiotiX formulations as biological solutions for root-knot nematode in bitter gourd

Viraja G¹, V Rajasree^{2*}, Anita B³, Virata G⁴, C Indu Rani¹, M Suganthi⁵ & Prasad Pawar⁶

¹Department of Vegetable Science, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

²Department of Plantation and Spices Crops, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

³Department of Nematology, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

⁴Department of Plantation, Spices, Medicinal & Aromatic Crops, College of Agriculture, Kerala Agricultural University, Vellanikkara 680 656, Kerala, India

⁵Nammazhvar Organic Farming Research Centre, Coimbatore 641 003, Tamil Nadu, India

⁶Zytech Biotech Private Limited, Mumbai 400 059, Maharashtra, India

*Email: dr.rajashreeprabhu@gmail.com



ARTICLE HISTORY

Received: 27 September 2024

Accepted: 01 November 2024

Available online

Version 1.0 : 19 December 2024

Version 2.0 : 29 January 2025



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Viraja G, Rajasree V, Anita. B, Virata. G, Indu Rani C, Suganthi M, Pawar P. Effectiveness of PlantbiotiX formulations as biological solutions for root-knot nematodes in bitter gourd. Plant Science Today.2024;11(sp4):01-09. <https://doi.org/10.14719/pst.5318>

Abstract

The root-knot nematode (*Meloidogyne incognita*) is a major pest that adversely affects bitter gourd production in India. Conventional chemical control methods, though effective, raise economic and environmental concerns, making it necessary to adopt integrated nematode management (INM) strategies. This study evaluates the bio-efficacy of PlantbiotiX formulations, including *Bacillus paralicheniformis* 5% WP ZBM5, *Bacillus subtilis* 2 % SP ZB87 ½, and Xplorer Glory (vesicular arbuscular mycorrhiza), for controlling root-knot nematodes in bitter gourd. Pot culture and field experiments were performed using various concentrations and application methods of PlantbiotiX formulations. Results revealed significant improvements in plant growth parameters, including vine length (9.13%), fruit weight (6.27%), number of fruits (25.21%), and yield per plant (36.36%) when *B. paralicheniformis* 5 % WP ZBM5 and *B. subtilis* 2 % SP ZB87 ½ were applied in combination. Furthermore, a substantial reduction in nematode populations was observed: soil nematode populations decreased by 72.9%, female nematodes by 65.01%, and egg mass by 60.59%. Scanning Electron Microscope (SEM) analysis of treated roots confirmed biofilm formation by *B. paralicheniformis*, demonstrating its capability to mitigate nematode infestation. This biofilm likely played a role in reducing nematode penetration and survival in the roots. The study indicates that PlantbiotiX formulations serve as a sustainable and environmentally friendly substitute for chemical nematicides, promoting plant growth and efficiently controlling root-knot nematodes, thereby presenting a viable solution for enhancing bitter gourd cultivation while reducing environmental harm.

Keywords

biological control; bitter gourd; management; root-knot nematode

Introduction

Bitter gourd (*Momordica charantia*) is an essential vegetable frequently eaten for its remarkable medicinal qualities. Despite being rich in essential nutrients, this vegetable is often underutilized due to its bitter taste. Bitter gourd contains active phytochemicals, including triterpenes, proteins, and steroids, and is rich in minerals such as copper (Cu), iron (Fe), magnesium (Mg), zinc (Zn), and calcium (Ca). It also contains fatty acids like lauric, myristic, palmitic, stearic, and linoleic acids. Charantin, a compound with

hypoglycemic properties found in this plant, aids in regulating blood glucose levels (1). Bitter gourd fruits are rich in iron and vitamin C, exhibiting significant antioxidant activity (2). It is also well known as a source of anti-diabetic drugs, in the pharmaceutical industry (3), and research is being conducted to enhance bitter gourd applications in areas such as colon cancer prevention and obesity management (4). The fruits and seeds contain over 60 phytochemicals that offer protection against more than 30 diseases, including cancer and diabetes (5).

Various biotic factors, such as fungi, bacteria, insect pests, and nematodes, affect bitter gourd production. Root-knot nematodes (*Meloidogyne* spp.) are particularly destructive, leading to significant yield losses and symptoms like leaf drying and stunted growth. Their wide host range, aggressive nature, and global presence make them a major cause of crop loss (6). Root-knot nematodes exhibit significant pathogenic potential in bitter gourd (7), resulting in output losses of up to 13.50 percent, equivalent to around Rs. 252.82 crores (8).

Chemical insecticides are widely used in India to increase yield, but they negatively affected the economy and environment of the nation (9). To reduce the harmful effect of chemical insecticides, INM and biological disease control strategies are gaining popularity, focusing on enhancing rhizosphere biodiversity with organic amendments and biocontrol agents. Biocontrol agents, such as *Purpureocillium lilacinum*, *Pseudomonas fluorescens*, *Trichoderma viride*, and *Pochonia chlamydosporia* have shown promising potential in managing root-knot nematode (10). In recent years, various plant growth-promoting rhizobacteria (PGPR), such as *Bacillus*, *Pseudomonas*, etc., have demonstrated antagonistic effects on plant pathogens (11). Various rhizobacteria produce biofilms in the rhizosphere, acting as biocontrol agents (12). Moreover, endosphere-forming *Bacillus* strains produce lipopeptides like surfactins, iturins, fengycins, plipastatins, etc., which act as broad-spectrum bio-control agents (13). This study aimed to evaluate the bio-efficacy and mode of action of PlantbiotiX formulations, including *B. paralicheniformis* 5 % WP ZBM5, *B. subtilis* 2 % SP ZB87 ½, and Xplorer Glory, as biocontrol options against root-knot nematode, *M. incognita*, in bitter gourd.

Materials and Methods

Experimental location

An experiment was conducted using PlantbiotiX formulations, namely *B. paralicheniformis* 5% WP formulation - ZBM5 (strain no. NAIMCC-SB-0043), *B. subtilis* 2 % SP ZB87 ½, and Xplorer Glory (Vesicular Arbuscular Mycorrhiza), to study the mode of action of PlantbiotiX formulations against root-knot nematodes. This experiment was carried out at the Department of Nematology, Glass House, TNAU Coimbatore.

Experiment details

Hybrid Palee was selected to assess the efficacy of the *B. paralicheniformis* 5 % WP ZBM5 formulation, applied *via* both soil and seed treatment. Seed treatments were conducted at concentrations of 2.5, 5.0, and 7.5g/Kg seeds. Soil applications were performed at concentrations of 1.0, 2.0, and 4.0 g /L of water.

Pot culture studies

Pots were disinfected and filled with sterilized soil. Plants were inoculated with root-knot nematodes (2J2 /g soil) 15 days after sowing. Soil application of the formulation was done at three concentration levels (1.0, 2.0, and 4.0 g /L of water), while seed treatment was conducted at concentrations of 2.5, 5.0, and 7.5 g / Kg of seed. Observations were recorded 45 days after plant emergence, focusing on vine length (cm), root length (cm), and nematode populations. Roots were washed, stained with 0.1% acid fuchsin, and then left in lactophenol for at least a day to discolor. Galls and egg masses per plant were counted using a stereoscopic binocular microscope. Soil from each treatment was thoroughly mixed, and 200 cc was taken to estimate nematode populations using Cobb's sieving and decanting technique, followed by Baermann's funnel method.

Scanning Electron Microscopy

The SEM utilized for analysis was the FEI Quanta 250, developed by the Field Electron and Ion Company, Czech Republic. This model is equipped with an Everhart-Thornley Detector (ETD), designed to collect secondary electrons for creating high-resolution images. The SEM operates with a tungsten electron source, which provides a stable electron required for imaging. Root samples were initially sputter-coated and then spread on double-sided conductive carbon tape, fixed onto a stub, and placed in the SEM sample chamber. After attaining a high vacuum pressure of 3.99e-4 Pa, the filament was turned on, and various parameters such as electron beam intensity, spot size, voltage, and emission current were adjusted. A magnification of 7220× provided detailed imaging of the microstructure of the samples. Images were then captured and depicted.

Field studies

The study was conducted using a completely randomized block design with seven treatments and five replications. *B. paralicheniformis* 5 % WP ZBM5 was applied both separately and in combination with *B. subtilis* 2 % SP ZB87 ½ and Xplorer Glory and was compared with a chemical nematicide (Fluopyrum 34.48% SC) and a bio-nematicide (*P. chlamydosporia*). Initial soil testing was done to evaluate the nematode populations. Fields were plowed, and pits were dug at 2×1 m spacing. Two to three hybrid Palee seeds were planted per pit and thinned to one plant after 15 days. The first treatment was applied 15 days after sowing (DAS), followed by additional treatments on the 30th and 45th day. Recommended practices were maintained throughout. Nematode populations and growth parameters were observed.

Statistical Analysis

The glasshouse experiment was conducted using a completely randomized block design with three replications, while the field experiments were carried out in a randomized block design with five replications to study the bioefficacy of PlantbiotiX products. Statistical significance was assessed using the F-value at the 5% level, along with the critical difference (CD). AGRESS software was used to perform the statistical analysis.

Results

Mode of action of PlantbiotiX products against *M. incognita* in bitter gourd: Pot culture studies

nita in bitter gourd: Pot culture studies

A pot culture experiment was conducted to study the mode of action of PlantbiotiX formulations against *M. incognita* in bitter gourd. Among the different methods, seed treatment performed better than soil application. The longest vine length was recorded in T3 (*B. paralicheniformis* 5 % WP ZBM5, seed treatment at 7.5g/kg seed) with a 12.10% increase over the control, followed by T6 (*B. paralicheniformis* 5 % WP ZBM5, soil application at 4g/kg) with a 10.36% increase over the control. The maximum root length was recorded in T3, showing a 12.99% increase over the control, followed by T6 with a 10.56 % increase (Table 1 & Figs. 1 - 3). The maximum reduction in soil nematode

Table 1. Effect of PlantbiotiX formulations on growth and nematode population of bitter gourd (Palee): Pot culture studies

Treatments	Vine length (cm)	Root length (cm)	Soil nematode population	Root gall index	Number of female nematodes /5g of roots (30 days)	Number of egg masses/ 5g of roots (30 days)
T ₁ - <i>B. paralicheniformis</i> 5 % WP ZBM5- seed treatment-2.5g/kg seed	178.15	8.79	156.10	3	9.8	8.1
T ₂ - <i>B. paralicheniformis</i> 5 % WP ZBM5- seed treatment-5g/kg seed	180.46	9.01	145.97	2	8.1	7.6
T ₃ - <i>B. paralicheniformis</i> 5 % WP ZBM5- seed treatment-7.5g/kg seed	190.64	9.74	95.97	1	7.0	5.7
T ₄ - <i>B. paralicheniformis</i> 5 % WP ZBM5- soil application-1g/ L	176.43	8.71	170.15	3	9.8	9.2
T ₅ - <i>B. paralicheniformis</i> 5 % WP ZBM5- soil application-2 g/ L	182.25	8.99	166.44	3	9.3	8.7
T ₆ - <i>B. paralicheniformis</i> 5 % WP ZBM5- soil application-4g/ L	186.93	9.53	124.43	2	7.9	6.5
T ₇ - Sterile soil without nematodes	173.21	8.7	0.00	0	0	0
T ₈ - Sterile soil with nematodes	167.56	8.6	345.90	4	17.4	12.6
SE(d)	4.24	0.15	4.25		0.18	0.18
CD (5%)	8.99	0.31	9.01		0.39	0.39

Sed: Standard error of deviation; **CD:** Critical difference. Source: AGRESS Software



Fig. 1. Effect of PlantbiotiX formulations on plant growth of bitter gourd (Palee): pot culture studies. **A:** T3 plant sample; **B:** T8 plant sample.



Fig. 2. Effect of PlantbiotiX formulations on roots of bitter gourd (Palee): Pot culture studies. **A:** T3 plant sample; **B:** T8 plant sample.




Fig. 3. Effect of PlantbiotiX formulations on vine length and root length of bitter gourd (Palee): Pot culture studies.

population (72.25%), female nematode population (59.77%), and egg mass (54.76%) was observed in T3, showing a significant reduction over the control. This was

followed by T6, with decreases of 64.03%, 54.60%, and 48.41%, respectively, compared to the untreated control (Table 1 & Fig. 4).

Scanning electron microscopic analysis

The surface morphology was closely examined using SEM images. Root samples from T6 (*B. paralicheniformis* 5 % WP ZBM5, soil application at 4g/kg) and T8 (sterile with

colonization patterns of *Bacillus amyloliquefaciens* FZB42 on the roots of *Arabidopsis thaliana* was previously examined using a SEM (14). The potential of biofilm formation by biological control agents and their role in effective pest management strategies is further supported by another




Fig. 4. Effect of PlantbiotiX formulations on soil nematode population, female nematodes, and egg masses: Pot culture studies.

nematodes) were observed at various magnification levels. The root sample from T6 (*B. paralicheniformis* 5 % WP ZBM5) showed root colonization, with *B. paralicheniformis* forming a biofilm on the root surface. This biofilm prevented the action of *M. incognita*, preventing further infection. In contrast, the root sample from T8 (sterile with nematodes) showed high infection, with root nematodes forming gall symptoms (Fig. 5). The biofilm formation and

investigation (15).

Bioefficacy of PlantbiotiX products against *M. incognita* in bitter gourd at field condition

Plant growth parameters

The longest vine length among the PlantbiotiX formulations was recorded in T3 *B. paralicheniformis* 5 % WP




Fig. 5 Effects of PlantbiotiX formulations on root samples of bitter gourd (Palee). A: T3 root sample showing biofilm formation of *B. paralicheniformis*, imaged at a magnification of 7220 \times ; B: T8 control without biofilm formation, imaged at a magnification of 7220 \times .

ZBM5, *B. subtilis* 2% SP ZB87 ½ (2 times each), showing a 7.84% increase over the control, followed by T4 (*B. paralicheniformis* 5% WP ZBM5, Xplorer Glory, applied 2 times each), with a 4.98% increase. The highest fruit weight was observed in T3 (*B. paralicheniformis* 5 % WP ZBM5, *B. subtilis* 2 % SP ZB87 ½ , applied 2 times each), with a 5.62% increase over control, followed by T4 (*B. paralicheniformis* 5 % WP ZBM5, Xplorer Glory, applied 2 times each), with a 3.10% increase. The maximum number of fruits was recorded in T3(*B. paralicheniformis* 5 % WP ZBM5, *B. subtilis* 2 % SP ZB87 ½, applied 2 times each), with a 25.21% increase over the control, followed by T4 (*B. paralicheniformis* 5 % WP ZBM5, Xplorer Glory, applied 2 times each), showing a 17.58% increase. The highest yield per plant was recorded in T3 (*B. paralicheniformis* 5 % WP ZBM5, *B. subtilis* 2 % SP ZB87 ½, applied 2 times each), with a 36.36% increase over control, followed by T4 (*B. paralicheniformis* 5 % WP ZBM5, Xplorer Glory, applied 2 times each), with a 24.32% increase. All the observations on growth parameters were comparable to the chemical control (Table 2).

Nematode population at 30 DAT

The maximum reduction among the PlantbiotiX formulations was observed with the combination of *B. paralicheniformis* 5 % WP ZBM5 and *B. subtilis* 2 % SP ZB87 ½ (2 times each), resulting in a decrease in the soil nematode popula-

ble to the chemical control (Fluopyrum 34.48% SC). This was followed by *B. paralicheniformis* 5 % WP ZBM5 and Xplorer Glory (2 times each), which reduced the soil nematode population by 57.3%, *B. paralicheniformis* 5 % WP ZBM5 (3 times), which showed a 55.07% reduction in the female nematode population, and *P. chlamydosporia*, which reduced egg mass by 35.99% compared to the untreated control (Table 3).

Nematode population at harvest

The maximum reduction among the PlantbiotiX formulations was observed with the combination of *B. paralicheniformis* 5 % WP ZBM5 and *B. subtilis* 2 % SP ZB87 ½ (2 times each), which resulted in a decrease in the soil nematode population (72.9%), female nematode population (65.01%), and egg mass (60.59%) compared to the untreated control. This combinations outperformed the chemical control (Fluopyrum 34.48% SC). It was followed by *B. paralicheniformis* 5 % WP ZBM5 and Xplorer Glory (2 times each), which reduced the soil nematode population by 57.3%, and *B. paralicheniformis* 5% WP ZBM5 (applied three times), which showed a 57.72% reduction in the female nematode population. *P. chlamydosporia* also showed a reduction in egg mass (32.87%) compared to the untreated control (Table 3).

Table 2. Effect of PlantbiotiX formulation on plant growth and yield of bitter gourd (Palee) under field conditions

Treatments	Vine length (cm)	Percent increase over control	Fruit weight (g)	Percent increase over control	No. of fruits per plant	Percent increase over control	Yield per plant (kg)	Percent increase over control
T ₁ <i>B. paralicheniformis</i> 5 % WP ZBM5 200g/acre (2 times)	581.48	2.32	153.31	1.86	25.95	14.17	2.73	17.00
T ₂ <i>B. paralicheniformis</i> 5 % WP ZBM5 200g/acre (3 times)	588.39	3.46	154.79	2.79	25.17	11.54	2.77	18.08
T ₃ <i>B. paralicheniformis</i> 5 % WP ZBM5 200g/acre, <i>B. subtilis</i> 2 % SP ZB87 ½ 500g/acre (2 times each)	616.34	7.84	159.42	5.62	29.78	25.21	3.52	35.66
T ₄ <i>B. paralicheniformis</i> 5 % WP ZBM5 200g/acre. Xplorer Glory 100g/acre (2 times each)	597.76	4.98	155.29	3.10	27.02	17.58	2.98	23.96
T ₅ Fluopyrum 34.48% SC 1ml /L	625.07	9.13	160.54	6.27	28.72	22.46	3.31	31.62
T ₆ <i>P. chlamydosporia</i> 2kg/ha	579.73	2.02	154.45	2.58	25.05	11.10	2.94	22.82
T ₇ Non treated	568.01	0.00	150.46	0.00	22.27	0.00	2.27	0.00
SE.d	6.48		1.79		0.24		0.11	
CD (5%)	13.16		3.64		0.49		0.23	

Sed: Standard error of deviation; **CD:** Critical difference. Source: AGRESS Software

tion (68.7%), female nematode population (48.99%), and egg mass (60.39%) compared to the untreated control, with results comparable to the chemical control (Fluopyrum 34.48% SC). This was followed by *B. paralicheniformis* 5 % WP ZBM5 and Xplorer Glory (2 times each), which showed reductions of 54.1%, 39.52%, and 31.55%, respectively, compared to the untreated control (Table 3).

Nematode population at 60 DAT

The maximum reduction among the PlantbiotiX formulations was observed with *B. paralicheniformis* 5 % WP ZBM5 and *B. subtilis* 2 % SP ZB87 ½ (2 times each), resulting in a decrease in the soil nematode population (70.7%), female nematode population (61.16%), and egg mass (60.36%) compared to the untreated control, with results compara-

Discussion

The beneficial effects of *B. paralicheniformis* and *B. subtilis* are well-documented in multiple studies on various plants. For instance, *B. paralicheniformis* TRQ65, isolated from wheat fields in Mexico, contains genes linked to stress response and growth-promoting substances (16). The *B. paralicheniformis* RP01 strain promoted growth on the root surface of *Brassica chinensis*, significantly increasing root length and brassinosteroid (BR) content (17). Similarly, (18) *B. paralicheniformis* SX21 was observed to significantly enhance root structures, photosynthetic parameters, growth rate, and biomass in cucumber. Supporting these findings, inoculation with *B. paralicheniformis* LBEndo1 and *Aeromonas caviae* KBect04 led to increased

Table 3. Effect of PlantbiotiX formulation on root-knot nematode populations at field condition

Treatments	Soil nematode population			Root gall index			Number of females/5g of root			Number of egg masses/ 5g of root		
	DAYS			DAYS			DAYS			DAYS		
	30	60	90	30	60	90	30	60	90	30	60	90
T ₁ <i>B. paralicheniformis</i> 5 % WP ZBM5 200g/acre (2 times)	170.1 (49.9)	164.4 (52.0)	155.6 (55.7)	3	3	3	9.64 (48.1)	9.09 (53.7)	8.93 (56)	9.19 (32.7)	8.96 (30.6)	8.47 (27.6)
T ₂ <i>B. paralicheniformis</i> 5 % WP ZBM5 200g/acre (3 times)	166.4 (51.0)	155.9 (54.4)	147.2 (58.1)	3	3	3	9.34 (49.7)	8.84 (55.0)	8.58 (57.7)	8.85 (35.2)	8.43 (34.6)	8.23 (29.7)
T ₃ <i>B. paralicheniformis</i> 5 % WP ZBM5 200g/acre, <i>B. subtilis</i> 2 % SP ZB87,½ 500g/acre (2 times each)	106.4 (68.7)	100.3 (70.7)	95.1 (72.9)	2	2	2	9.48 (48.9)	7.64 (61.1)	7.10 (65.0)	5.42 (60.3)	5.12 (60.3)	4.61 (60.5)
T ₄ <i>B. paralicheniformis</i> 5 % WP ZBM5 200g/acre, Xplorer Glory 100g/acre (2 times each)	156.1 (54.1)	146.1 (57.3)	132.8 (62.2)	3	3	3	11.24 (39.5)	9.23 (53.1)	8.71 (57.0)	9.36 (31.5)	8.29 (35.7)	7.89 (32.6)
T ₅ Fluopyrum 34.48% SC 1ml /L	86.0 (74.7)	82.5 (75.9)	120.2 (65.8)	1	1	1	6.43 (65.4)	5.62 (71.4)	8.29 (59.1)	4.71 (65.5)	4.44 (65.5)	8.85 (24.4)
T ₆ <i>P. chlamydosporia</i> 2kg/ha	193.0 (43.2)	185.6 (45.8)	149.5 (57.4)	3	3	3	11.64 (37.3)	10.11 (48.6)	9.63 (52.5)	8.63 (36.8)	8.26 (35.9)	7.86 (32.8)
T ₇ Non treated	339.9 (0.0)	342.2 (0.0)	351.1 (0.0)	4	4	4	18.58 (0.00)	19.68 (0.00)	20.29 (0.00)	13.68 (0.00)	12.90 (0.00)	11.71 (0.00)
SE.d	1.70	1.78	2.36				0.13	0.17	0.15	0.12	0.11	0.12
CD (5%)	3.46	3.62	4.79				0.27	0.35	0.30	0.25	0.22	0.25

Sed: Standard error of deviation; **CD:** Critical difference. Source: AGRESS Software

fruit production in tomato plants (19). In line with these observations, our study also found that *B. paralicheniformis* improved plant growth and yield in bitter melon. Moreover, *B. paralicheniformis* MDJK30 was shown to promote plant growth by enhancing of carbohydrate and amino acid metabolism (20).

The inoculation of the *B. paralicheniformis* RP01 strain in cotton plants led to elevated concentrations of growth hormones, specifically brassinosteroids (BRs) and auxin (IAA), with increases of 43.9% and 5.1%, respectively (17). Genomic analysis of this strain identified the trpABCDEF pathway for L-TRP, spermine synthase (speE), and agmatinase (speB), which can catalyze the conversion of amino acids into plant growth-promoting compounds (17). The microbial diversity of the rhizosphere is an important ecological bioindicator for the healthy plant growth (18). After 44 days of inoculation with *B. paralicheniformis* 2R5, the diversity of beneficial microorganisms such as *Nitrospira*, *Ramlibacter*, *Sphingomonas*, *Massilia*, and *Terrimonas* increased in the rhizosphere of canola (21). Genomic analysis of *B. paralicheniformis* BP9 revealed the presence of genes for secondary metabolites such as bacillibactin, lantibiotics, and bacitracin, contributing to its antibacterial potential (22). *B. subtilis* enhances plant growth by improving nutrient uptake through mechanisms such as nitrogen fixation, phosphorus solubilization, and siderophore production (23). Volatile substances synthesized by *B. subtilis* SST2, such as albuterol and 1,3-propanediol, enhance plant development by elevating gibberellin and auxin levels, as well as photosynthetic rates (24). Additionally, *B. subtilis* AH18 and *B. licheniformis* K11 have been shown to promote growth in tomatoes and peppers by producing auxins (IAA, IBA, IPA), antifungal β -glucanase, siderophores, and solubilizing phosphates (25). Complementing this, a study emphasized the effec-

tiveness of using multiple microorganisms together, highlighting the genomic and plant growth-promoting traits of *B. paralicheniformis* ES- 1 (26).

The nematicidal properties of *B. paralicheniformis* and *B. subtilis* have been extensively studied, showcasing their potential in managing plant-parasitic nematodes. *B. paralicheniformis* has demonstrated potential as a biocontrol agent against phytopathogenic nematodes, a degrader of polystyrene, and a plant growth promoter (27). A study found that the secretome of *B. paralicheniformis* TB197 exhibited over 95% nematicidal activity against *M. incognita* and effectively suppressed *M. enterolobii* infections in tomatoes (28). Similarly, it was reported from another study that *B. paralicheniformis* FMCH001 and *B. subtilis* FMCH002 hampered the establishment of nematodes by reducing egg-hatching and juvenile survival (29). This finding supports the notion that *B. paralicheniformis* is effective in inhibiting the life cycle of nematodes, thus providing a viable biocontrol strategy. Furthermore, a comparative study between *B. paralicheniformis* TB197 and *B. subtilis* ATCC 21332 showed that TB197 exhibited greater nematicidal action (30). The nematicidal activity of *B. paralicheniformis* TB197 is primarily attributed to its high polarity, which disrupts the cell membranes of *Caenorhabditis elegans* larvae, affecting their viability. Mutagenesis experiments in *B. subtilis* OKB105 identified the purL gene as crucial for nematicidal activity (31). In another study, tomato plants treated with bacterial suspensions of *B. subtilis* (MTCC-441) and *Pseudomonas putida* (MTCC-102) before J2 inoculation exhibited enhanced growth characteristics and high nematicidal activity (32).

Gas chromatography-mass spectrometry (GC-MS) analysis identified several nematicidal volatile compounds, like ketones, alkyls, sulfides, and heterocyclic compounds, in *B. subtilis* Bs-1 strain (33). Additionally, the

genomic study of *B. subtilis* GEB5 identified 21 homologs of nematode-virulent proteases, suggesting a potential mechanism for its nematocidal activity (34). This further underscores the superior efficacy of *B. paralicheniformis* and *B. subtilis* in managing nematode populations.

Nonetheless, one drawback of the pot cultivation study is the controlled environmental condition, which may not accurately represent the variability found in wild field situations. Additionally, the study duration was limited to a single growing season, which may not capture the long-term effects of the formulations on nematode activity and plant growth performance.

Conclusion

The study compared PlantbiotiX formulations with a chemical nematicide (Fluopyrum 34.48% SC) and a biological control agent (*P. chlamydosporia*). Fluopyrum 34.48% SC effectively reduced the nematode population; however, its effectiveness declined after 60 days. In contrast, the combination of *B. paralicheniformis* 5 % WP ZBM5 and *B. subtilis* 2 % SP ZB87 ½ (T3) demonstrated sustained efficacy throughout the crop cycle.

Future research could explore the combined use of *B. paralicheniformis* with other biocontrol agents such as *B. subtilis*, *P. chlamydosporia*, *P. lilacinum*, *P. fluorescens*, and *T. viride* to further reduce nematode populations and enhance plant growth in bitter melon. Additionally, the sustainability of *B. paralicheniformis* formulations can be examined through long-term field trials across different regions and environmental conditions. Future studies could also investigate the molecular mechanisms behind the nematocidal activity of *B. paralicheniformis* to identify key bioactive compounds. Further research on formulations that enhance shelf life and stability across various soil types, as well as studies on their synergistic effect with other organic amendments, would be valuable for practical agricultural applications. Moreover, assessing the impact of PlantbiotiX formulations on other economically important crops could expand its potential in INM strategies.

In conclusion, cumulative evidence from these studies suggests that PlantbiotiX formulations hold significant potential for reducing root-knot nematode damage while improving plant growth and yield of bitter melon.

Acknowledgements

The authors would like to extend their sincere thanks to the Department of Vegetable Science for providing the scholarship needed for the successful completion of research. The authors are also grateful to Tamil Nadu Agricultural University for offering the laboratory facilities required for the SEM analysis. Special thanks to Chairperson, Dr V Rajasree, and the advisory committee for their unwavering support throughout the research process and their invaluable assistance in writing the article. The study was funded by Zytex Biotech, Mumbai, and is gratefully acknowledged.

Authors' Contributions

VG¹ conceptualized the study, developed the research methodology, played a key role in data interpretation and contributed to overall writing of the manuscript. VG¹ also carried out the SEM analysis, including sample preparation, data acquisition and responsible for analyzing and interpreting the results. Additionally, VG¹ helped draft the manuscript. AB offered technical assistance during the SEM analysis, aided in data interpretation and contributed to revising the manuscript with a focus on critical intellectual aspects. VG² helped in revising the manuscript and data analysis. CIR reviewed and edited the manuscript, offering valuable insights into the discussion and conclusions. MS conducted the literature review, supported data interpretation and contributed to drafting, revising the manuscript and assisted in the final submission process. PP supported the research financially. (VG¹- Viraja G & VG²- Virata G)

Compliance with Ethical Standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

References

1. Krawinkel MB, Keding GB. Bitter melon (*Momordica charantia*): A dietary approach to hyperglycemia. *Nutr Rev*. 2006;64(7 Pt 1):331-37. <https://doi.org/10.1301/nr.2006.jul.331-337>
2. Behera TK, Behera S, Bharathi LK, John KJ, Simon PW, Staub JE. Bitter melon: Botany, horticulture, breeding. In: Janic J, editor. *Horticultural Reviews*. Wiley-Blackwell;2010.v37.p.101-41. <http://dx.doi.org/10.1002/9780470543672.ch2>
3. Krawinkel MB, Ludwig C, Swai ME, Yang RY, Chun KP, Habicht SD. Bitter melon reduces elevated fasting plasma glucose levels in an intervention study among prediabetics in Tanzania. *J Ethnopharmacol*. 2018;216:1-7. <https://doi.org/10.1016/j.jep.2018.01.016>
4. Kwatra D, Subramaniam D, Ramamoorthy P, Standing D, Moran E, Velayutham R, et al. Methanolic extracts of bitter melon inhibit colon cancer stem cells by affecting energy homeostasis and autophagy. *Evid Based Complement Alternat Med*. 2013;2013(1):702869. <https://doi.org/10.1155/2013/702869>
5. Gayathry KS, John JA. A comprehensive review on bitter melon (*Momordica charantia* L.) as a gold mine of functional bioactive components for therapeutic foods. *Food Production, Processing and Nutrition*. 2022;4(1):10. <https://doi.org/10.1186/s43014-022-00089-x>
6. Khan J, Baheti B, Sharma HK, Bairwa HL, Verma A, Nama C. An eco-friendly management option against root-knot nematode (*Meloidogyne incognita*) infecting bitter melon (*Momordica charantia* L.). *Biol Forum*. 2023;15(12):45-52.
7. Singh SK, Conde B, Hodda M. Root-knot nematode (*Meloidogyne incognita*) on bitter melon (*Momordica charantia*) near Darwin, Australia. *Australasian Plant Dis Notes*. 2012;7:75-78. <http://dx.doi.org/10.1007/s13314-012-0052-z>
8. Kumar V, Khan MR, Walia RK. Crop loss estimations due to plant-parasitic nematodes in major crops in India. *Natl Acad Sci Lett*. 2020;43:409-12. <https://doi.org/10.1007/s40009-020-00895-2>

9. Singh S. Integrated approach for the management of the root-knot nematode, *Meloidogyne incognita*, on eggplant under field conditions. *Nematology*. 2013;15(6):747-57. <https://doi.org/10.1163/15685411-00002715>
10. Mane PB, Mhase N. Bioefficacy of different bioagents against root-knot nematode, *Meloidogyne incognita* infesting bottle gourd under laboratory conditions. *Int J Plant Prot.*2017;10(1):87-91. <http://dx.doi.org/10.15740/HAS/IJPP/10.1/87-91>
11. Mhatre PH, Karthik C, Kadirvelu K, Divya KL, Venkatasalam EP, Srinivasan S, et al. Plant growth promoting rhizobacteria (PGPR): A potential alternative tool for nematodes bio-control. *Biocatal Agric Biotechnol*. 2019;17:119-28. <http://dx.doi.org/10.1016/j.bcab.2018.11.009>
12. Sang MK, Kim KD. Biocontrol activity and root colonization by *Pseudomonas corrugata* strains CCR04 and CCR80 against Phytophthora blight of pepper. *BioControl*. 2014;59:437-48. <http://dx.doi.org/10.1007/s10526-014-9584-9>
13. Ongena M, Jacques P. *Bacillus* lipopeptides: Versatile weapons for plant disease biocontrol. *Trends Microbiol*. 2008;16(3):115-25. <https://doi.org/10.1016/j.tim.2007.12.009>
14. Bhattacharyya A, Mavrodi O, Bhowmik N, Weller D, Thomashow L, Mavrodi D. Bacterial biofilms as an essential component of rhizosphere plant-microbe interactions. In: Gurtler V, Patrauchan M, editors. *Methods in Microbiology*. Elsevier;2023.v53.p. 3-48. <https://doi.org/10.1016%2Fbs.mim.2023.05.006>
15. Keswani C, Singh HB, García-Estrada C, Caradus J, He Y-W, Mezaache-Aichour S, et al. Antimicrobial secondary metabolites from agriculturally important bacteria as next-generation pesticides. *Appl Microbiol Biotechnol*. 2020;104(3):1013-34. <https://doi.org/10.1007/s00253-019-10300-8>
16. Valenzuela-Ruiz V, Robles-Montoya RI, Parra-Cota FI, Santoyo G, Ma del Carmen Orozco-Mosqueda M, Rodríguez-Ramírez R, et al. Draft genome sequence of *B. paralicheniformis* TRQ65, a biological control agent and plant growth-promoting bacterium isolated from wheat (*Triticum turgidum* subsp. *durum*) rhizosphere in the Yaqui Valley, Mexico. *3 Biotech*. 2019;9:436. <https://doi.org/10.1007/s13205-019-1972-5>
17. Xu J, Qin L, Xu X, Shen H, Yang X. *B. paralicheniformis* RP01 enhances the expression of growth-related genes in cotton and promotes plant growth by altering microbiota inside and outside the root. *Int J Mol Sci*. 2023;24(8):7227. <https://doi.org/10.3390/ijms24087227>
18. Wang J, Qu F, Liang J, Yang M, Hu X. *Bacillus velezensis* SX13 promoted cucumber growth and production by accelerating the absorption of nutrients and increasing plant photosynthetic metabolism. *Sci Hortic*. 2022;301:111151. <https://doi.org/10.1016/j.scienta.2022.111151>
19. Palacio-Rodríguez R, Nava-Reyes B, Sánchez-Galván H, Quezada-Rivera JJ, Sáenz-Mata J. Effect of plant growth-promoting rhizobacteria inoculation on tomato under commercial shade-house conditions. *Rev Mexicana Cienc Agric*. 2022;13(SPE28):231-42. <https://doi.org/10.29312/remexca.v13i28.3278>
20. Du Y, Ma J, Yin Z, Liu K, Yao G, Xu W, et al. Comparative genomic analysis of *B. paralicheniformis* MDJK30 with its closely related species reveals an evolutionary relationship between *B. paralicheniformis* and *B. licheniformis*. *BMC Genomics*. 2019;20:283. <https://doi.org/10.1186/s12864-019-5646-9>
21. Świątczak J, Kalwasińska A, Szabó A, Brzezinska MS. The effect of seed bacterization with *B. paralicheniformis* 2R5 on bacterial and fungal communities in the canola rhizosphere. *Microbiol Res.*2023;275:127448. <https://doi.org/10.1016/j.micres.2023.127448>
22. Asif M, Li-Qun Z, Zeng Q, Atiq M, Ahmad K, Tariq A, et al. Comprehensive genomic analysis of *B. paralicheniformis* strain BP9, pan-genomic and genetic basis of biocontrol mechanism. *Comput Struct Biotechnol J*. 2023;21:4647-62. <https://doi.org/10.1016/j.csbj.2023.09.043>
23. Hashem A, Tabassum B, Abd_Allah EF. *Bacillus subtilis*: A plant-growth promoting rhizobacterium that also impacts biotic stress. *Saudi J Biol Sci*. 2019;26(6):1291-97. <https://doi.org/10.1016/j.sjbs.2019.05.004>
24. Tahir HA, Gu Q, Wu H, Raza W, Hanif A, Wu L, et al. Plant growth promotion by volatile organic compounds produced by *Bacillus subtilis* SYST2. *Front Microbiol*. 2017;8:171. <https://doi.org/10.3389%2Ffmicb.2017.00171>
25. Lim J-H, Kim S-D. Synergistic plant growth promotion by the indigenous auxins-producing PGPR *Bacillus subtilis* AH18 and *Bacillus licheniformis* K11. *J Korean Soc. Appl Biol Chem*. 2009;52:531-38. <http://dx.doi.org/10.3839/jksabc.2009.090>
26. Iqbal S, Qasim M, Rahman H, Khan N, Paracha RZ, Bhatti MF, et al. Genome mining, antimicrobial and plant growth-promoting potentials of halotolerant *B. paralicheniformis* ES-1 isolated from salt mine. *Mol Genet Genomics*. 2023;298:79-93. <https://doi.org/10.1007/s00438-022-01964-5>
27. Can-Ubando LC, Ramírez-Durán N, Aranda E, Manzanares-Leal GL, Sánchez-Reyes A, de Paz GA, et al. Complete genome sequence of the *B. paralicheniformis* strain HAS-1. *Microbiol Resour Announc*. 2024;13:e00337-24. <https://doi.org/10.1128/mra.00337-24>
28. Chavarria-Quicaño E, Contreras-Jáquez V, Carrillo-Fasio A, De la Torre-González F, Asaff-Torres A. Native *B. paralicheniformis* isolate as a potential agent for phytopathogenic nematodes control. *Front Microbiol*. 2023;14:1213306. <https://doi.org/10.3389/fmicb.2023.1213306>
29. Díaz-Manzano FE, Amora DX, Martínez-Gómez Á, Moelbak L, Escobar C. Biocontrol of *Meloidogyne* spp. in *Solanum lycopersicum* using a dual combination of *Bacillus* strains. *Front Plant Sci*. 2023;13:1077062. <https://doi.org/10.3389/fpls.2022.1077062>
30. Chavarria-Quicaño E, De la Torre-González F, González-Riojas M, Rodríguez-González J, Asaff-Torres A. Nematicidal lipopeptides from *B. paralicheniformis* and *Bacillus subtilis*: A comparative study. *Appl Microbiol Biotechnol*. 2023;107:1537-49. <https://doi.org/10.1007/s00253-023-12391-w>
31. Xia Y, Xie S, Ma X, Wu H, Wang X, Gao X. The *purL* gene of *Bacillus subtilis* is associated with nematicidal activity. *FEMS Microbiol. Lett*. 2011;322(2):99-107. <https://doi.org/10.1111/j.1574-6968.2011.02336.x>
32. Nadeem H, Niazi P, Asif M, Kaskavalcı G, Ahmad F. Bacterial strains integrated with surfactin molecules of *Bacillus subtilis* MTCC441 enrich nematocidal activity against *Meloidogyne incognita*. *Plant Biol*. 2021;23(6):1027-36. <https://doi.org/10.1111/plb.13301>
33. Cao H, Jiao Y, Yin N, Li Y, Ling J, Mao Z, et al. Analysis of the activity and biological control efficacy of the *Bacillus subtilis* strain Bs-1 against *Meloidogyne incognita*. *Crop Prot*. 2019;122:125-35. <http://dx.doi.org/10.1016/j.cropro.2019.04.021>
34. Ganeshan S, Settu V, Mannu J, Annaiyan S, Muthusamy G, Arun A, et al. Genomic analysis of *Bacillus subtilis* sub sp. *subtilis* GEB5 reveals its genetic assets for nematicidal and plant growth promoting mechanisms. *Rhizosphere*. 2024;31:100953. <http://dx.doi.org/10.1016/j.rhisph.2024.100953>