



REVIEW ARTICLE

Millet bran: The underrated ingredient with a potential to transform human nutrition

Preeti Kittur¹, V Meenakshi^{2*}, R Vijayalakshmi³, ML Mini⁴, A Kalaiselvan⁵ & M Ilamaram¹

¹Department of Food Science and Nutrition, Community Science College and Research Institute, Tamil Nadu Agricultural University, Madurai 625 104, Tamil Nadu, India

²Department of Horticulture, VO Chidhambaranar, Agricultural College and Research Institute Tamil Nadu Agricultural University, Killikulam 628 252, Tamil Nadu, India

³Department of Family Resource Management and Consumer Science, Community Science College and Research Institute, Tamil Nadu Agricultural University, Madurai 625 104, Tamil Nadu, India

⁴Department of Biotechnology, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Madurai 625 104, Tamil Nadu, India

⁵Department of Extension Education and Communication, Community Science College and Research Institute, Tamil Nadu Agricultural University, Madurai 625 104, Tamil Nadu, India

*Email: meenakshi.v@tnau.ac.in

OPEN ACCESS

ARTICLE HISTORY

Received: 01 October 2024

Accepted: 02 November 2024

Available online

Version 1.0 : 29 December 2024



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Kittur P, Meenakshi V, Vijayalakshmi R, Mini ML, Kalaiselvan A, Ilamaram M. Millet bran: The underrated ingredient with a potential to transform human nutrition. Plant Science Today. 2024;11(sp4):01-10. <https://doi.org/10.14719/pst.5431>

Abstract

Millets are consumed by people across the globe. Millet bran (MB), a byproduct of preliminary processes like dehulling, debranning and milling, is often discarded or utilized as animal feed. Foxtail millet bran (FMB) consists of 9.39% crude oil, 12.48% crude protein, 51.69% crude fiber, 7.50% ash, and 8.29% moisture. Kodo millet bran (KMB) has a nutrient profile of 4.92% protein, 79.84% carbohydrates, 2.83% fat, 48.42% overall dietary fiber, 5.33% ash and 7.07% moisture. Little millet bran has a phenolic concentration of 465.67 µg, whole grain contains 148.53 µg, and pearled grain has 78.63 µg. Proso millet bran is composed of 9% fat, 26% carbohydrates, 36% dietary fiber and 14% protein, along with 3 mg gallic acid equivalent/g of phenolics. Bran is a promising ingredient for creating innovative functional and therapeutic foods since it contains good nutrients like protein, fat, dietary fiber, phenols, phytonutrients, flavonoids, and antioxidants. Hydrolytic rancidity is the primary challenge in using bran as food, and the only way to prevent it is through a process called stabilization, which inactivates the enzymes responsible for this issue. Stabilization of bran is crucial to preventing the formation of free fatty acids (FFA) due to the action of lipase. This review addresses stabilization techniques, health and therapeutic benefits and industrial applications for developing MB-based food products like bakery products, beverages and bran oil.

Keywords

bran; debranning; industrial applications; stabilisation; therapeutic property

Introduction

Millets provide a notable return on investment, even under the pressures of climate change. Their climate-resilient features contribute to sustainable agriculture without chemical fertilizers (1). Sorghum (*Sorghum bicolor*), Pearl millet (*Pennisetum glaucum* L.), Finger millet (*Eleusine coracana* L.) and small millets like Barnyard millet (*Echinochloa frumentacea* L.), Proso millet (*Panicum miliaceum* L.), Kodo millet (*Paspalum scrobiculata* L.), Little (Kutki) millet (*Panicum sumatrense*) and Foxtail millet (*Setaria italica* L.) are widely cultivated in India that are adaptable to different agroclimatic conditions. Historically, the nation produced and consumed

many millets, covering nearly the same region as wheat and rice (2). Millets are distinctive among cereals for their high calcium levels, dietary fiber, polyphenols, and protein. In recent years, millet-based dishes have increasingly been promoted for their role in a balanced diet and managing various health issues (3). However, the area designated for cultivating nutri cereals experienced a significant decline in the post-Green Revolution era. Between 1950-1951 and 2018-2019, this reduction amounted to 41.65%, indicating a substantial shift in agricultural priorities and practices that impacted the production of these important crops (2). The primary factors contributed to this decline include low yield compared to other food crops, absence of price incentives and input subsidies, subsidized supply of high-quality cereals via the Public Distribution System (PDS), shift in consumer preference since consequences of processing challenges, low longevity of flour and low social standing associated with millets (4). Additionally, the importance given to rice and wheat during the Green Revolution (5)

Bran is the tough outer layer of millet obtained during the initial processing stages, such as husking, polishing, dehulling, debranning, and milling (6,7). These processes make grains more palatable and easier to cook or further process (6,8).

The quantity of bran extracted varies from grain to grain according to the grain's structure and processing variables such as time, equipment and degree of dehulling or milling. For instance, bran constitutes 13-19% of the total grain weight in wheat (9) and 5-8% in rice (10). Millet bran, rich in fiber, can aid digestion and promote gut health. It can help control blood sugar levels, especially beneficial for diabetics or pre-diabetics. Millet bran's omega-3 to omega-6 ratio may reduce inflammation blood pressure and enhance blood vessel function, potentially reducing heart disease and stroke risk. Alpha-linolenic acid may aid cognitive function, memory, and learning, especially during childhood. Studies suggest omega-3 fatty acids may benefit individuals with neurological diseases. Further research is needed to understand their health benefits and impact on human health (11).

The husk and bran content of minor millets vary, with foxtail millet having 13.5% husk and 1-2% bran (12), while kodo millet and barnyard millet have up to 37% and 23% of each, respectively (13-15). Primary operations such as milling, dehulling, dehusking and debranning typically result in the removal of around 16-17% of bran in small millets. The production statistics pertaining to small millets demonstrate that roughly 0.062 to 0.066 metric tons of bran is yielded post-dehulling. This bran, which is acquired as a by-product, is either discarded or utilized as livestock feed (16)

During the refinement process, bran is removed, significantly reducing nutrients, dietary fiber, tannins, flavonoids, antioxidants, and polyphenols, suggesting that bran is an excellent source of these vital elements (17-21). The objective of this study on utilizing MB is to enhance nutrition through its fiber and vitamins, support health by reducing chronic disease risk, and promote sustainability by repurposing an agricultural byproduct. Additionally, it aims to assess the potential benefits for local farmers, provide gluten-free options and preserve culinary heritage, all while exploring market opportunities in the growing health food sector.

Production Scenario of Millet in India

Millets in India are broadly categorized into major and minor millets based on their size and scale of cultivation. Pearl millet and sorghum are major millets, while finger millet, foxtail millet, kodo millet, proso millet, small millet, barnyard millet and browntop millet are minor millets. India is the world's biggest millet producer. Most of India's states cultivate one or more types of millet crops (22). India produces 16.9 million tons of millets, or around 6% of the nation's total food grain supply, on approximately 12.7 million hectares. The production of 10.1 million tonnes of pearl millet is cultivated on around 7.4 million hectares. Finger millet (1.1 million ha) yields 1.58 million tonnes, sorghum (4.35 million ha) yields 4.63 million tonnes, and other millets (0.44 m ha, yielding 0.35 million tonnes) are the next most productive crops. Over 95% of the region planted under millet cultivation is composed of sorghum, pearl millet and finger millet and less than 5% is made up of small millets, such as barnyard millet, foxtail millet, little millet, kodo millet and proso millet (23).

Debranning Techniques Adopted for Millets and Cereals

Every minor millet grain has the same fundamental structure as other millet grains: an endosperm, a germ, a pericarp, a seed coat or testa and an aleurone layer (24). According to botanical characteristics, millet is divided into caryopses and utricles. In caryopses, the pericarp adheres to the seed firmly, but in utricles, the pericarp and seed coat are only connected at one point, making the pericarp easily removed. Foxtail, proso and barnyard are utricule types, whereas kodo millet is caryopsis (25,26). However, some studies have categorized proso under the utricule type and kodo, foxtail and barnyard under the caryopsis type (27). This disparity emphasizes the intricacy and diversity of botanical classifications, implying that additional study could be required to create a firmer categorization scheme for millets.

Traditional manual hand pounding is still used for basic processing steps, such as dehulling and decortication. Three distinct methods for dehulling small millets are abrasive decortication, rubbing technique and metal friction decortication. Small millets can be ground into fine powder using hammer mills, burr mills and plate or disk mills (28). Due to their small size, a significantly larger amount of the grain-specifically, the bran, bran-rich fraction and husk-is lost during processing, resulting in a higher loss of nutrients (29). Because of its nutritional value, bran (or hull) from millets, including proso, foxtail, pearl millet, jowar, etc., are processed as livestock feed (30).

A secondary product of millet processing produces MB (31,32). Various researchers have documented the utilization of rice dehullers, millers, barley pearlers, or abrasives because of smaller grain sizes and the absence of specifically designed dehullers for processing minor millet grains (28). This is the limiting factor for using bran in value-added products. Several scientists have reported the procedures involved in dehusking, dehulling, polishing, or milling grains to extract hull and bran from millets for research purposes, as depicted in Fig. 1.

Stabilisation of Bran

Despite the nutritional advantages, bran is primarily disposed of due to its shortened shelf life caused by the synthesis of FFA (38). Therefore, bran needs to be stabilized immediately after being obtained. Stabilization is the process of inhibiting hydrolytic degradation. A FFA level of more than 10 mEq/kg for human




Fig. 1. Different debranning machineries used for processing of millets.

consumption is unsuitable (39). Lipase activity and formation are the primary indicators of hydrolytic degradation in rice bran (RB). Lipoxygenase and peroxidase activities are less common indicators of RB deterioration. However, FFA is the most frequently used measure due to its straightforward determination (40). Many authors have documented the successful stabilization of cereal bran using various techniques, as shown in Table 1.

Millet bran was stabilized using a 900W microwave for 2.5 minutes (49). Chemicals like acetic acid, phosphoric acid and sodium metabisulphite failed to limit the rise in FFA. However, Hydrochloric acid at a concentration of about 30 ml/Kg helps substantially in managing the lipase activity and decreases the FFA (50). Rice bran was microwave-heated (2450MHz for 3min) to stabilize it. Stabilized RB was infected with lactic acid Bacillus culture to make probiotic-treated RB. The FFA percentage in RB that had been probiotic-treated ranged from 4.35 to 7.95 over the first four weeks of storage, while it varied from 4.10 to 7.50 in RB that has been stabilized from the first to the fourth week (51). Microwave heating at 100% power for 1-2 minutes and dry heating at 120°C for 10-15 minutes is used to stabilize the RB. Both heat stabilization and microwave heating were more effective than dry heating at keeping RB from rancid (52).

Free fatty acid, peroxide, anisidine and iodine levels were examined to track chemical changes during Proso millet bran (PMB) 180 days of storage at room temperature and in a refrigerator. During the duration of the storage, there was no discernible alteration in the iodine value of 137 ± 2 . There was a substantial interaction between the storage circumstances and time for the FFA levels, peroxide and anisidine. The initial thirty days of preservation saw a significant rise in FFA content until the ninetieth day. In both storage conditions, a similar trend was seen. Hydrolytic rancidity development results in an increase in the content of FFA (53).

Numerous attempts have been made to enhance bran stabilization by utilizing microwaves, ohmic heating, and infrared treatments. Stabilization and increased rice bran oil (RBO) extraction yield can be obtained by using ohmic heating with an

electrical field of 100 V/cm intensity and an alternating electric current of 60 Hz (54).

To stabilize RB and assess the stability efficiency by residual lipase activities, a comparison was made between 11 distinct treatments, comprising 6 heating treatments and 5 non-heating treatments. The findings suggested that RB may be stabilized by microwave, extrusion, steaming, -80°C, ultraviolet light and infrared heating (55). While oil peroxide values and color may increase, extrusion and microwave methods are appropriate for large-scale, fast industrial processing. The RB stabilized by steaming and oil extracted by immersion had the lowest α -tocopherol ($11.12 \pm 0.48 \mu\text{g/g}$) and γ -oryzanol ($957.18 \pm 53.7 \mu\text{g/g}$) compared to the greatest α -tocopherol ($42.38 \pm 0.53 \mu\text{g/g}$) and γ -oryzanol ($1190.1 \pm 89.3 \mu\text{g/g}$) in the ohmically stabilized RB. Ohmically stabilized RBO has a greater overall antioxidant capacity than steaming (56).

On the other hand, the UV irradiation process is an easy and energy-efficient stabilization technique that doesn't affect nutrients or oil quality. The RB stabilization through extrusion processing enhanced its functional attributes. It increased the quantity of dietary fiber and improved the capacity to hold water and foam but lowered the capacity to hold oil. Phytic acid concentration was also decreased by stabilization in contrast to unstabilized bran (57).

FSSAI Specifications for Millet bran /Cereal bran

The RBO must exhibit clarity and be devoid of rancidity, impurities, sediments, foreign substances, water and artificial colorings or flavorings. Clarity is determined by the absence of cloudiness after storing the filtered sample at 35°C for 24 hours. Refining is necessary to make RBO suitable for human consumption (58). Specification for RBO is given in Table 2.

Nutritional Profile of Different Millet Bran

The nutritional composition of MB differs across various millet types. Here's an overview of the nutritional profiles of some common MB (Table 3).

Table 1. Summary of different stabilization methods and their effect

Technique for stabilization	Criteria for stabilization	Assessment of stabilization	Result of stabilization	Reference
Extrusion	70°C, 90 °C, 110 °C and 130°C, with screw speed set at 300 rpm	Acid value Peroxide value Lipoxygenase activity Lipase activity	After 28 days of accelerated storage at 37°C, the acid value of untreated RB was approximately 6 mg KOH/100 g. In contrast, the acid values of stabilized samples were around 1.7 mg KOH/100 g, except for the atmospheric steam-treated RB.	(41)
Microwave heating	700 W, 2450 MHz for 1, 3 and 5 min (moisture content was adjusted to 21% before the treatment). MW power densities were 2, 4 and 6 W/g	FFA Peroxide value	The FFA content in RB treated with microwave (MW) at a power density of 6 W/g for 5 minutes was 1.12%, while the untreated control had an FFA content of 58.5% after 28 days of storage at room temperature.	(42)
Microwave heating	850, 925, and 1000 W for 3, 4.5 and 6 min	FFA Acid value Peroxide value	Stabilization was examined in three RB milling fractions and composite RB. The findings suggest stabilizing at 925 W for 3 minutes is optimal for the RB milling fractions.	(43)
Infrared heating	Medium wave IR radiation at 500, 600, and 700 W IR power for 3.0, 5.5 and 7.0 min	FFA	Stabilization was studied in three RB milling fractions. Stabilizing at 700 W using infrared power for 7 minutes resulted in a shelf life of 90 days, with no significant change in the FFA content of the RB fraction obtained from the first whitening step.	(44)
Infrared heating	2400 W, unspecified radiation intensity at 100, 120 and 140°C for 5, 10, 15 and 20 min.	Peroxide value FFA	The FFA content of raw RB rose from 4.4% to 62.8% after 6 months of storage at 25-30 °C. In contrast, RB was treated with infrared (IR) at 140 °C for 15 and 20 minutes and its FFA content was maintained at approximately 7%.	(45)
Hot air-assisted radio frequency (HA-RF)	A 6 kW, 27.12 MHz pilot-scale free-running oscillator RF system integrated with a hot air oven.	FFA Peroxide value Lipoxygenase activity Lipase activity	Rice bran treated with hot air-radio frequency (HA-RF) heating to 100 °C for 15 minutes can be stored at 35 °C for up to 60 days while remaining below acceptable quality thresholds without any negative effects on product quality.	(46)
Infrared heating	Infrared drying apparatus until the surface temperature of the RB reached 85 °C.	FFA Peroxide value Lipase activity Lipoxygenase activity Peroxidase activity FFA Peroxide value	The FFA content of untreated RB rose from 5.41% to 49.87%, while that of infrared (IR)-treated RB increased from 5.21% to 14.11% after 20 days of storage at 20 °C. Lipase, lipoxygenase, and peroxidase activities were significantly higher in the IR-treated samples by the end of the storage period.	(47)
Ohmic heating Electrical	The initial moisture contents were 20%, 30% and 40%. The applied voltages were 132 V, 150 V, 168 V, 189 V and 216 V, with a frequency of 50 Hz. The field strengths used were 44, 50, 56, 63 and 72 V/cm.	FFA Peroxide value	After 75 days of storage, the FFA content of ohmically heated bran was 4.77%, while raw RB had an FFA content of 41.84%.	(48)

Table 2. FSSAI Standards for rice bran oil

Parameter	Specification
Moisture and Volatile Matter	Not more than 0.1% by weight
Saponification value	180 - 195
Acid value	The accepted threshold is a maximum of 0.5
Iodine value (Wij's method)	90 - 105
Refractive Index at 40°C Or Butyro-refractometer reading at 40°C	1.4600 - 1.4700 51.0 - 66.4
Unsaponifiable matter,% by weight (a) for chemically refined (b) for physically refined Oryzanol Content	(a) Not more than 3.5% (b) Not more than 4.5% Not less than 1.0%
Flash Point (Pensky Marten Closed method)	Not less than 250°C

Table 3. Nutritional profile of various millet brans

Millet bran	Moisture (%)	Protein (%)	Fat (%)	Fiber (%)	Ash (%)	reference
Foxtail millet bran	6.10 - 9.07	6.78 - 12.93	5.65 - 9.63	42.56- 51.69	2.15 - 7.78	(59)
Proso millet bran	7.32	1.78-26.33	1.92- 8.96	31.63	5.77- 8.44	(53)
Kodo millet bran	5.53 - 7.07	4.92	2.83	48.42	5.33 - 7.74	(60)
Sorghum bran	6.13	6.7	4.6-4.9	18.04	2.15 - 7.78	(61)
Barnyard bran	6.79	5.15 - 5.19	4.04	38.4%	9.11 - 23.10	(62)

Proso millet bran

The quantity of fat in various types of proso millet and bran varied. White-coloured proso MB types had 3.95 - 6.84% (free) and 0.33 - 0.49% (bound) lipids, while red variants had 3.45 - 4.29% and 0.30 - 0.70%, respectively. Bran of the red and white kinds had a total lipid content ranging from 4.14 to 7.27 %. Ninety percent of the fatty acids in proso millet flour and bran are composed of oleic linoleic and palmitic acid (33).

Proximate analysis of four different types of proso MB was studied (Table 4). Crude fiber, crude protein and crude fat content were highest in yellow proso MB, starch content was highest in white proso MB and ash content was highest in black proso MB (63).

Functional properties like oil holding capacity (OHC) and water holding capacity (WHC) were studied. WHC and OHC were higher in the husks than in the bran. The white proso millet husks had the highest levels of WHC (2.53 g/g) and OHC (1.61 g/g). The OHC of the yellow and black proso millet brans was the greatest, while the yellow proso millet brans' WHC was maximum (63).

Foxtail millet bran

The phenol content of milled FMB was higher than that of whole or pearled grain; the highest phenol content (mg/100 g) was found in foxtail bran (510.53 ± 50.34), followed by whole grain (132.76 ± 8.66) and the lowest phenol content (104.00 ± 0.00) in pearled grain (64).

It was noted that 9.39 ± 0.17% crude oil, 12.48 ± 0.41% crude protein, 51.69 ± 2.14% crude fiber, 7.50 ± 0.18% ash and 8.29 ± 0.16% moisture were found in FMB. Tocopherol in the oil extracted from FMB was 64.83 ± 0.83 mg/100 g. It was mostly made up of 15.53 ± 0.31 mg/100 g oil of α -tocopherol and 48.79 ± 0.46 mg/100 g of oil of γ -tocopherol. Additionally, linolenic (66.5%), oleic (13.1%) and saturated fatty acids, including stearic and palmitic acid, were present in foxtail millet bran oil (FMBO) (16).

Foxtail millet bran can serve as a natural source of antioxidants, both as a nutraceutical and for the food industry. The predominant unsaturated fatty acids in FMBO are linoleic and oleic acid, making up more than 80% of the oil. Saturated fatty acids comprise a small part of the fatty acid, accounting for about 14% of the total. Furthermore, it was discovered that FMO possesses comparatively elevated levels of tocopherols, sterols, γ -oryzanol and squalene-all of which have the potential to be advantageous (65).

The functional and physicochemical characteristics of FMB dietary fiber (FMBDF) were examined, and the possibility of extracting dietary fiber from the bran using enzyme techniques was examined. The findings showed that FMB was an excellent way to produce dietary fiber. The results of the functional and physicochemical property experiments indicated that the FMBDF had a swelling power of 2.06 ml/g and a water-holding capacity of 3.24 g/g. Its capacity to adsorb bile salts (76.65 mmol/g for sodium taurocholate and 143.03 mmol/g for sodium cholate) explains its capacity to decrease cholesterol (59)

Table 4. Proximate analysis of brans of different color proso millet bran

Parameter	Red	White	Yellow	Black
Crude protein (%)	14.75±0.08	14.56±0.04	16.23±0.03	13.13±0.01
Crude fat (%)	20.43±0.01	20.29±0.07	24.27±0.36	22.24±0.18
Crude fiber (%)	35.80±0.37	36.23±2.31	39.92±0.91	39.19±0.43
Starch (%)	6.36±0.27	7.11±0.34	6.62±0.38	4.63±0.10
Ash (%)	8.54±0.11	8.51±0.09	6.17±0.03	8.69±0.13
Moisture (%)	7.62±0.06	6.74±0.23	6.41±0.15	6.82±0.17

According to fatty acid composition analysis, defatted foxtail millet bran (DFMB) oil has a higher mono-unsaturated fatty acid content than RB, peanut and olive oils. Large amounts of mono-unsaturated fatty acid, which can lower cholesterol, were present in the FMBO. Still, there was a lack of poly-unsaturated fatty acids necessary for human health (16). Fig 2 illustrates the high concentration of mono-unsaturated fatty acid (87% mono-unsaturated, 9% saturated fatty acid 4% and poly-unsaturated) in FMBO (66). Therefore, FMBO, which is also high in mono-unsaturated fatty acids, may be a more nutrient-dense dietary source than known olive oil on the market.

Kodo millet bran

A study revealed the quantity of protein (4.92%), carbohydrates (79.84%), fat (2.83%), overall dietary fiber (48.42%), ash (5.33%) and moisture (7.07%) in 100 g of KMB (60). Kodo millet bran exhibited the maximum antioxidant potential due to significantly greater levels of dietary fiber (61.52%), phytic acid (630 mg/100 g), flavonoids (22.37 μ g retinol equivalent/g) and total phenols (449.27 mg gallic acid equivalent/100 g) (67).

Little millet

Upon milling little millet, 27% of the hull was produced. Compared to whole grain (148.53 ± 6.31) and pearled grain (78.63 ± 1.79), bran had a phenol concentration (465.67 ± 20.1). For both millets, there was no noticeable difference in the phytic content, zinc, or copper in the whole grain or its milled fractions (64)

Sorghum bran

Sorghum bran can function as a nutraceutical because it is an excellent source of fiber, antioxidants and phytonutrients. The percentage of dietary fiber in sorghum bran varied from 40% to 45%. When sorghum bran is employed in product development, its high fiber and phytate content lowers the product's calorific value (68). 3-Deoxyanthocyanidins were found in sorghum and sorghum bran, which have antioxidant qualities equivalent to anthocyanins and were resilient to pH changes. Black sorghum bran had twice as much anthocyanins (10.1 mg/g) as red (3.6 mg/g) and brown (3.6 mg/g) sorghum bran. Bran contained three to four times more anthocyanins than whole grains reported. Thus, it was discovered that black sorghum was the most promising crop to use as a natural food coloring source (69).




Fig. 2. Comparison of different oil composition(66).

Application of Millet Bran in Food Processing

Breads and buns

Sorghum bran supplement (15%) gave bread outstanding phenolic and antioxidant content without compromising the bread's physical characteristics, such as specific volume or appearance. The bran gave the loaves a dark brownish color (dahlberg 55). Similarly, bread with 12% brown sorghum bran, 5% flaxseed and 2% soy flour added dark brown color to the loaves and improved dietary fiber, antioxidants and palatability (70).

Breads supplemented with 0, 5, 10, 15 and 20% RB showed that up to 10% incorporation was acceptable. Sensory acceptance fell as the percentage of bran increased. Based on gravimetric tests, each unit of bran showed an increase in water absorption, a decrease in volume and a rise in weight. Farinograph revealed that when bran addition increased, dough mixing stability decreased (71).

Muffins

Rusks and muffins were made by incorporating barnyard millet bran (BMB) with wheat flour at varying percentages (0, 5, 10, 15, 20, 25 and 30%). When BMB was added, the hardness of the rusk and muffin increased. Muffins at 15% and rusks at 30% were suitable (49).

Foxtail millet bran (0, 10, 15, 20, 25 and 30%) made muffins instead of refined wheat flour. Physical properties like height, weight and baking loss were measured over five replications. The muffin height increased as bran content increased and the color of the muffin became darker with bran incorporation. The panel members accepted muffins enriched with 30% MB (72).

Beverages

To create an antioxidant-rich wine from FMB, research found that foxtail millet bran wine (MBW) had six times more polyphenols than millet wine (MW) and demonstrated stronger antioxidant activity in the 2, 2-Diphenyl-1-picrylhydrazyl (DPPH), Trolox Equivalent Antioxidant Capacity (TEAC) and Ferric Reducing Antioxidant Power (FRAP) tests. It was determined that vanillic acid, p-coumaric acid, syringic acid, and ferulic acid were the four phenolics in it. Additionally, MBW has more amino acids than commercially produced millet wine (CAMW) and MW. Results showed that it is possible to use MB wine as a nutritional supplement, nutraceutical, or functional beverage (73).

Production of Millet Bran Oil and Extraction Methods

Millet bran oil

Originally from China, foxtail millet is one of the earliest crops to be cultivated and is currently grown worldwide (66). The nutritional and medicinal benefits of millet have been well documented (74). The oil extracted from FMB is high in linoleic acid (66.5%) and oleic acid (13.1%). Stearic acid (6.3%) and palmitic acid (6.4%) are two of the saturated fatty acids. Linoleic acid (71.2%) was the predominant fatty acid in the millet oil. Dilinoleoyl-monoolein (LLO, 17.2%) and trilinoleate (LLL, 29.3%) were identified as the two predominant triacylglycerols, as determined by the 1,3-random-2-random hypothesis (16).

Supercritical solvent extraction techniques

In recent years, there has been a growing interest in supercritical carbon dioxide extraction (SCE). This technology employs

carbon dioxide in a supercritical state, above its critical temperature and pressure, making it an attractive solvent for temperature-sensitive materials (75). The subcritical carbon dioxide (CO₂) Soxhlet method is a novel approach that combines the advantages of Soxhlet extraction and CO₂ extraction to yield RBO effectively. Using the subcritical CO₂ process, the oil contains about ten times more oryzanol and tocol compounds and lower FFA levels and peroxide values than oil extracted with hexane (76). Traditional methods mainly use solvents and cold-pressing techniques to extract bran oil. Using commercial-grade hexane in solvent extraction (SE) presents several disadvantages, including poor color quality in RBO, substantial air pollution, and health risks due to its toxicity (77). Subcritical propane extraction (SPE), SCE and traditional SE were used to extract the FMBO. Examinations were conducted on the yield, physicochemical characteristics, fatty acid profile, tocopherol content, and oil oxidative stability. Foxtail millet bran oil yields by SPE, SCE, and SE were 21.79%, 19.65% and 17.14% of the raw material's weight, respectively. These values corresponded to 75.54%, 86.60% and 96.03% of the total quantity of oil. Based on Gas chromatography (GC) analysis, the variation in the fatty acid profile was insignificant. Still, the methods of extraction had a substantial impact on the physicochemical attributes (peroxide value, color, and saponification value) (78). Through HPLC analysis, the primary vitamin E components found in the produced oils were determined to be α - and β -tocopherols. SPE proved to be more effective than SE and SCE in extracting tocopherols. The oil recovered by SPE demonstrated the highest oil oxidative stability in the Rancimat test, which may have been caused by its low peroxide value and high tocopherol content. Compared to SCE and SE, SPE used smaller periods and lower pressures in consideration of the oil quality. SPE proved an effective and focused technique for obtaining oil from FMB (79).

Solvent extraction techniques

Solvents are used to extract oil from seeds that have relatively little oil or pre-pressed oil cakes to increase oil yields. The economic benefits of commercial-grade hexane have made it a popular choice worldwide. It is a powerful solvent that is easy to recover and has good solubility. Hexane, however, is linked to several issues. It causes serious environmental problems such as air pollution and ends up producing bran oil with low color quality. Its toxicity also presents health risks. Numerous efforts have explored alternative solvents, including halogenated hydrocarbons, alcohols, carbon dioxide, hydrocarbons (utilized in supercritical fluid extraction), and water (80). Because of their higher safety, other short-chain alcohols, such as isopropanol, and ethanol, have also been suggested as an alternate extraction solvent. (81,82). For high yields of high-quality crude edible oil, d-limonene, an agricultural by-product of the citrus sector, is a good substitute for hexane-based solvents (83). However, due to its high boiling point and elevated latent heat, limonene is not ideal for edible oil extraction because of its high energy cost. Energy-efficient solvent recovery techniques, like membrane separations, may help overcome this issue, making limonene-based oil extraction more cost-effective.

Mechanical extraction techniques

The mechanical or cold pressing method doesn't utilize heat treatment or organic solvents. It is an alternative to traditional practices and shows promise in oil extraction due to its lower labor

and cost requirements than SE. Cold pressing offers advantages in terms of safety, efficiency and simplicity. Furthermore, oils extracted through this method boast superior nutritional properties and chemical-free end products, increasing interest in cold-pressed plant oils. (84). Given the drawbacks of SE-such as the toxicity and cost of solvents and significant environmental impacts -alternative extraction methods are gaining traction. The desire for high-quality crude oil, a focus on environmental sustainability and the ability of mechanical systems to support continuous processes are essential factors driving this shift (85).

Therapeutic Properties of Millet Bran

Consuming whole grains is linked to a lower chance of developing several diseases, including constipation, type II diabetes, hypercholesterolemia, cardiovascular disease and several gastrointestinal malignancies (86). Whole grains are nutrient-rich, including dietary fiber, vitamins, minerals, phytochemicals and phenolic compounds. The grain's bran contains most of these nutrients (87,88). Therefore, a considerable amount of fiber, minerals and phenolic chemicals are present in millet and wheat brans, which contribute to their natural antioxidant potential and may be used to prevent or treat certain illnesses. The interaction between fiber and antioxidants may contribute to fiber's health benefits.

In addition, FMB's bound polyphenols of the inner shell (BPIS) showed anti-inflammatory properties in HT-29 cells and nude mice produced by lipopolysaccharide (LPS). When BPIS was applied to LPS-induced HT-29 cells, pro-inflammatory cytokines like IL-1 β (interleukin-1 β) and IL-6 (interleukin-6) were downregulated, and anti-inflammatory cytokines like IL-10 (interleukin-10) were increased. By using BPIS, tumor tissue growth was also inhibited. In light of this, BPIS may be employed as a pro-oxidant agent to reduce inflammation (79).

A new 35 kDa protein was isolated and purified from FMB (89). The findings demonstrated that by triggering cell cycle G1-phase arrest, the crude protein dramatically inhibited colon cancer cell growth *in vitro* and in xenografted tumors in nude mice. Foxtail millet bran-derived peroxidase (FMBP) inhibited colon cancer cells' migration ability, making it a potential treatment for colon cancer patients (90). The FRAP (ferric reducing antioxidant power) and DPPH (2, 2-Diphenyl-1-picrylhydrazyl) assays demonstrated significant antioxidant activity in FMBO. It was also beneficial in reducing ethanol-induced liver damage in mice (91).

Challenges Of Utilizing Millet Bran

Utilizing MB presents several challenges. Firstly, there is a lack of efficient debranning machinery, which hampers processing effectiveness. Additionally, the high costs of extracting MB oil make it less competitive than other oil sources. Compounding these issues is the insufficient research on the toxicity and therapeutic benefits of MB, which limits its acceptance in the market. Furthermore, stakeholders often have low awareness of the nutritional advantages of MB, affecting its potential for wider adoption. Lastly, in some social circles, using MB for human consumption is viewed unfavorably, hindering its utilization (11).

Conclusion

Emerging trends in health-conscious diets and sustainable food sources align well with the benefits of MB. As consumers seek nutritious options, MB's high fiber, vitamins and minerals support digestive health and overall well-being. Its resilience and low water requirements enhance its sustainability, attracting environmentally conscious consumers. Despite being underutilized compared to other cereal brans like RB, MB has significant potential for developing functional and nutraceutical foods due to its nutritional properties and high monounsaturated fatty acid content. Rich in fiber, minerals and phenolic compounds, it offers natural antioxidant benefits and can help prevent various health issues, including cardiovascular disease and type II diabetes. However, research into food products such as beverages, bran oil, and baked goods remains limited. Advances in debranning and stabilizing technologies could accelerate MB's industrial applications, making it a compelling ingredient for modern diets.

Acknowledgements

We sincerely thank Tamil Nadu Agricultural University for the technical support, and we would like to thank the Social Education Economical Development Society (SEEDS) Farmer Producer Organization (FPO) Virudhunagar, Tamil Nadu, for providing financial support.

Authors' contributions

PK carried out writing and proofreading of the article. VM carried out conceptualization, literature review, and correction of the article. RV designed the article. ML helped collect literature and gave technical help. AK contributed to literature collection and MI contributed in prefinal correction.

Compliance with ethical standards

Conflict of interest: All authors do not have any conflict of interest to declare.

Ethical issues: None

Declaration of generative AI and AI-assisted technologies in the writing process

While preparing this work, we used the Grammarly AI tool to improve grammar and increase the article's readability.

References

1. Thilakarathna MS, Raizada MN. A review of nutrient management studies involving finger millet in the semi-arid tropics of Asia and Africa. *Agronomy*. 2015;5(3):262-90. <https://doi.org/10.3390/agronomy5030262>.
2. Sukumaran Sreekala AD, Anbukkani P, Singh A, Dayakar Rao B, Jha GK. Millet production and consumption in India: Where do we stand and where do we go?. *Natl Acad Sci Lett*. 2023;46(1):65-70. <https://doi.org/10.1007/s40009-022-01164-0>
3. Anitha S, Givens DI, Subramaniam K, Upadhyay S, Kane-Potaka J, Vogtschmidt YD, et al. Can feeding a millet-based diet improve the growth of children?-A systematic review and meta-analysis. *Nutrients*. 2022;14(1):225. <https://doi.org/10.3390/nu14010225>

4. Indian Institute of Millets Research (ICAR-Millets). Vision 2050 [Internet]. Indian Council of Agricultural Research; 2015 [Cited 2024 Dec 24]. Available from: <https://krishi.icar.gov.in/jspui/handle/123456789/2675>
5. Directorate of Millets Development. Status paper on coarse cereals [Internet]. Department of Agriculture and Cooperation, Ministry of Agriculture, Government of India; March 2014 [Cited 2024 Dec 24]. Available from: <https://www.nfsm.gov.in/StatusPaper/StatusMillets2016.pdf>
6. Saleh ASM, Zhang Q, Chen J, Shen Q. Millet grains: nutritional quality, processing and potential health benefits. *Compr Rev Food Sci Food Saf.* 2013;12(3):281-95. <https://doi.org/10.1111/1541-4337.12012>
7. Kalpanadevi C, Singh V, Subramanian R. Influence of milling on the nutritional composition of bran from different rice varieties. *J Food Sci Technol.* 2018;55:2259-69. <https://doi.org/10.1007/s13197-018-3143-9>
8. Patel S. Cereal bran fortified-functional foods for obesity and diabetes management: Triumphs, hurdles and possibilities. *J Funct Foods.* 2015;14:255-69. <https://doi.org/10.1016/j.jff.2015.02.010>
9. Onipe OO, Jideani AI, Beswa D. Composition and functionality of wheat bran and its application in some cereal food products. *Int J Food Sci Technol.* 2015;50(12):2509-18. <https://doi.org/10.1111/ijfs.12935>
10. Arendt EK, Zannini E. Cereal grains for the food and beverage industries. Woodhead Publishing Series in Food Science, Technology and Nutrition; 2013. <https://doi.org/10.1533/9780857098924>
11. Vashishth N, Maurya NK. Millet bran: Unveiling its potential health benefits. *International Journal of Nutritions.* 2024;1(2):11-19.
12. Pawar VD, Machewad GM. Processing of foxtail millet for improved nutrient availability. *J Food Process Preserv.* 2006;30(3):269-79. <https://doi.org/10.1111/j.1745-4549.2006.00064.x>
13. Taylor JR, Emmambux MN. Gluten-free foods and beverages from millets. In: Arendt EK, Dal Bello F, editors. *Gluten-free cereal products and beverages.* Academic Press; 2008. p. 119-48. Available from: <https://doi.org/10.1016/b978-012373739-7.50008-3>
14. Dharmaraj U, Rao BS, Sakhare SD, Inamdar AA. Preparation of semolina from foxtail millet (*Setaria italica*) and evaluation of its quality characteristics. *J Cereal Sci.* 2016;68:1-7. <https://doi.org/10.1016/j.jcs.2015.11.003>
15. Sharma N, Niranjana K. Foxtail millet: Properties, processing, health benefits and uses. *Food Rev Int.* 2018;34(4):329-63. <https://doi.org/10.1080/87559129.2017.1290103>
16. Liang S, Yang G, Ma Y. Chemical characteristics and fatty acid profile of foxtail millet bran oil. *J Am Oil Chem Soc.* 2010;87(1):63-67. <https://doi.org/10.1007/s11746-009-1475-3>
17. Yu L, Nanguet AL, Beta T. Comparison of antioxidant properties of refined and whole wheat flour and bread. *Antioxidants.* 2013;2(4):370-83. <https://doi.org/10.3390/antiox2040370>
18. Devi PB, Vijayabharathi R, Sathyabama S, Malleshi NG, Priyadarisini VB. Health benefits of finger millet (*Eleusine coracana* L.) polyphenols and dietary fiber: a review. *J Food Sci Technol.* 2014;51:1021-40. <https://doi.org/10.1007/s13197-011-0584-9>
19. Goudar G, Sathisha GJ. Effect of processing on ferulic acid content in foxtail millet (*Setaria italica*) grain cultivars evaluated by HPTLC. *Orient J Chem.* 2016;32(4):2251-58. <https://doi.org/10.13005/ojc/320458>
20. Ahmad F, Pasha I, Saeed M, Asgher M. Biochemical profiling of Pakistani sorghum and millet varieties with special reference to anthocyanins and condensed tannins. *Int J Food Prop.* 2018;21(1):1586-97. <https://doi.org/10.1080/10942912.2018.1502198>
21. Bora P, Das P, Bhattacharyya R, Saikia A. Effect of processing on the phytochemical content and antioxidant capacity of proso millet (*Panicum miliaceum* L.) milled fractions. *Int J Chem Stud.* 2018;6(4):18-22.
22. McDonough CM, Rooney LW, Serna-Saldivar SO. The millets. In: Kulp K, editor. *Handbook of cereal science and technology.* Revised and expanded. 2nd ed. Boca Raton: CRC Press; 2000. p. 177-201.
23. Bhat BV, Hariprasanna K, Ratnavathi CV. Global and Indian scenario of millets. *Indian Farming.* 2023;73(1):16-18.
24. Akanbi TO, Timilsena Y, Dhital S. Bioactives from millet: properties and effects of processing on bioavailability. In: Wang J, Sun B, Tsao R, editors. *Bioactive factors and processing technology for cereal foods.* Singapore: Springer; 2019. p. 171-83. https://doi.org/10.1007/978-981-13-6167-8_10
25. Taylor JR, Duodu KG. Effects of processing sorghum and millets on their phenolic phytochemicals and the implications of this to the health-enhancing properties of sorghum and millet food and beverage products. *J Sci Food Agric.* 2014;95(2):225-37. <https://doi.org/10.1002/jsfa.6713>
26. Dayakar Rao B, Ananthan R, Hariprasanna K, Bhatt KV, Rajeswari SS, et al. Nutritional and health benefits of nutri cereals [Internet]. Hyderabad: Nutri Hub TBI, ICAR Indian Institute of Millets Research (IIMR); 2018 [Cited 2024 Dec 24]. Available from: https://nutricereals.dac.gov.in/Publication/Pub_HealthyBenefits_Sep2018.pdf
27. Serna-Saldivar SO, Espinosa-Ramírez J. Grain structure and grain chemical composition. In: Taylor JR, Duodu KG, editors. *Sorghum and millets: chemistry, technology and nutritional attributes.* 2nd ed. Cambridge: Woodhead Publishing, Elsevier; 2019. p.85-129. <https://doi.org/10.1016/b978-0-12-811527-5.00005-8>
28. Chandhi GK, Annor GA. Millet minor: overview. In: Wrigley C, Corke H, Seetharaman K, Faubion J, editors. *Encyclopedia of food grains.* 2nd ed. San Diego: Academic Press; 2016. p. 199-208. <https://doi.org/10.1016/b978-0-12-394437-5.00012-7>
29. Gulia SK, Wilson JP, Carter J, Singh BP. Progress in grain pearl millet research and market development. In: Janick J, Whipkey A, editors. *Issues in new crops and new uses.* Alexandria, VA: ASHS Press; 2007. p. 196-203.
30. Tran G. Millet hulls [Internet]. Feedipedia, a programme by INRAE, CIRAD, AFZ and FAO; 2015 [Cited 2024 Dec 24]. Available from: <https://feedipedia.org/node/15695>
31. Devittori C, Gummy D, Kusy A, Colarow L, Bertoli C, Lambelet P. Supercritical fluid extraction of oil from millet bran. *J Am Oil Chem Soc.* 2000;77(6):573-79. <https://doi.org/10.1007/s11746-000-0092-7>
32. Taylor JRN, Kruger J. Millets. In: Caballero B, Finglas PM, Toldrá F, editors. *Encyclopedia of food and health.* Oxford: Academic Press; 2016. p. 748-57. <https://doi.org/10.1016/B978-0-12-384947-2.00466-9>
33. Lorenz K, Hwang YS. Lipids in proso millet (*Panicum miliaceum*) flours and brans. *Cereal Chem.* 1986;63(5):387-90.
34. Rhodes D. Diversity, genetics and health benefits of sorghum grain [dissertation on the Internet]. Columbia: University of South Carolina; 2014 [cited 2024 Dec 24]. Available from: <https://scholarcommons.sc.edu/etd/3005>
35. Devisetti R, Yadahally SN, Bhattacharya S. Nutrients and antinutrients in foxtail and proso millet milled fractions: Evaluation of their flour functionality. *LWT.* 2014 ;59(2):889-95. <https://doi.org/10.1016/j.lwt.2014.07.003>
36. Kumari D, Madhujith T, Chandrasekara A. Comparison of phenolic content and antioxidant activities of millet varieties grown in different locations in Sri Lanka. *Food Sci Nutr.* 2017;5(3):474-85. <https://doi.org/10.1002/fsn.3.415>
37. Palaniappan A, Yuvaraj SS, Sonaimuthu S, Antony U. Characterization of xylan from rice bran and finger millet seed coat for functional food applications. *J Cereal Sci.* 2017;75:296-305. <https://doi.org/10.1016/j.jcs.2017.03.032>
38. Sharma S, Kaur S, Dar BN, Singh B. Storage stability and quality assessment of processed cereal brans. *J Food Sci Technol.* 2014;51:583-88. <https://doi.org/10.1007/s13197-011-0537-3>

39. Orthoefer FT, Eastman J. Rice bran oil. In: Shahidi F, editor. *Bailey's industrial oil and fat products*. 6th ed. New York: John Wiley and Sons, Ltd; 2005.p.465-89. <https://doi.org/10.1002/047167849X.bio015>
40. Tao J, Rao R, Liuzzo J. Microwave heating for rice bran stabilization. *J Microw Power Electromagn Energy*. 1993;28(3):156-64. <https://doi.org/10.1080/08327823.1993.11688217>
41. Li Y, Gao C, Wang Y, Fan M, Wang L, Qian H. Analysis of the aroma volatile compounds in different stabilized rice bran during storage. *Food Chem*. 2023;405:134753. <https://doi.org/10.1016/j.foodchem.2022.134753>
42. Patil SS, Kar A, Mohapatra D. Stabilization of rice bran using microwave: Process optimization and storage studies. *Food Bioprod Process*. 2016;99:204-11. <https://doi.org/10.1016/j.fbp.2016.05.002>
43. Lavanya MN, Saikiran KC, Venkatachalapathy N. Stabilization of rice bran milling fractions using microwave heating and its effect on storage. *J Food Sci Technol*. 2019;56(2):889-95. <https://doi.org/10.1007/s13197-018-3550-y>
44. Yilmaz N. Middle infrared stabilization of individual rice bran milling fractions. *Food Chem*. 2016;190:179-85. <https://doi.org/10.1016/j.foodchem.2015.05.094>
45. Irakli M, Kleisariis F, Mygdalia A, Katsantonis D. Stabilization of rice bran and its effect on bioactive compounds content, antioxidant activity and storage stability during infrared radiation heating. *J Cereal Sci*. 2018;80:135-42. <https://doi.org/10.1016/j.jcs.2018.02.005>
46. Ling B, Lyng JG, Wang S. Effects of hot air-assisted radio frequency heating on enzyme inactivation, lipid stability and product quality of rice bran. *LWT*. 2018;91:453-59. <https://doi.org/10.1016/j.lwt.2018.01.084>
47. Yan W, Liu Q, Wang Y, Tao T, Liu B, Liu J, Ding C. Inhibition of lipid and aroma deterioration in rice bran by infrared heating. *Food Bioproc Tech*. 2020;13:1677-87. <https://doi.org/10.1007/s11947-020-02503-z>
48. Dhingra D, Chopra S, Rai DR. Stabilization of raw rice bran using ohmic heating. *Agric Res*. 2012;1:392-98. <https://doi.org/10.1007/s40003-012-0037-3>
49. Nazni P, Karuna TD. Development and quality evaluation of barnyard millet bran incorporated rusk and muffin. *J Food Ind Microbiol*. 2016;2(116):2. <https://doi.org/10.4172/2572-4134.1000116>
50. Akhter M, Afzal N, Haider Z, Raza MA. Inactivation of lipase enzyme by using chemicals to maximize rice bran shelf life and its edible oil recovery. *J Nutr Food Sci*. 2015 Sep 16;12:S12002. <https://doi.org/10.15761/ifnm.1000123>
51. Bhosale S, Vijayalakshmi D. Processing and nutritional composition of rice bran. *Curr Res Nutr Food Sci*. 2015;3(1):74-80. <https://doi.org/10.12944/crnfsj.3.1.08>
52. Lv SW, He LY, Sun LH. Effect of different stabilisation treatments on preparation and functional properties of rice bran proteins. *Czech J Food Sci*. 2018;36(1). <https://doi.org/10.17221/100/2017-cjfs>
53. Mustač NČ, Novotni D, Habuš M, Drakula S, Nanjara L, Voučko B, et al. Storage stability, micronisation and application of nutrient-dense fraction of proso millet bran in gluten-free bread. *J Cereal Sci*. 2020;91:102864. <https://doi.org/10.1016/j.jcs.2019.102864>
54. Lakkakula NR, Lima M, Walker T. Rice bran stabilization and rice bran oil extraction using ohmic heating. *Bioresour Technol*. 2004;92(2):157-61. <https://doi.org/10.1016/j.biortech.2003.08.010>
55. Yu C, Hu Q, Wang H, Deng Z. Comparison of 11 rice bran stabilization methods by analyzing lipase activities. *J Food Process Preserv*. 2020;44(4):e14370. <https://doi.org/10.1111/jfpp.14370>
56. Loypimai P, Moongngarm A, Chottanom P. Impact of stabilization and extraction methods on chemical quality and bioactive compounds of rice bran oil. *Emir J Food Agric*. 2015;27(11):849-56. <https://doi.org/10.9755/ejfa.2015.09.738>
57. Rafe A, Sadeghian A, Hoseini-Yazdi SZ. Physicochemical, functional and nutritional characteristics of stabilized rice bran from tarom cultivar. *Food Sci Nutr*. 2017;5(3):407-14. <https://doi.org/10.1002/fsn3.407>
58. https://www.fssai.gov.in/upload/uploadfiles/files/Compendium_Food_Additives_Regulations_08_09_2020-compressed.pdf
59. Zhu Y, Chu J, Lu Z, Lv F, Bie X, Zhang C, Zhao H. Physicochemical and functional properties of dietary fiber from foxtail millet (*Setaria italica*) bran. *J Cereal Sci*. 2018;79:456-61. <https://doi.org/10.1016/j.jcs.2017.12.011>
60. Sarma SM, Khare P, Jagtap S, Singh DP, Baboota RK, Podili K, et al. Kodo millet whole grain and bran supplementation prevents high-fat diet induced derangements in a lipid profile, inflammatory status and gut bacteria in mice. *Food Funct*. 2017;8(3):1174-83. <https://doi.org/10.1039/c6fo01467d>
61. Nirmala BY, Hanchinal RR, Basarkar PW. Antioxidant contents of whole grain cereals, millets and their milled fractions. *Journal of Dairying Foods and Home Sciences*. 2011;30(3):191-96.
62. Bisoi PC, Sahoo G, Mishra SK, Das C, Das KL. Hypoglycemic effects of insoluble fiber rich fraction of different cereals and millets. *J Food Process Technol*. 2012;3(11). <https://doi.org/10.4172/2157-7110.1000191>
63. Li M, Chang L, Ren J, Jiang F, Zhao N, Liu Y, et al. Nutritional, physical, functional properties and antioxidant potential of different colors proso millet husks and brans. *LWT*. 2022;171:114092. <https://doi.org/10.1016/j.lwt.2022.114092>
64. Sridevi. Evaluation of antioxidant properties of whole grains and processed regional cereals of Karnataka [master's dissertation on the Internet]. Dharwad: Department of Food Science and Nutrition, College of Rural Home Science, University of Agricultural Sciences; 2007 [cited 2024 Dec 24]. Available from: <https://krishikosh.egranth.ac.in/items/6fc85766-39c3-4615-8490-cb76269c8f2a>
65. Ji J, Liu Y, Ge Z, Zhang Y, Wang X. Oleochemical properties for different fractions of foxtail millet bran. *J Oleo Sci*. 2019;68(8):709-18. <https://doi.org/10.5650/jos.ess19063>
66. Le G, Amadou I, Amza T, Shi YH, Le GW. Chemical analysis and antioxidant properties of foxtail millet bran extracts. *Songklanakar J Sci Technol*. 2011;33(5):509-15.
67. Barbhai MD, Hymavathi TV. Nutrient, phytonutrient and antioxidant potential of selected underutilized nutri-cereal brans. *J Food Meas Charact*. 2022;16(3):1952-66. <https://doi.org/10.1007/s11694-022-01301-9>
68. Dahlberg JA, Wilson JP, Snyder T. Sorghum and pearl millet: health foods and industrial products in developed countries. In: *Alternative uses of sorghum and pearl millet in Asia: proceedings of an expert meeting; ICRISAT; Patancheru, Andhra Pradesh, India; 1-4 July 2003*. Patancheru: International Crops Research Institute for the Semi-Arid Tropics (ICRISAT); 2004.p. 42-59.
69. Awika JM, Rooney LW, Waniska RD. Properties of 3-deoxyanthocyanins from sorghum. *J Agric Food Chem*. 2004;52(14):4388-94. <https://doi.org/10.1021/jf049653>
70. Hines LR. Development of specialty breads as nutraceutical products [master's thesis]. College Station: Texas A and M University; 2007. 111 p.
71. Espinales C, Cuesta A, Tapia J, Palacios-Ponce S, Peñas E, Martínez-Villaluenga C, et al. The effect of stabilized rice bran addition on physicochemical, sensory and techno-functional properties of bread. *Foods*. 2022;11(21):3328. <https://doi.org/10.3390/foods11213328>
72. Mrunal B. Physico- sensory quality of foxtail millet (*Setaria italica*) bran enriched muffins. *Indian J Pure Appl Biosci*. 2020;8(6):28-33. <https://doi.org/10.18782/2582-2845.8438>
73. Guo X, Sha X, Rahman E, Wang Y, Ji B, Wu W, Zhou F. Antioxidant capacity and amino acid profile of millet bran wine and the synergistic interaction between major polyphenols. *J Food Sci Technol*. 2018;55:1010-20. <https://doi.org/10.1007/s13197-017-3014-9>

74. Hegde PS, Rajasekaran NS, Chandra TS. Effects of the antioxidant properties of millet species on oxidative stress and glycemic status in alloxan-induced rats. *J Oleo Sci.* 2005;25(12):1109-20. <https://doi.org/10.1016/j.nutres.2005.09.020>
75. Sookwong P, Mahatheeranont S. Supercritical CO₂ extraction of rice bran oil—the technology, manufacture and applications. *J Oleo Sci.* 2017;66(6):557-64. <https://doi.org/10.5650/jos.ess17019>
76. Chia SL, Boo HC, Muhamad K, Sulaiman R, Umanan F, Chong GH. Effect of subcritical carbon dioxide extraction and bran stabilization methods on rice bran oil. *J Am Oil Chem Soc.* 2015;92:393-402. <https://doi.org/10.1007/s11746-015-2596-5>
77. Lucas EW. Oilseeds and oil-bearing materials. In: Kulp K, editor. *Handbook of cereal science and technology*, revised and expanded. Boca Raton: CRC Press; 2000 Mar 28.p.297-362.
78. Sachdev N, Goomer S, Singh LR. Foxtail millet: a potential crop to meet future demand scenario for alternative sustainable protein. *J Sci Food Agric.* 2021;101(3):831-42. <https://doi.org/10.1002/jsfa.10716>
79. Shi J, Shan S, Li H, Song G, Li Z. Anti-inflammatory effects of millet bran derived-bound polyphenols in LPS-induced HT-29 cell via ROS/miR-149/Akt/NF-κB signaling pathway. *Oncotarget.* 2017;8(43):74582. <https://doi.org/10.18632/oncotarget.20216>
80. Fraterrigo Garofalo S, Tommasi T, Fino D. A short review of green extraction technologies for rice bran oil. *Biomass Convers Biorefin.* 2021;11:569-87. <https://doi.org/10.1007/s13399-020-00846-3>
81. Patel M, Naik SN. Gamma-oryzanol from rice bran oil-A review. *J Sci Ind Res.* 2004;63:569-78.
82. Zhang ZS, Wang LJ, Li D, Jiao SS, Chen XD, Mao ZH. Ultrasound-assisted extraction of oil from flaxseed. *Sep Purif Technol.* 2008;62(1):192-98. <https://doi.org/10.1016/j.seppur.2008.01.014>
83. Liu SX, Mamidipally PK. Quality comparison of rice bran oil extracted with d-limonene and hexane. *Cereal Chem.* 2005;82(2):209-15. <https://doi.org/10.1094/cc-82-0209>
84. Uquiche E, Jeréz M, Ortíz J. Effect of pretreatment with microwaves on mechanical extraction yield and quality of vegetable oil from Chilean hazelnuts (*Gevuina avellana* Mol). *Innov Food Sci Emerg Technol.* 2008;9(4):495-500. <https://doi.org/10.1016/j.ifset.2008.05.004>
85. Azmir J, Zaidul IS, Rahman MM, Sharif KM, Mohamed A, Sahena F, et al. Techniques for extraction of bioactive compounds from plant materials: A review. *J Food Eng.* 2013;117(4):426-36. <https://doi.org/10.1016/j.jfoodeng.2013.01.014>
86. Jones JM, Engleson J. Whole grains: benefits and challenges. *Annu Rev Food Sci Technol.* 2010;1(1):19-40. <https://doi.org/10.1146/annurev.food.112408.132746>
87. Dykes L, Rooney LW. Phenolic compounds in cereal grains and their health benefits. *Cereal Foods World.* 2007;52(3):105-11. <https://doi.org/10.1094/cfw-52-3-0105>
88. De Munter JS, Hu FB, Spiegelman D, Franz M, Van Dam RM. Whole grain, bran and germ intake and risk of type 2 diabetes: a prospective cohort study and systematic review. *PLoS Med.* 2007;4(8):e261. <https://doi.org/10.1371/journal.pmed.0040261>
89. Shan S, Li Z, Newton IP, Zhao C, Li Z, Guo M. A novel protein extracted from foxtail millet bran displays anti-carcinogenic effects in human colon cancer cells. *Toxicol Lett.* 2014a;227(2):129-38. <https://doi.org/10.1016/j.toxlet.2014.03.008>
90. Shan S, Li Z, Guo S, Li Z, Shi T, Shi J. A millet bran-derived peroxidase inhibits cell migration by antagonizing STAT3-mediated epithelial-mesenchymal transition in human colon cancer. *J Funct Foods.* 2014;10:444-55. <https://doi.org/10.1016/j.jff.2014.07.005>
91. Pang M, He S, Wang L, Cao X, Cao L, Jiang S. Physicochemical properties, antioxidant activities and protective effect against acute ethanol-induced hepatic injury in mice of foxtail millet (*Setaria italica*) bran oil. *Food Funct.* 2014;5(8):1763-70. <https://doi.org/10.1039/c4fo00106k>