

RESEARCH ARTICLE

# Exploring genetic diversity in Indian ginger (*Zingiber officinale* Rosc.) through microsatellite markers

BA Vaishnavi<sup>1</sup>, K Venkatesan<sup>1\*</sup>, N Senthil<sup>2</sup>, M Mohanalakshmi<sup>1</sup>, V Parantharan<sup>3</sup> & SP Thamaraiselvi<sup>4</sup>

<sup>1</sup>Department of Spices and Plantation Crops, Horticultural College and Research Institute, Tamil Nadu Agricultural University, Coimbatore-641 003, Tamil Nadu, India

<sup>2</sup>Centre for Plant Molecular Biology and Biotechnology, Tamil Nadu Agricultural University, Coimbatore-641 003, Tamil Nadu, India

<sup>3</sup>Department of Plant Pathology, Tamil Nadu Agricultural University, Coimbatore, 641 003, Tamil Nadu, India

<sup>4</sup>Department of Floriculture and Landscaping, Horticultural College and Research Institute, Tamil Nadu Agricultural University, Coimbatore, 641 003, Tamil Nadu, India

\*Email: [venkatesanhort@tnau.ac.in](mailto:venkatesanhort@tnau.ac.in)



## ARTICLE HISTORY

Received: 09 October 2024

Accepted: 13 November 2024

Available online

Version 1.0 : 24 December 2024



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Vaishnavi BA, Venkatesan K, Senthil N, Mohanalakshmi M, Parantharan V, Thamaraiselvi SP. Exploring genetic diversity in Indian ginger (*Zingiber officinale* Rosc.) through microsatellite markers. Plant Science Today.2024;11(sp4):01-07. <https://doi.org/10.14719/pst.5552>

## Abstract

Ginger (*Zingiber officinale* Rosc.) holds significant value as a rhizomatous spice known for its distinct taste and aroma. India is the world's largest producer, exporter, and consumer of ginger. Local names commonly know many ginger cultivars, and since the crop is propagated vegetatively, the chances of mixing are very high. This complicates the maintenance of purity and distinct characteristics of each variety. In the present study, thirty-two ginger genotypes were procured from various regions nationwide and assessed using 49 SSR (Simple Sequence Repeats) markers to evaluate their genetic diversity patterns. Among the 49 markers, 19 primers were amplified and produced 23 polymorphic bands, resulting in a polymorphism percentage of 52.63 %. Additionally, the unweighted pair group method (UPGMA) cluster analysis grouped the genotypes into five distinct clusters, with similarity coefficients ranging from 0.31 to 1.00. This suggests that each genotype exhibits substantial variability. Genotypes Maran and Acc. 581 showed a similarity value 1.00, indicating perfect similarity (100 %) in their genetic characteristics. These findings emphasize the critical role of SSR markers in germplasm conservation and highlight the potential for utilizing genetic diversity in breeding programs to develop improved ginger varieties with desirable traits.

## Keywords

germplasm fingerprinting; Indian ginger; molecular characterization; phylogenetic diversity; SSR markers

## Introduction

Ginger, botanically known as *Zingiber officinale* Rosc. (2n=22), is a member of the Zingiberaceae family and originates from Southeast Asia. The underground rhizome, renowned for its distinct flavour, aroma, and pungency, is the ginger of commerce. This versatile crop holds immense cultural, culinary, and medicinal significance. Rhizomes are marketed in different forms, including fresh, dried, and powder. Being one of the oldest and most significant spices, its cultivation traces back to ancient civilization (1). Ginger was among the earliest oriental spices introduced to Europe and remains highly sought after today (2).

In India, over 75 recognized varieties of ginger are under cultivation, alongside approximately 500 indigenous varieties being maintained and

grown across various regions (3, 4). These cultivars are generally named after their respective localities or areas where they are grown. However, identifying distinct morphological traits among these varieties poses challenges, leading to confusion among farmers and breeders regarding their maintenance and utilization (5).

Since ginger is a vegetatively propagated crop, genotype mixing is highly likely. While genotypes are identified based on morphological traits, assessing these traits is often tricky and subjective, especially considering the close genetic relationships between most cultivars. Additionally, phenotyping traits are commonly influenced by environmental variations (6). Though molecular markers, primarily through Marker Assisted Selection (MAS), enhance breeding programs, they supplement the selection process based on morphological and quantitative traits. These traits remain crucial, especially in studying Quantitative Trait Loci (QTLs) for variety development in breeding programs.

Molecular marker technology is a robust tool for assessing genetic variation in ginger genotypes, revealing substantial differences at the DNA level. This approach provides a direct, reliable, and efficient means for characterizing, conserving, and managing germplasm unaffected by environmental factors (7). Several genetic markers assess genetic variations in asexually reproducing plants (8). Today, molecular markers are commonly employed to differentiate genotypes in horticultural crops (9). Among these, simple sequence repeat (SSR) markers stand out due to their reproducibility and effectiveness in understanding genetic diversity and identifying cultivars. This is attributed to their co-dominance and highly polymorphic behaviour (10-12).


SSRs are extensively utilized as versatile tools in plant breeding programs and evolutionary studies because they can reveal diversity among cultivars (13). Studies have demonstrated that SSRs in coding sequences (EST-SSRs) exhibit polymorphism across species, making them valuable for phylogenetic analysis (14, 15). For instance, twelve SSR markers were used in a recent study to evaluate the genetic diversity among 100 ginger genotypes (16). 13 SSR primers produced 160 polymorphic bands and 63.29 % polymorphism (17). Considering the above context, this study utilized SSR markers to assess genetic differentiation among 32 ginger genotypes sourced from various regions across India. This is the first research effort to characterize extensive ginger germplasm diversity using molecular markers in Tamil Nadu, aiming to support the identification of genotypes with potential for commercial use and breeding programs.

## Materials and Methods

### Plant material

Thirty-two ginger genotypes were procured from various regions across the country and maintained at the germplasm centre of the Hybrid Rice Evaluation Centre, Gudalur, in the Nilgiris District of Tamil Nadu, and utilized in this study (Fig. 1 and Table 1). This centre is situated at

an elevation of 1300 m above mean sea level (MSL) and 11.5° N latitude and 76.5° E longitude. The climate is tropical and moderately humid, with temperatures between 18-28 °C and relative humidity around 70 %. The area receives an average annual rainfall of 2000 mm. The experimental plot has well-drained sandy loam soil.



**Fig. 1.** Geographical map of India, indicating areas of the collection of the 32 ginger genotype

### Genomic DNA extraction

Leaves from plants aged 45 to 60 days were selected for DNA isolation. The genomic DNA extraction process followed the CTAB (Cetyltrimethylammonium bromide) method, as outlined (18). One gram of fresh, young leaf material was finely ground to initiate the isolation procedure using a pestle and mortar. Subsequently, the DNA was extracted using the CTAB extraction buffer. Following extraction, DNA quantification was done using a nanodrop spectrophotometer.

### SSR analysis

Based on previous studies on ginger and turmeric, belonging to the family 'Zingiberaceae,' a set of SSR markers has been identified as reproducible and effective in assessing genetic diversity. In light of these findings, the same set of SSR markers was employed in the present study to evaluate the genetic diversity of ginger germplasm. 49 SSR primers were used, consisting of four genomic SSR primers, six EST SSR primers specific to ginger (19, 20), and 39 EST SSR primers derived from *Curcuma longa* (Table S1) (21). These primers were obtained from Bio-Serve Biotechnologies Pvt. Ltd., Hyderabad (Table S1). The PCR reaction mixture had a total volume of 20 µl and included the following components: 2.0 µl of dNTP (10 mM), 2 µl of primer (10 mM), 2.5 µl of 10X reaction buffer, 0.2 µl of Taq DNA

**Table 1.** List of 32 ginger germplasm and their source of collection

Sl.no.	Genotypes	Description	Longitude	Latitude
<b>Released varieties</b>				
1	IISR Mahima	Clonal selection, Released from ICAR-IISR, Kozhikode	75° 49' 22" E	11° 36' 20" N
2	IISR Rejatha	Clonal selection, Released from ICAR-IISR, Kozhikode	75° 49' 22" E	11° 36' 20" N
3	IISR Varada	Clonal selection, Released from ICAR-IISR, Kozhikode	75° 49' 22" E	11° 36' 20" N
4	IISR Vajra	Clonal selection, Released from ICAR-IISR, Kozhikode	75° 49' 22" E	11° 36' 20" N
5	IISR 1 GB	Clonal selection, Released from ICAR-IISR, Kozhikode	75° 49' 22" E	11° 36' 20" N
6	Sourabh	Mutant (V1S1 2) from HARS, OUAT, Pottangi, Odisha	82° 57' 53.82" E	18° 33' 47.376"N
7	Suruchi	Released from HARS, OUAT, Pottangi, Odisha	82° 57' 53.82" E	18° 33' 47.376"N
8	Suravi	An induced mutant of Rudrapur local, Released from HARS, OUAT, Pottangi, Odisha	82° 57' 53.82" E	18° 33' 47.376"N
9	Athira	Selection form somaclones of cultivar Maran, Released from KAU, Thrissur, Kerala	76° 12' 18.36" E	10° 32' 55.3" N
10	Aswathy	Single plant selection from somaclones of cultivar Rio-de-Janeiro. Released from KAU, Thrissur, Kerala	76° 12' 18.36" E	10° 32' 55.3" N
<b>Promising lines</b>				
11	Acc. 578			
12	Acc. 247	Promising germplasm collections, Sourced from National Active Germplasm Site (NAGS), ICAR-Indian Institute of Spices Research, Kerala	75° 49' 22" E	11° 36' 20" N
13	Acc. 581			
14	GCP 49			
<b>Local types</b>				
15	Maran	Landrace from Assam	92° 53' 15.7974" E	26° 11' 9.7212" N
16	Thandimaran	Landrace from Kerala	76° 15' E	10° 30' N
17	Rio De Janeiro	Popular ginger cultivar in South India, Collected from HREC, Gudalur	79° 45' 14.4" E	11° 45' 14.4" N
18	Gudalur Local	Local cultivars collected from Gudalur, Nilgiri District, Tamil Nadu	79° 45' 14.4" E	11° 45' 14.4" N
19	Nadia	Local cultivar, collected from Assam Agriculture University, Jorhat	94° 12' 3.5928" E	26° 45' 5.4888" N
20	Bhaise	Local cultivar, collected from KVK, Ranipool, North Eastern Sikkim	88° 36' 58.73" E	27° 21' 57.75" N
21	Nagaland Local	Local cultivar, collected from Dimapur, Nagaland	93° 44' 28.8" E	25° 54' 20.4" N
22	Manipur Local	Local cultivar, collected from Manipur	93° 43' E	24° 43' N
23	Himachal	Landrace from Himachal Pradesh	77° 12' E	31° 53' N
24	H D Kote Local	Local cultivar, collected from Mysore, Karnataka	76° 8' 38.5" E	12° 14' 32.3" N
25	Hassan Local	Local cultivar, collected from Hassan, Karnataka	76° 5' 48.1" E	12° 59' 52.3" N
26	Chikkamagalore Local-1	Local cultivar, collected from Chikkamagalore, Karnataka	75° 46' 44.7" E	13° 19' 41.7" N
27	Chikkamagalore Local-2	Local cultivar, collected from Chikkamagalore, Karnataka	75° 46' 44.7" E	13° 19' 41.7" N
28	Chikkamagalore Local-3	Local cultivar, collected from Chikkamagalore, Karnataka	75° 46' 44.7" E	13° 19' 41.7" N
29	Humnabadh Local	The popular local cultivar of North Karnataka, collected from KVK, Bidar, Karnataka	77° 32' 28.4" E	17° 56' 22.9" N
30	Thalavadi Local	Local cultivar, collected from Erode, Tamil Nadu	77° 16' 40.7" E	11° 36' 13.8" N
31	Gorubathane	Local cultivar, collected from BCKV, Mohanpur, West Bengal	88° 30' 51.6" E	23° 22' 20.4" N
32	Palakkad local	Local cultivar, collected from Palakkad, Kerala	76° 39' 59.6" E	10° 46' 52.8" N

polymerase (3 U/μl), 1.5 μl of template DNA (30-40 ng μl<sup>-1</sup>), and 11 μl of nuclease-free water. The PCR protocol, executed in a thermocycler, began with an initial denaturation at 94 °C for 5 minutes. This was followed by 35 cycles of 45 denaturation at 94 °C, 45 seconds at 94 °C, 45 seconds of annealing at 52-65 °C, and 1 minute extension at 72 °C. A final extension was performed at 72 °C for 20 minutes (7, 22). The amplified products were subsequently analyzed using 3 % agarose gel electrophoresis. The gel, stained with ethidium bromide, was visualized using a Bio-Imaging system (Syngene, GBOXCHEMI, England).

#### Data collection and statistical analysis

The SSR data generated for 32 genotypes were analyzed individually and collectively using statistical methods. The SSR products were visually evaluated and recorded based

on the bands' presence (1) or absence (0). The genetic association was assessed by calculating the Jaccard's similarity coefficient using the marker data, following the Unweighted pair group method (UPGMA), and the dendrogram was generated using NTSYS-PC Ver. 2.2. software (23). The Polymorphic information content (PIC) was calculated using the formula proposed (24).

### Results and Discussion

In this study, 49 SSR markers were employed to examine genetic diversity in 32 ginger genotypes. These SSR markers successfully identified polymorphic and monomorphic alleles in ginger genotypes, highlighting their effectiveness in assessing genetic variability among the genotypes. Primers were selected based on their ability to produce distinct, measurable, and consistent

amplified banding patterns. Out of the 49 SSR primers studied, 19 were successfully amplified, with each primer showing different amplification patterns. The amplified products ranged in size from 50 to 1200 bp. Five of these 19 primers (ZOC-100, ZOC-28, Clon 16, Clon 01, and CLEST 07) exhibited 100 % polymorphism, as detailed in Table 2. Out of the 41 total alleles, 23 were polymorphic. The primer GB-ZOM-103 detected the highest number of alleles (six), while primers such as ZOC-11, CSSR07, Clon 16, Clon 01, CSSR 24, CLEST SSR-02, and CLEST 07 produced the minimum number of alleles (one each). Similarly, comparable findings were noticed by using SSR markers to analyze the genetic variation and diversity within ginger collections (7). SSR primers demonstrated polymorphisms, indicating their effectiveness in detecting genetic variation in ginger genotypes. The high levels of polymorphism detected by the SSR markers indicate substantial genetic diversity among the genotypes studied (5,7,16). The number and frequency of alleles in a population affected the heterozygosity of the markers (6). The polymorphic information content (PIC), which measures gene diversity, averaged 0.70, ranging from 0.50 for the ZOC-100 primer to 0.98 for the GB-ZOM-103 primer (Table 3). Primers such as GB-ZOM-103, ZOC-11, and CSSR 07, which had PIC values

exceeding 0.90, were highly influential in distinguishing ginger genotypes. The high allele polymorphism could be attributed to factors such as large genome size, outcrossing behaviour, or inherent genetic diversity of the species. The primer GB-ZOM-103 exhibited the highest PIC value (0.98) with 6 alleles. A similar finding is consistent with the results (6, 7, 17,21).

The SSR patterns were identified based on the appearance or disappearance of bands produced by these primers. The SSR patterns of each genotype were then compared with those of the others, and an Euclidean distance matrix was computed for all 32 genotypes. The relationship among the genotypes was depicted in a dendrogram created using the UPGMA method (Fig. 2). From Jaccard's similarity coefficient, the dendrogram and the relatedness between the genotypes can be understood. This square matrix shows the similarity or distance between the 32 genotypes (Table 3). The similarity coefficient ranges from about 0.31 to 1.00, indicating a wide range of similarity among the genotypes. The genotypes near 0.31 showed less moderate similarity, while values closer to one indicate more similarity.

The genotypes were grouped into five main clusters, as illustrated in Fig 2. Cluster I, II, and III each contained a

**Table 2.** Polymorphism among the ginger genotypes detected by SSR markers

Primers	Repeat motifs	Primer sequence (5'-3')	Na	Ne	I	He	MB	PB	% MM	% PM	Allele range	PIC value
GB-ZOM-064	(GGA) <sub>5</sub>	F-CGTAGGATCTTCCCGACC R-CGAGTGAACCCATGGAGA	1	1.98	0.69	0.50	1	0	100	0	250	0.81
GB-ZOM-040	(GGC) <sub>5</sub>	F-TCTCCCTCTCGGATCCAT R-ATCCATTGCCTGATGGTG	2	1.85	0.65	0.46	1	1	50	50	210	0.75
GB-ZOM-103	(CCT) <sub>2</sub> (CTT) (CCT) <sub>4</sub>	F-GCTGCGGACTAAATGCTG R-ACGCTAGGGAACAGGGAG	6	1.65	0.62	0.50	1	5	16.66	83.33	150-1200	0.98
ZOC 11	(CTG) <sub>9</sub>	F-GGAGTATCTTCACCTCTGTGCC R-ACCCTCACCTTCTCCAAGC	1	1.82	0.64	0.45	1	0	100	0	250	0.93
ZOC-100	(CGA) <sub>9</sub>	F-CATCCCACTGGAAGCGTACAAC R-AGGTCGGAGGTGAAGTCTCTG	2	1.85	0.65	0.46	1	1	50	100	150-170	0.50
ZOC 98	(CT) <sub>14</sub>	F-GTAGTCCCCAACAGAACTCG R-AGATCGAGGTGGTCAGCAAT	3	1.58	0.55	0.36	1	2	33.33	66.66	250-280	0.68
ZOC 28	(GCCTTC) <sub>4</sub>	F-GCCTTCTTCGGAGTGTCTTT R-AACCAAGCCTAATCCAAAACC	3	1.55	0.54	0.35	0	3	0	100	150-280	0.65
CSSR07	(CCT) <sub>4</sub>	F-CGCAGCTGACACTTCTTCT R-AAGTCCGGGAGTTCTAAAGG	1	1.82	0.64	0.45	1	0	100	0	120-300	0.93
Clon 16	(AGA) <sub>8</sub>	F-TTGTGCCAAGTGAGGATTTG R-ACTCGCTTCTGCTCATCCAT	1	1.98	0.69	0.49	0	1	0	100	50-120	0.62
Clon 01	(TA) <sub>6</sub> TTG (TC) <sub>16</sub>	F-ACTGGACTGTCCGAGAGCAT R-TCGTTTAGCGACAACGGATT	1	1.36	0.43	0.26	0	1	0	100	120-250	0.46
CSSR 18	(AGA) <sub>13</sub>	F-CTTTTGGCTGATAATGGAAGG R-AAGAAAGAACTGACATCTCCG	2	1.98	0.69	0.49	1	1	50	50	250	0.59
CSSR 27	(GGAG) <sub>3</sub>	F-TCTCCCGAGTGATTCTTTGA R-TCTTCTCCATATCCCTGAT	1	1.00	0.45	0.42	1	0	100	0	180	0.81
CuMiSat-20	(AC) <sub>6</sub>	F-CGATACGAGTCCATCTCTTCG R-CCTTGCTTTGGTGGCTAGAG	3	1.95	0.68	0.48	2	1	66.66	33.33	200	0.90
CLEST SSR-02	(AAG) <sub>12</sub>	F-ACCGTAGCAAAGAAATAGGAC R-AAGGTGGAAGGAACTCG	1	1.00	0.44	0.41	1	0	100	0	250	0.90
Clon 14	(CTT) <sub>7</sub>	F-CTGCGGTCCAAGTACAAGATC R-CTAGCTGGTGGCGGTGGT	3	1.98	0.69	0.49	1	2	33.33	66.66	250	0.78
CLEST 07	(AT) <sub>8</sub> ...(T) <sub>10</sub>	F-CGCAGCTGACACTTCTTCT R-AAGTCCGGGAGTTCTAAAGG	1	1.82	0.64	0.45	0	1	0	100	150	0.90
Clon 09	(AC) <sub>14</sub>	F-GGAGGAGGCGAGTTGATTTGT R-GCTTTGGTGGCTAGAGATGC	3	1.65	0.58	0.39	2	1	66.66	33.33	50-120	0.65
Clon 15	(AAG) <sub>7</sub>	F: GTCGCCGATCTATTGTAGC R: GATCCATCTCCCTAAAGC	2	1.93	0.67	0.48	1	1	50	50	210	0.68
Clon 8	(GT) <sub>10</sub>	F-CCGGTGAGGGTGATATCTTG R-AAGCTCAAGCTCAAGCCAAT	3	1.99	0.69	0.50	1	2	33.33	66.66	120-300	0.87
<b>Total</b>	-		<b>41</b>	<b>1.72</b>	<b>0.61</b>	<b>0.44</b>	<b>17</b>	<b>23</b>	<b>949.97</b>	<b>999.97</b>		<b>13.42</b>
<b>Mean</b>	-		<b>2.15</b>				<b>0.89</b>	<b>1.26</b>	<b>49.99</b>	<b>52.63</b>		<b>0.70</b>

Note: Na-Total number of alleles; Ne-Number of effective alleles; I; Shannons' index; He-Expected heterozygosity; MB-Monomorphic bands; PB-Polymorphic bands; % MB-Per cent monomorphism; % PB-Per cent polymorphism; PIC-Polymorphism Information Content




Fig. 2. UPGMA dendrogram based on SSR markers using Jaccard's similarity coefficient

Table 3. Similarity matrix of ginger genotypes generated by SSR markers

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32			
1	1.00																																		
2	0.84	1.00																																	
3	0.73	0.57	1.00																																
4	0.73	0.68	0.89	1.00																															
5	0.68	0.63	0.94	0.94	1.00																														
6	0.68	0.73	0.73	0.73	0.78	1.00																													
7	0.63	0.57	0.89	0.89	0.94	0.73	1.00																												
8	0.68	0.63	0.84	0.84	0.89	0.68	0.84	1.00																											
9	0.78	0.73	0.84	0.84	0.89	0.78	0.84	0.89	1.00																										
10	0.68	0.73	0.73	0.73	0.78	0.78	0.84	0.78	0.78	1.00																									
11	0.68	0.73	0.52	0.52	0.57	0.78	0.52	0.68	0.68	0.68	1.00																								
12	0.68	0.73	0.63	0.73	0.68	0.78	0.63	0.68	0.68	0.78	0.68	1.00																							
13	0.57	0.63	0.52	0.52	0.57	0.78	0.63	0.57	0.68	0.78	0.68	0.68	1.00																						
14	0.68	0.73	0.73	0.73	0.78	0.78	0.73	0.78	0.78	0.89	0.68	0.78	0.78	1.00																					
15	0.68	0.73	0.73	0.73	0.78	0.78	0.84	0.78	0.78	1.00	0.68	0.78	0.78	0.89	1.00																				
16	0.73	0.78	0.68	0.68	0.73	0.94	0.68	0.73	0.84	0.84	0.84	0.84	0.84	0.84	0.84	1.00																			
17	0.73	0.68	0.68	0.78	0.73	0.73	0.78	0.73	0.84	0.73	0.63	0.73	0.73	0.63	0.73	0.73	1.00																		
18	0.57	0.73	0.42	0.42	0.47	0.57	0.52	0.47	0.57	0.68	0.57	0.57	0.68	0.68	0.68	0.63	0.52	1.00																	
19	0.73	0.78	0.57	0.57	0.63	0.73	0.57	0.73	0.73	0.73	0.84	0.63	0.63	0.73	0.73	0.78	0.57	0.73	1.00																
20	0.68	0.63	0.52	0.63	0.57	0.47	0.63	0.68	0.68	0.57	0.57	0.47	0.47	0.47	0.57	0.52	0.73	0.57	0.73	1.00															
21	0.57	0.63	0.52	0.52	0.57	0.78	0.63	0.68	0.68	0.78	0.78	0.68	0.78	0.68	0.78	0.84	0.73	0.68	0.84	0.68	1.00														
22	0.63	0.57	0.57	0.68	0.63	0.52	0.68	0.73	0.73	0.63	0.63	0.63	0.52	0.52	0.63	0.57	0.79	0.52	0.57	0.73	0.63	1.00													
23	0.73	0.89	0.57	0.68	0.63	0.73	0.57	0.73	0.73	0.73	0.73	0.73	0.63	0.73	0.73	0.78	0.68	0.63	0.78	0.63	0.73	0.68	1.00												
24	0.63	0.68	0.57	0.57	0.63	0.84	0.68	0.73	0.73	0.84	0.84	0.73	0.84	0.73	0.84	0.89	0.78	0.63	0.78	0.63	0.94	0.68	0.78	1.00											
25	0.68	0.63	0.63	0.63	0.68	0.57	0.73	0.78	0.78	0.68	0.68	0.47	0.57	0.57	0.68	0.63	0.73	0.68	0.84	0.89	0.78	0.73	0.63	0.73	1.00										
26	0.47	0.52	0.52	0.63	0.57	0.47	0.63	0.68	0.57	0.57	0.47	0.68	0.47	0.47	0.57	0.52	0.73	0.57	0.52	0.68	0.68	0.73	0.63	0.63	0.68	1.00									
27	0.36	0.52	0.52	0.52	0.57	0.68	0.52	0.57	0.47	0.57	0.57	0.57	0.57	0.57	0.57	0.63	0.42	0.47	0.63	0.47	0.68	0.31	0.63	0.63	0.47	0.57	1.00								
28	0.57	0.52	0.84	0.84	0.89	0.78	0.94	0.78	0.78	0.78	0.57	0.68	0.68	0.68	0.78	0.73	0.84	0.47	0.52	0.57	0.68	0.63	0.52	0.73	0.68	0.68	0.57	1.00							
29	0.63	0.68	0.78	0.78	0.84	0.84	0.89	0.73	0.73	0.94	0.63	0.73	0.73	0.84	0.94	0.78	0.68	0.63	0.68	0.52	0.73	0.57	0.68	0.78	0.63	0.52	0.63	0.84	1.00						
30	0.57	0.73	0.73	0.73	0.78	0.78	0.84	0.68	0.68	0.89	0.57	0.68	0.68	0.78	0.89	0.73	0.63	0.68	0.63	0.47	0.68	0.52	0.73	0.73	0.57	0.57	0.68	0.78	0.94	1.00					
31	0.73	0.78	0.78	0.78	0.84	0.84	0.78	0.84	0.84	0.94	0.73	0.84	0.73	0.94	0.94	0.89	0.68	0.63	0.78	0.52	0.73	0.57	0.78	0.78	0.63	0.52	0.63	0.73	0.89	0.84	1.00				
32	0.68	0.63	0.63	0.63	0.57	0.68	0.63	0.47	0.57	0.68	0.47	0.57	0.57	0.57	0.68	0.63	0.63	0.57	0.63	0.68	0.52	0.63	0.63	0.57	0.47	0.57	0.57	0.57	0.68	0.63	0.63	1.00			

Note: 1-IISR Varada; 2-IISR Rejatha; 3- IISR Mahima; 4- Nagaland local; 5-GCP-49; 6-Manipur local; 7 Sourabh; 8-Chikkamagalore local- 1; 9-Aswathy; 10-Maran; 11-Acc. 247; 12-Thadimaran; 13- IISR 1 GB; 14-IISR Vajra; 15-Acc. 581; 16-Gudalur Local; 17-Acc. 578; 18-Hassan local; 19-HD Kote local; 20-Athira; 21-Bhaise; 22-Chikkamagalore local-2; 23-Nadia; 24-Humnabad local; 25-Suruchi; 26-Rio De Janeiro; 27-Suravi; 28-Palakkad Local; 29-Himachal; 30-Chikkamagalore local-3; 31-Thalavadi local; 32-Gorubathan

single genotype, such as Suravi, Hassan local and Gorubathane, respectively, and shared a similarity ranging from 0.56 % to 0.61 %. Cluster IV was further divided into subclusters A and B with a similarity of 72 %. Cluster A comprised two genotypes (Athira and Suruchi) sharing a similarity of approximately 89 %. Cluster B included three genotypes, Acc. 578, CKM-2 (Chikkamagalore local-2), and Rio De Janeiro, with a similarity of 73 %. Cluster V, the

largest, comprised 24 genotypes and was further subdivided into Cluster C and D, which exhibited 68 % similarity. Cluster C was further split into Cluster E and F, with a similarity of 73 %. Cluster E entailed 11 genotypes, viz., Manipur Local, Gudalur Local, Bhaise, Humnabad Local, IISR 1GB, Maran, Acc. 581, Thalavadi Local, IISR Vajra, Himachal and CKM-3 (Chikkamagalore Local-3). Among 11 genotypes, Maran and Acc. 581 showed a

similarity value 1.00, indicating perfect similarity (100 %) in their genetic characteristics. Seven genotypes, such as Mahima, GCP-49, Nagaland Local, Sourabh, Palakkad Local, CKM-1 (Chikkamagalore Local-1), and Aswathy, fall under Cluster F sharing a similarity of 84 %. Cluster D was additionally divided into Cluster G and H, which shared a similarity of around 74 %. Cluster G included two genotypes (Acc. 247 & HD Kote Local) that shared 84 % similarity, while Cluster H included three genotypes (IISR Varada, IISR Rejatha & Nadia) with a similarity of 89 %.

Notably, two genotypes (Maran and Acc. 578) in Cluster IV exhibited a very close relationship, indicating high similarity in their genetic characteristics. Genotypes within the same cluster demonstrate closer genetic relationships, while those in different clusters or that diverge early (Suravi, Hassan Local, and Goribathane) may represent unique or highly divergent varieties. These unique varieties hold potential value for breeding programs to introduce new traits. Interestingly, most genotypes from nine states did not cluster exclusively by geographic origin. Instead, they formed mixed groups, with genotypes from various states intermingling. This finding underscores that the genetic diversity among ginger varieties is not primarily determined by their geographical source. Consequently, ginger genotypes have minimal or no location specificity (5-7,16).

Genetic variation is determined by geographical origin and human-driven selection processes, and the spread of desirable genetic variants plays a crucial role in determining the genetic makeup of plant populations (25). Accordingly, this diverse genetic material has the potential to enrich the local genetic resources and provide valuable insights for future breeding efforts. Genotypes that form distinct groups are promising germplasms that can be utilized to broaden the genetic base (26). Conversely, genetically similar genotypes are valuable in breeding programs for maintaining consistency, stabilizing desirable traits, and providing a reliable foundation for developing improved crop varieties.

## Conclusion

Using SSR markers has revealed significant genetic variation in ginger genotypes sourced from diverse agricultural and climatic regions across the country. Of the nineteen SSR primers tested, primers ZOC-100, ZOC-28, Clon 16, Clon 01, and CLEST 07 exhibited 100% polymorphism. Primers such as GB-ZOM-103, ZOC-11, and CSSR 07, with PIC values exceeding 0.90, were highly influential in distinguishing ginger genotypes. Regardless of their collection sites or geographical origins, the 32 ginger genotypes clustered into distinct groups. This clustering highlights each genotype's considerable variability and potential, influenced by genetic similarities, human selection, or environmental adaptation to local climates. These findings are invaluable for breeders and germplasm curators in characterizing, conserving, and protecting the ginger gene pool and implementing targeted crop improvement programs.

## Acknowledgements

We express our deepest gratitude to the scientists, farmers, and experts who generously provided the ginger genotypes for our study, enabling this investigation's successful completion.

## Authors' contributions

BAV conducted the experiment, recorded observations, and analyzed the data. KV guided the research, formulated the data, and approved the final manuscript. MM reviewed, edited, summarized, and revised the manuscript. NS, VP, and SPT assisted with editing and revising the manuscript.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interest to declare.

**Ethical issues:** None.

## Declaration of generative AI and AI-assisted technologies in the writing process

We acknowledge using the AI tool "Grammarly" to check the grammar in the manuscript.

## Supplementary data

Table S1. A list of 49 SSR primers was used to study the genetic diversity of ginger genotypes.

## References

- Purseglove JW, Brown EG, Green CL, Robbins SRJ. Ginger. In: Purseglove JW, editor. Spices. New York: Longman Inc;1981. p. 447-532.
- Nandi S, Rahman MZ, Siraj S, Ahmed S, Ahsan MA, Sultana N, Roy SK. Quality composition and biological significance of the Bangladeshi and China ginger (*Zingiber officinale* Rosc). J Microbiol Biotech Food Sci. 2013;2(5):2283-90. <https://office2.jmbfs.org/index.php/JMBFS/article/view/7109>.
- Rattan RS. Improvement of ginger. In: Chadha KL, Rethinam P, editors. Advances in Horticulture. Plantation in Spice Crops. New Delhi: Malhotra Publishing House;1994. p.333-34.
- Kumari M, Kumar M, Solankey SS. *Zingiber officinale* Rosc Ginger. In: Novak J, Bluthner WD, editors. Medicinal, Aromatic and Stimulant Plants-Handbook of Plant Breeding. Switzerland: Springer Cham; 2020. p. 12-16. [https://doi.org/10.1007/978-3-030-38792-1\\_20](https://doi.org/10.1007/978-3-030-38792-1_20).
- Das A, Gaur M, Barik DP, Subudhi E. Genetic diversity analysis of 60 ginger germplasm core accessions using ISSR and SSR markers. Plant Biosys. 2017;151(5):822-32. <https://doi.org/10.1080/11263504.2016.1211197>.
- Ismail NA, Rafii MY, Mahmud TMM, Hanafi MM, Miah G. Genetic diversity of torch ginger (*Etilingera elatior*) germplasm revealed by ISSR and SSR markers. Biomed Res Int. 2019;1:5904804. <https://doi.org/10.1155/2019/5904804>.
- Akshitha HJ, Prasath D, Umesha K, Mohammad Faisal P, Venkataravanappa V. Molecular characterization of ginger genotypes using RAPD and SSR markers. J Hort Sci. 2022;17:95-106. <https://doi.org/10.24154/jhs.v17i1.1052>

8. Bardakci F. Random amplified polymorphic DNA (RAPD) markers. *Turkish J Biol.* 2001; 25:185-96.
9. Ansari AM, Singh YV. Molecular diversity of brinjal (*Solanum melongena* L. and *S. aethiopicum* L.) genotypes revealed by SSR markers. *Electron J Plant Breed.* 2014;5(4):722-28.
10. Pomper KW, Lowe JD, Lu L, Crabtree SB, Dutta S. Characterization and identification of pawpaw cultivars and advanced selections by simple sequence repeat markers. *J American Society Hort Sci.* 2010;135:143-49. <https://doi.org/10.21273/jashes.135.2.143>.
11. Nas MN, Bolek Y Bardak A. Genetic diversity and phylogenetic relationships of *Prunus microcarpa* C.A. Mey. subsp. *tortosa* analyzed by simple sequence repeats (SSRs). *Sci Hortic.* 2011;127:220-27. <https://doi.org/10.1016/j.scienta.2010.09.018>.
12. Ismail NA, Rafii MY, Mahmud TMM, Hanafi MM, Miah G. Molecular markers: a potential resource for ginger genetic diversity studies. *Mol Bio Rep.* 2016;43:1347-58. <https://doi.org/10.1007/s11033-016-4070-3>.
13. Adato A, Sharon D, Lavi U. Application of DNA fingerprints for identification and genetic analyses of mango (*Mangifera indica*) genotypes. *J American Society Hort Sci.* 1995;120:259-64. <https://doi.org/10.21273/jashes.120.2.259>.
14. Gandhi SG, Awasthi P, Bedi YS. Analysis of SSR dynamics in chloroplast genomes of Brassicaceae family. *Bioinform.* 2010;5(1):16. <https://doi.org/10.6026/97320630005016>
15. Awasthi P, Singh A, Sheikh G, Mahajan V, Gupta AP, Gupta S, et al. Mining and characterization of EST-SSR markers for *Zingiber officinale* Roscoe with transferability to other species of Zingiberaceae. *Physiology Mol Bio Plants.* 2017;23:925-31. <https://doi.org/10.1007/s12298-017-0472-5>.
16. Gezahegn G, Rezene Y, Feyissa T. Genetic diversity analysis of Ethiopian ginger (*Zingiber officinale* Rosc) accessions using simple sequence repeat (SSR) markers. *Gen Res Crop Evol.* 2024;1-12. <https://doi.org/10.1007/s10722-024-01972-x>
17. Das A, Sahoo RK, Barik DP. Identification of duplicates in ginger germplasm collection from Odisha using morphological and molecular characterization. *Proc Natl Acad Sci India Sect B Biol Sci.* 2020;90:1057-66. <https://doi.org/10.1007/s40011-020-01178-y>.
18. Akshitha HJ. Molecular and chemo-profiling of ginger (*Zingiber officinale* Rosc) genotypes. PhD Dissertation. Bagalkot: University of Horticultural Sciences Press. 2018.
19. Lee SY, Fai WK, Zakaria M, Ibrahim H, Othman RY, Gwag JG, et al. Characterization of polymorphic microsatellite markers isolated from ginger (*Zingiber officinale* Rosc). *Mol Ecol Notes.* 2007;7(6):1009-11. <https://doi.org/10.1111/j.1471-8286.2007.01757.x>.
20. Anu C. Development of microsatellite markers for small cardamom (*Elettaria cardamomum* Maton.). PhD Dissertation. India: Mangalore University; 2010
21. Siju S, Dhanya K, Syamkumar S, Sheeja TE, Sasikumar B, Bhat AI, Parthasarathy VA. Development, characterization and utilization of genomic microsatellite markers in turmeric (*Curcuma longa* L.). *Biochem Syst Ecol.* 2010;38:641-46. <https://doi.org/10.1016/j.bse.2010.08.006>.
22. Nandkangre H, Konate MNG, Kambou DJ, Ouoba A, Zida WPFMS, Bado S, et al. Comparative study of genetic diversity parameters within ginger ecotypes from Burkina Faso using microsatellite markers developed on rice and ginger. *Afr J Biol Sci.* 2022;4(4):62-69. 10.33472/AFJBS.4.4.2022.62-69.
23. Rohlf FJ. NTSYS-pc numerical taxonomy and multivariate analysis system. New York. Appl Biostat Inc. 1992.
24. Anderson JA, Churchill GA, Autrique JE, Tanksley SD, Sorrels ME. Optimizing parental selection for genetic linkage maps. *Genome.* 1993;36:181-86. <https://doi.org/10.1139/g93-024>
25. Jatoi SA, Kikuchi A, San SY, Naing KW, Yamanaka S, Watanabe JA, Watanabe KN. Use of rice SSR markers as RAPD markers for genetic diversity analysis in Zingiberaceae. *Breed Sci.* 2006;56(2):107-11. <https://doi.org/10.1270/jsbbs.56.107>
26. Ashraf K, Ahmad A, Chaudhary A, Mujeeb M, Ahmad S, Amir M, Mallick N. Genetic diversity analysis of *Zingiber officinale* Rosc by RAPD collected from subcontinent of India. *Saudi J Biol Sci.* 2014;21(2):159-65. <https://doi.org/10.1016/j.sjbs.2013.09.005>