



REVIEW ARTICLE

Silicon as a key driver of phytolith and phytolith-occluded carbon sequestration for climate change mitigation in rice ecosystems - a review

Manonmani M¹, Christy Nirmala Mary P^{1*}, Sheeba S¹, Raghavan T², Amudha R³, Jasmine⁴, Jayabalakrishnan RM¹, Murugaragavan R¹, Ramamoorthy P⁵ & Rajkumar K⁶

¹Department of Soils and Environment, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Madurai 625 104, Tamil Nadu, India.

²Department of Agronomy, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Madurai 625 104, Tamil Nadu, India.

³Department of Seed Science and Technology, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Madurai 625 104, Tamil Nadu, India.

⁴Department of Biochemistry, Agricultural College and Research Institute, Tamil Nadu Agricultural University, Madurai 625 104, Tamil Nadu, India.

⁵Department of Soils Science, Don Bosco College of Agriculture, Takkolam, Ranipet 631 151, Tamil Nadu, India.

⁶Department of Soils Science, Tamil Nadu, India.

*Email: nirmalajeysee@gmail.com



ARTICLE HISTORY

Received: 11 October 2024

Accepted: 13 November 2024

Available online

Version 1.0 : 28 December 2024

Version 2.0 : 21 February 2025



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Manonmani M, Christy Nirmala Mary P, Sheeba S, Raghavan T, Amudha R, Jasmine, Jayabalakrishnan RM, Murugaragavan R, Ramamoorthy P, Rajkumar K. Silicon as a key driver of phytolith and phytolith-occluded carbon sequestration for climate change mitigation in rice ecosystems - a review. Plant Science Today.2024;11(sp4):01-10. <https://doi.org/10.14719/pst.5815>

Abstract

Anthropogenic activities have increased atmospheric greenhouse gases, especially carbon dioxide, leading to global warming and climate change in recent years. Silica, a principal element in the Earth's crust, is an essential resource for plant growth and development. Silica is absorbed by plants as mono-silicic acid (H₄SiO₄) and deposited as opal stone/phytolith in the cellular spaces and vascular bundles of plant parts. The highest phytolith content of rice plants is observed in straw, which is incorporated into the soil during harvest and acts as a resilience material. The Lsi1 and Lsi2 transporters and significant and secondary plant nutrients underwent Polymerization with adsorbed Si to form a phytolith structure in rice. These phytoliths give structural support, function as a defence mechanism, impart biotic and abiotic stresses and reduce the toxicity of certain heavy metals and salinization of soil. Phytolith-occluded carbon (PhytOC) is formed through occlusion, contributing to the geochemical carbon cycle and climate change mitigation. Over the past 60 years, the annual carbon sequestration varied between 0.81 × 10⁶ and 3.88 × 10⁶ Mg-e-CO₂ and a maximum of 37 × 10⁸ Mg-e-CO₂ within phytoliths in rice crops in China. Research in archaeology, palaeobotany, geology and paleoecology has focused on phytoliths because silica is a non-degradable base preserved as microfossils. Using silicon-rich organic and inorganic sources enhances the Aboveground Net Primary Productivity (ANPP) and Phytolith C sequestration in the rice ecosystem.

Keywords

carbon sequestration; climate change; phytolith; phytolith occluded carbon

Introduction

Augmenting greenhouse gases, especially carbon dioxide (CO₂), in the atmosphere due to human interventions, industrial activities, land use conversion and soil cultivation is the reason for the elevated temperatures observed both in the atmosphere and on the Earth's surface. Terrestrial carbon sequestration is crucial for mitigating climate change because 30% of CO₂ is emitted by terrestrial ecosystems (1, 2). Soil carbon storage decides the CO₂ level in the air and on earth

and indirectly affects climate change (3). Silica is considered a beneficial element. Some examples of silica-accumulating plants are rice, maize, wheat, sugarcane and barley. Plants absorb silicon as monosilicic acid (H_4SiO_4), especially in these silica-accumulating plants and assimilate it as a phytolith (4). Phytoliths are silica or opal stones formed during the biomineralization of silica in the plant, especially those that accumulate silica (5). Phytoliths predominantly comprise silicon dioxide (SiO_2), organic carbon (OC), water (H_2O), aluminium (Al), and iron (Fe), with proportions ranging from 66% to 91%, 1 to 6%, 0 to 11%, 0.01 to 4.55% and 0 to 2.1% respectively (6) and their composition varies with plant species (7).

Phytolith structure and composition, particularly in silica and calcium carbonate content, is crucial for identifying plant species and reconstructing past ecosystems. Different phytoliths reflect specific taxonomic attributes, aiding in understanding plant diversity and environmental adaptations. These variations also indicate ecological conditions, influencing nutrient cycling and habitat preferences (8). Phytoliths are composed of silica and other elements, can survive for long periods and are resilient to decomposition in rice ecosystems (9), helping plants withstand various stresses (10). Phytolith helps to overcome multiple biotic and abiotic stresses. Archaeology, palaeobotany, geology and pale ecological research on Phytoliths are carried out because silica is a non-degradable pedestal preserved as microfossils in various substrates (11). The maximum CO_2 sequestration of 0.26 Tg CO_2 per year in rice straw phytolith was noticed in China. Recent research pointed out that carbon is occluded as PhytOC during the absorption of phytolith in rice plant parts and is considered a carbon pool in the rice ecosystem and bang on the global carbon cycle (12).

PhytOC has persisted in soil for thousands of years, and research on radioactive carbon (C) isotopes has been carried out over the last decades. PhytOC carbon sequestration ranged from 7.2-8.8 kg/ha/yr (13). Soil properties, nutrients and climate are crucial in forming phytoliths and (PhytOC). Nutrient-rich soils, especially those containing sufficient silica and essential minerals, encourage strong phytolith production, which enhances carbon storage. In contrast, nutrient-poor or highly acidic soils can impede phytolith formation due to limited resources. Climatic factors, including moisture levels and temperature, also impact this process; generally, warmer and wetter climates promote the more significant accumulation of phytoliths and PhytOC. Therefore, both nutrient availability and climatic conditions work together to influence the efficiency of PhytOC formation in plants (12). The phytolith-enriched sheath and rice leaves can produce biochar as an amendment (14). PhytOC enhances phytolith content in soil and plant parts, varying from 0.94×10^6 tonnes CO_2 /yr to 2.17×10^6 tonnes CO_2 /yr. Fertilization and selection of rice varieties impact phytOC in geochemical sequestration and climate change mitigation (15).

Application of silicate fertilizers at 1.87 gm Si/pot and NPK fertilizers increased the uptake of rice's nitrogen, phosphorus, phytolith and phytOC production compared to the control (16). The application of silica fertilizers to paddy crops, especially in silica-deficient soil, increases the PhytOC fluxes from 8.04×10^4 to 7.41×10^5 Mg-e- CO_2 and basalt powder from 0.005 to 0.041 Mg CO_2 /ha (17, 18). Phytolith and PhytOC production was increased due to applying phosphorus fertilizer, silica-rich amendment,

organic mulching, basalt powder and NPK fertilizers (17-21). The rice phytolith potential in the long-term carbon sequestration process and carbon sequestration rate was 0.12 Mg-e- CO_2 /ha/year.

Implication of Silicon on Rice and its Phytolith Content

Rice is an indispensable crop for food and nutrition, covering around 164 million hectares globally as of 2014 (22). Silicon is crucial for silica-accumulating plants, especially in the Poaceae family, as it strengthens cell walls and enhances structural support. It improves resistance to pests, diseases and stresses like drought while aiding nutrient uptake for overall plant health. In crops like rice, wheat and maize, silicon is critical to maintaining productivity and resilience. Its protective role makes it vital for both agriculture and ecosystems. Crop wastes derived from rice (Rice straw and husks) are used for cattle feed and firewood in factories. In the Earth's crust, silicon comprises 27.7% of the Earth's crust by mass and ranks as the second most abundant element. Rice requires silica fertilization, removes 230-470 kg/ha of silicon and increases the availability of primary and secondary nutrients. Silicon induces photosynthesis, aids in phosphorus absorption from the soil, reduces nutrient toxicities and helps rice withstand water logging conditions (Fig. 1) (23). Silica positively influences phosphorus availability by absorption, solubilization and phosphorus release from plant to soil (24).

Silica occurs as silicon dioxide and silicate compounds in the Earth and plants. It should be absorbed as ortho silicic acid ($Si(OH)_4$), stored in the leaves, stems and culms of rice and other silica-accumulating plants (25,33,52). The assimilated silicic acid is stored as plant opal or silica stones or "Phytolith" in rice plants' leaves, stem culms and roots and is considered one of the silicon sources (26). Silica assimilates in the cell wall, intra and inter-cellular lumen and inter-cellular cortex region (27). Silicon provides mechanical strength and rigidity to the rice plants and prevents lodging (28). The production of chlorophyll and oxidative mechanisms are responsible for increasing the productivity of rice crops (29). Rice responds well to silicate fertilizers and the genes *LSi1*, *LSi2* and *LSi6* are responsible for transporting silicon from the roots to panicles through apoplast and vascular bundles (Fig. 2.) (30, 31). Polymerization occurs in the rice plant when excess mono silicic acid (H_4SiO_4) or $Si(OH)_4$ forms amorphous silica particles (Phytolith) in the cuticle along with cellulose and hemicellulose (32).

The deposited phytolith gives mechanical strength, decreases the feeding habits of herbivores and imparts biotic and biotic stress (Fig. 1) in rice plants. Phytoliths, when occluded with small amounts of carbon (C), nitrogen (N) and sulphur (S), enhance their functionality in various ways. Carbon contributes to phytoliths' structural integrity and stability, while nitrogen supports plant metabolism and promotes growth. Sulphur enhances phytolith resilience against biotic and abiotic stresses, including metal toxicity and salinization. Together, these elements improve soil health by facilitating nutrient cycling and interacting beneficially with soil microorganisms. Incorporating C, N and S into phytoliths supports plant health and environmental adaptability (34, 35).




Fig. 1 Rice crop needs silicon for its nutrition availability and for defence.




Fig. 2. Silicon aids in the establishment of rice phytolith production in different plant parts via silicon transporters.

Contrivance of Phytolith Advancement in Silica Accumulating Plants

Silicon is absorbed by the plant as mono-silicic acid in a solution with water and is transported through the vascular system (36). The silicic acid is accumulated as solid hydrated silica as phytolith (37). In the ocean, the phytolith gets re-solubilized and sequestered (38). The *Lsi₁* and *Lsi₂* transporters, along with primary and secondary nutrients, are involved in the accumulation of silica in soil solution and further Polymerization takes place during the formation of the phytolith structure (39). The development of phytolith structure includes silicon transfer from the root to vascular, silicon assimilation in the phytolith wall, advancement of phytolith structure by creating spaces for transport of silicon and other nutrients and nutrients isolation in the phytolith by desiccation and coagulation (Fig. 3). The organic matter, carbon, silicon and nutrients are fused through chemical bonds in rice (40). The gene transporters, viz., *OsLsi₁*, *OsLsi₂*, *OsLsi₃* and *OsLsi₆* are involved in the phytochrome gene structure for phytolith formation (41).




Fig. 3. Fruition of phytolith in plant's cell – silicon in the soil as a sign.

Modules of Phytolith in Different Organs of Rice Plants and its Cultivars

The maximum silica accumulation was noticed in the bracts, leaf blade and leaf sheath and the minimum was seen in the roots. The phytolith content of rice straw ranged from 29.3 to 143.3 mg/g with a mean value of 80 mg/g (15). The Al and Fe contents of the phytolith ranged from 0.003 to 0.009% and 0.012 and 0.034% respectively and varied with rice cultivars. The Al and Fe contents of the phytolith decreased in the order of sheaths > leaves = stems = grains and sheaths > stems = leaves = grains respectively (42). The Phytolith content of paddy straw was analyzed using EDS and distinct solid bodies indiscrete solid deposits around phytolith and small round spike-shaped phytolith (43) Fe, Al, K Si, O and C were observed (34).

Si, Al, Fe and water play a significant role in establishing organic carbon in rice phytoliths (44). The mean water content of the phytolith varied from 5.4 to 14.5% and the highest and lowest were noticed in stems and grains respectively. The average Al content of phytoliths ranged from 0.012 to 0.043% with the maximum phytolith Al content observed in rice sheaths and leaves, followed by grains and stems (42). The SiO₂ content of phytolith in grains, leaf sheaths, leaves and stems were 94.71, 92.50, 88.47 and 83.04% respectively. The Mg, Fe and OC contents in phytoliths varied from 0.0043 to 0.0082%, 0.017 to 0.009% and 1.93 to 2.46% respectively and the contents decreased from sheaths and leaves = grains > stems (42). The results on phytolith correlation with the elemental composition of rice cultivars envisaged that the R² value for Phytolith Vs. Al, Phytolith Vs. Fe and Phytolith Vs. Mg were 0.6494, 0.572 and 0.3194 respectively (15). Application of silicon (150, 300 and 600 kg/ha) increased the phytolith from 24.23 to 53.85 g/kg in rice and the maximum accumulation was in the stem (17).

A study on 71 different rice cultivars across nine provinces confirmed that the concentration of phytolith in rice straw ranged from 29.3 to 143.3 mg/g with a mean phytolith concentration of 80 mg/g (15). Geographical location, soil composition, climate and agricultural management practices influence the phytolith content in plants (45). The silicon dynamics in soil-rice systems varied based on the phytolith's solubility and parts of the plants. The phytolith content of straw, root, husk and grain in 15 rice cultivars ranged from 0.14 to 26.4

mg/g, with the highest content recorded in rice straw followed by husk, root and grain (Fig. 4).

Edifice of Rice Phytolith in Different Organs of Rice Plants and Disparity with Cultivars and Species

The rice phytolith shapes include double-peaked, bulliform and paralleled dumbbell structures, which vary across different organs (47). Bulliform cells, fucoid cells, or prickle hairs of rice phytoliths are found in the silica cells of vascular bundles or as silica bodies in various cells. The highest silica accumulation occurs during the reproductive period (32). The shapes of phytolith in husks and leaves were double-peaked and bulliform (49). The phytolith shapes were described by *O. sativa* and *O. rufipogon* as bulliform with unique buff-shaped ribbons encircling the half side and bulliform containing a spherical fan and round bow with a tubercle on the divergent side, respectively.

The contours of *O. sativa*, *O. rufipogon* and Xishuangbanna *O. rufipogon* from 20 different cultivars are presented. The results deliberated that bulliform phytoliths with minor variations in the lower part and lateral side were noticed in all the rice cultivars (48). The SEM images of paddy straw phytolith were observed. The shapes varied as globular (6-9 μ m), small round spike (2-3 μ m), a cluster of 2/3/5, cross-shaped, boat-shaped, 6-8 number of encircled lens-shaped, lens shape merged with round shape, cross-shaped on the veins (6-7 μ m), hook-shaped appear equidistantly on the veins, irregular silica bodies with small round (50) spike-shaped on smooth-surface and arrangement of regular pattern of globular (34). Double-peaked, bulliform and paralleled dumbbell phytolith shapes were observed in paddy straw (47).

Phytolith Occluded Carbon in Phytolith Assembly-A Basis for Terrestrial Soil Organic Carbon (SOC) Pool

Silica-accumulating plants have silica in their internal structure as mono silicic acid (Si (OH)₄), forming phytolith and occluding carbon within these phytoliths, called PhytOC. The PhytOC persist in soil for decades, contributing to carbon sequestration and climate change mitigation (51). The PhytOC content in rice varies by species, ranging from 0.13 to 0.21 mg/g, as determined by elemental analysis (53).




Fig. 4. Phytolith amassing in organs of rice cultivars and its impact on rice ecosystem.

The sub-compartmentation of OC and phytolith occluded carbon using Tomography. The results revealed that silica augmented compactly packed bundle-sheath cells mingled. Loosely arranged mesophyll cells are seen in rice plants (54). The CO_2 fluxes of $8.62 \mu\text{mol}/\text{m}^2/\text{s}$, which diffused over the rice field, were observed during the aerobic decomposition of assimilated fresh phytolith (55), ash phytolith and aged phytolith soil OC in the root zone and later it gets penetrated with water and then transmuted.

The phytolith accumulated in the rice straw underwent some chemical reactions, as shown in Fig. 5. The soluble forms of CO_2 , including H_2CO_3 , HCO_3^- and CO_3^{2-} are produced in this reaction. During its decomposition and its straw cycle, an acidic environment is created. The pH should be very low and drive the CO_2 buffering capacity. During this process, CO_2 buffer capacity is decreased. It enhances emission (Reaction. 1). If the paddy straw areas are burnt in the field, the organic matter will be converted to CO_2 and produce alkaline conditions with high pH. The open burning of rice straw in the field produces ash with high soil pH and increases the solubility of CO_2 . Finally, phytolith-occluded alkali ions are formed (56).

Plant growth environment (climate, surface vegetation, and soil) and plant genetics increase the carbon content of phytolith and are linked to carbon sequestration efficiency (57). Silicon absorption mechanisms and root silicification processes are responsible for phytolith-occluded carbon content in crops (58). The phytolith and phytOC ranged from 12.46 to 23.6% and 1.4 to 3.3% for straw, 5.5 to 11.4% and 1.1 to 2.7% for roots, 13.1 to 24.3% and 2.1 to 6.3% for husk and 0.2 to 1.9% and 0.7 to 1.4% for grains respectively (Fig. 6). Rice straw has high PhytOC than husk, roots and grains and high phytolith content was observed in husk and straw (59) (Fig. 6). The carbon sequestration rate of $76.4 \text{ g e-CO}_2 \text{ pot}^{-1}$ was noted in high phytolith content rice cultivars and mean PhytOC content was 5.0 mg/g (60).

The maximum Si-phytolith content was found in rice straw and the minimum in grains ranging from 14.47 to 26.39% in straw, 7.05 to 11.4% in roots, 13.13 to 24.38% in husk and 0.14 to 1.94% in grains (61). The rice husk recorded the highest PhytOC in phytoliths, followed by straw, root and grains, with the ranged values of 1.87-3.37%, 1.13-2.27%, 2.13-6.3% and 0.7-1.43% respectively. The utmost phytOC content of 3.37% was




Fig. 5. Chemistry behind phytolith occluded carbon development in soil -as a source of CO_2 flux in rice ecosystem.




Fig. 6. Rice plant organs with different rates of Phytolith and PhytOC absorption as silicon-enriched sustainable carbon.

noticed in the rice cv. M-219 straw than others (59).

A study on the relationship between phytolith content and carbon content in intensively cultivated soils in India showed no significant positive correlation between phytoliths and carbon content in phytoliths. However, a strong correlation was observed between phytolith and PhytOC content ($R^2 = 0.76$, $p < 0.01$), with a significant positive correlation ($R^2 = 0.40$, $p < 0.01$) between phytolith carbon content and PhytOC content in rice (17).

Preservation of Carbon Pool and Fluxes in Rice Ecosystem through Phytolith and Phytolith-Occluded Carbon for Climate Change Mitigation

Climate change relies upon the soil carbon pool, and phytOC is an essential source (63). PhytOC is very sustainable and skirmishes to decay compared to other crude C fractions in the soil (64). After the rice harvest, the straw waste is incorporated into the soil and the phytolith is released into the earth. The PhytOC in phytolith persists in the soil for over 6000 years (65). This persistence contributes significantly to terrestrial C sinks, accounting for up to 82% of total carbon sequestration (66). The sustainable carbon sequestration mechanism for atmospheric CO_2 is established through phytolith storage in plant biomass and soil pools. The phytolith-occluded carbon pools and fluxes varied from 0.01% to 0.04% and 0.14 - 2.32 teramoles (Tmol). Annual net land PhytOC accumulation rate was $< 0.01 - 0.05$ Tmol/yr. The soil had the most significant quantities of PhytOC from 8.69 - 128.25 Tmol and rice biomass from 1.11 - 23.05 Tmol.

Silicon fertilizers can be applied to recover phytolith carbon sink in rice ecosystems and reduce CO_2 emission into the atmosphere (67). The phytolith and PhytOC production flux in the rice ecosystem ranged from 643.30 to 1893.99 kg/ha/yr and 16.10 to 109.51 kg CO_2 ha/yr respectively. The phytolith and PhytOC production fluctuation were recorded to be the highest in long duration varieties than in medium and short duration rice cultivars. For example, Japonica rice varieties display a coefficient of variation of 29.16 Mg CO_2 /ha/year, compared to 24.1 Mg CO_2 /ha/year in Indica varieties (53).

Carbon sequestration Potential of Rice through Phytoliths and PhytOC Sink

PhytOC exhibits high resistance to decomposition, potentially persisting in soil for thousands of years after plant decay, thus serving as a significant long-term carbon sink (5, 68-71). This soil phytOC accumulation represents a crucial and enduring carbon sink mechanism in croplands and grass-dominated ecosystems (45, 60, 66, 72-73). The carbon sequestration potential in rice ranged from 0.03 to 0.13 Mg-e- CO_2 /ha/year across 5 rice cultivars. Globally, rice-growing regions sequester about 1.94×10^7 Mg-e- CO_2 annually. The Carbon fluctuation of rice phytoliths, annual carbon sequestration and sequestration within phytolith were 0.03–0.13 Mg-e- CO_2 /ha/year, 0.81×10^6 to 3.88×10^6 Mg-e- CO_2 and 2.37×10^8 Mg-e- CO_2 respectively during the 60 years in China (60). The annual C flux of rice phytoliths from the atmosphere was 1.94×10^7 Mg-e- CO_2 and is involved in climate change mitigation (74). The CO_2 sequestered in Chinese croplands was 4.39 ± 1.56 T g/yr and crop management practices could enhance the phytolith potential and carbon sequestration (75). The distribution of silica in various plant species found in China, which plays an important role in the carbon sequestration

process, was studied and results showed that herbs had the highest silicon accumulator, followed by trees and shrubs (76). The carbon sequestration potential in PhytOC (77) reached $0.48 \pm 0.20 \times 10^6$ t CO_2 /year and 44% rest upon the vegetation. Applying silicate fertilizers significantly enhanced the phytOC level and carbon bio-sequestration in paddy soil from 0.94×10^6 tonnes CO_2 /yr to 2.17×10^6 tonnes CO_2 /yr (46). The phytolith content of rice cultivars encompassed a carbon sequestration rate that ranged from 0.05 to 0.12 Mg-e- CO_2 /ha/year (79).

Worldwide, the annual latent sink rate of phytOC through rice phytolith was estimated at 16.4 Tg-e- CO_2 , with a biome total of 156.7 ± 91.6 Tg CO_2 /yr (46). The phytolith capacity in rice for carbon bio-sequestration ranged from 0.03 to 0.13 Mg-e- CO_2 /ha/year in 5 different rice cultivars (60). Silicate fertilizers boost PhytOC levels by supplying extra silica, which enhances phytolith formation in rice plants. This increased production leads to more significant carbon sequestration as more phytoliths accumulate in the soil. Silica also helps plants withstand environmental stresses, promoting overall health and carbon capture. Successful applications in rice-growing regions like Indonesia and China have shown improved PhytOC levels and soil health, demonstrating the potential of silicate fertilizers to enhance carbon sequestration while supporting sustainable agriculture (80-82). Phytolith C sequestration recovery in the rice ecosystem was achieved by selecting cultivars, applying silicon-rich fertilizers and applying organic manure, especially biochar (83-85).

The PhytOC content ranged from 0.002 to 0.82 mg/g in different rice organs and Carbon sequestration rates were 0.05 to 0.12 Mg-e- CO_2 /ha/year; the disparity might rest on silicon content and dry matter production. Rice plays a significant role in enduring C-seizure and climate transformation through the PhytOC Asia, Africa and South America contribute to the Phytolith carbon sink (46). Rice cultivars decide the PhytOC sequestration and the report of (46) depicted in (Fig. 7) that the Kavya cultivar recorded the lowest PhytOC sequestration value of 0.055 Mg-e- CO_2 /ha/year and the Surekha rice cultivar with highest PhytOC sequestration value of 0.119 Mg-e- CO_2 /ha/year. The PhytOC in soils through rice phytolith enhances the approximate universal annual latent sink rate of 1.64×10^7 Mg-e- CO_2 /ha/year. Application of slag-type silicate fertilizers in rice crops suppressed the methane emission from 158 mg CH_4 /m²/h to 116 mg CH_4 /m²/h due to the increased carbon dioxide activity and iron concentration in soil (86). Methane flux declined by 20% in control tillage and 36% in the no-tillage system in all seasons.

Bio-silicification Processes- a Dynamic Role in Long-Term Capture of Carbon Dioxide as Phytoc in Climate Change Mitigation

Phytolith occluded carbon refers to carbon restricted within phytoliths, which are microscopic silica structures found in plants (87) and derived from photosynthesis, converting atmospheric carbon dioxide into organic forms (88). PhytOC has the prospective a striking task in carbon sequestration in rice ecosystems (87) and indirectly on climate change mitigation (89). PhytOC operates as a sink for soil carbon, reducing CO_2 emissions and supplying nutrients for rice production. Improved management practices and mitigation technologies in the rice ecosystem could help reduce emissions (90).




Fig. 7. Rice cultivars with varied phytOC sequestration potential.

The development of no or no carbon fuel and natural and engineering strategies are needed to lower CO₂ emissions (91). PhytOC has come forward as a significant tool in the long-term seizure and retention of carbon (92, 93), presenting a promising boulevard for addressing climate change (57). Phytolith in silica accumulating plants acquire considerable latent for climate change and mitigation owing to their buoyancy against decomposition (94). The bio-silicification process forms PhytOC and is stored within rice plant tissue, representing a stable form of carbon sequestration (95).

Conclusion and Future Perspective

Rice is a staple food for almost all Asian countries, generating approximately one billion tons of straw annually. The incorporation of rice straw waste plays a crucial role in recycling nutrients, organic matter and phytoliths (silica bodies) deposited within rice plants' inter and intra-cellular tissues. Phytoliths, silica structures, persist in the soil for thousands of years and enhance photosynthesis while helping rice plants cope with various stresses. Phytoliths improve photosynthesis and conflict with multiple stresses in rice plants. Carbon, along with some nutrients deposited into the phytolith called phytOC, is involved in geochemical carbon sequestration in the rice ecosystem. Phytoliths have played an enormous role in the documentation of plants and the age of the microfossils.

Rice phytolith sequestered the carbon as 0.05-0.12 Mg-e-CO₂/ha/year and wide-ranging based on varieties, climate, rhizosphere and dry matter production. Approximately 1.64×10^7 Mg-e-CO₂/ha/year of estimated global annual potential sink rate of PhytOC in soils produced through rice phytolith. The enduring earthly carbon sequestration and climate change desires phytOC and rice growing countries viz., Asia, Africa and South America are principal carbon sink backers. Improved management practices and mitigation technologies to be practised in the rice ecosystem could help reduce emissions and help mitigate

climate change. Selecting high PhytOC-yielding rice cultivars presents a valuable opportunity to enhance the production of phytoliths, phytOC and the overall soil carbon pool. Applying silicon-rich fertilizer, bio wastes and organic matter improves the readily soluble Si pools and enhances phytolith, PhytOC production and soil physicochemical properties. There is an urgent need to augment phytolith and PhytOC production in the rice ecosystem, which occupies a large area worldwide, through aboveground Net Primary Productivity (ANPP) management practices.

However, research on the PhytOC accumulation in the rice ecosystem will be needed to standardize the phytOC production in different rice growing regions for global C sequestration, thereby reducing greenhouse gasses, especially carbon dioxide, in the atmosphere for mitigating climate change. The selection of long-duration rice cultivars with high PhytOC occlusion potential will pave the phytOC descend rate in India above the present sequestration rate.

Acknowledgements

Authors acknowledge the support rendered by the Department of Agricultural Microbiology, Agricultural College and Research Institute, Madurai.

Authors' contributions

MM drafted the original manuscript, investigated and analyzed. CNM supervised and did project administration reviewed and edited. SS collected resources and performed data curation. JB and MR collected resources and investigated. RT involved formal analysis. AR, JR, RM and RK did formal analysis.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None

Declaration of generative AI and AI-assisted technologies in the writing process

While preparing this work, the author(s) used Grammarly to correct the grammar. After using this tool/service, the author (s) reviewed and edited the content as needed and take(s) full responsibility for the publications' content.

References

- Le Quéré C, Moriarty R, Andrew RM, Canadell JG, Sitch S, Korsbakken JI, et al. Global carbon budget 2015. *Earth System Science Data*. 2015;7(2):349-96. <https://doi.org/10.5194/essd-7-349-2015>
- Tang X, Zhao X, Bai Y, Tang Z, Wang W, Zhao Y, et al. Carbon pools in China's terrestrial ecosystems: new estimates based on an intensive field survey. *Proceedings of the National Academy of Sciences*. 2018;115(16):4021-26. <https://doi.org/10.1073/pnas.1700291115>
- Chen L, Smith P, Yang Y. How has soil carbon stock changed over recent decades?. *Global Change Biology*. 2015;21(9):3197-99.
- Kaczorek D, Puppe D, Busse J, Sommer M. Effects of phytolith distribution and characteristics on extractable silicon fractions in soils under different vegetation—an exploratory study on loess. *Geoderma*. 2019;356:113917. <https://doi.org/10.1016/j.geoderma.2019.113917>
- Parr JF, Sullivan LA. Soil carbon sequestration in phytoliths. *Soil Biology and Biochemistry*. 2005;37(1):117-24. <https://doi.org/10.1016/j.soilbio.2004.06.013>
- Blecker SW, McCulley RL, Chadwick OA, Kelly EF. Biologic cycling of silica across a grassland bioclimosequence. *Global Biogeochemical Cycles*. 2006;20(3). <https://doi.org/10.1029/2006GB002690>
- Kameník J, Mizera J, Řanda Z. Chemical composition of plant silica phytoliths. *Environmental Chemistry Letters*. 2013;11:189-95.
- Shakoos S, Bhat M, Mir S, Soodan A. Investigations into phytoliths as diagnostic markers for the grasses (Poaceae) of Punjab. *Univ J Plant Sci*. 2014;2(6):107-22. <https://doi.org/10.13189/ujps.2014.020602>
- Matoh T, Kairusmee P, Takahashi E. Salt-induced damage to rice plants and alleviation effect of silicate. *Soil Science and Plant Nutrition*. 1986;32(2):295-304. <https://doi.org/10.1080/00380768.1986.10557506>
- Sharma R, Gupta A, Nandan G, Dwivedi G, Kumar S. Life span and overall performance enhancement of solar photovoltaic cell using water as coolant: a recent review. *Materials Today Proceedings*. 2018;5(9):18202-10. <https://doi.org/10.1016/j.matpr.2018.06.156>
- Albert RM, Bamford MK, Stanistreet IG, Stollhofen H, Rivera-Rondón CA, Njau JK, Blumenschine RJ. River-fed wetland palaeovegetation and palaeoecology at the HWK W site, Bed I, Olduvai Gorge. *Review of Palaeobotany and Palynology*. 2018;259:223-41. <https://doi.org/10.1016/j.revpalbo.2018.09.010>
- Song Z, McGrouther K, Wang H. Occurrence, turnover and carbon sequestration potential of phytoliths in terrestrial ecosystems. *Earth Science Reviews*. 2016;158:19-30. <https://doi.org/10.1016/j.earscirev.2016.04.007>
- Pan W, Song Z, Liu H, Van ZL, Li Y, Yang X, et al. The accumulation of phytolith-occluded carbon in soils of different grasslands. *Journal of Soils and Sediments*. 2017;17:2420-27. <https://doi.org/10.1007/s11368-017-1690-8>
- Nguyen AT, Nguyen MN. Straw phytolith for less hazardous open burning of paddy straw. *Scientific Reports*. 2019;9(1):20043. <https://doi.org/10.1038/s41598-019-56735-x>
- Tan L, Fan X, Yan G, Peng M, Zhang N, Ye M, et al. Sequestration potential of phytolith occluded carbon in China's paddy rice (*Oryza sativa* L.) systems. *Science of the Total Environment*. 2021;774:145696. <https://doi.org/10.1016/j.scitotenv.2021.145696>
- Singh P, Kumar V, Sharma J, Saini S, Sharma P, Kumar S, et al. Silicon supplementation alleviates the salinity stress in wheat plants by enhancing the plant water status, photosynthetic pigments, proline content and antioxidant enzyme activities. *Plants*. 2022;11(19):2525. <https://doi.org/10.3390/plants11192525>
- Sun X, Liu Q, Tang T, Chen X, Luo X. Silicon fertilizer application promotes phytolith accumulation in rice plants. *Frontiers in Plant Science*. 2019;10:425. <https://doi.org/10.3389/fpls.2019.00425>
- Guo F, Song Z, Sullivan L, Wang H, Liu X, Wang X, et al. Enhancing phytolith carbon sequestration in rice ecosystems through basalt powder amendment. *Science Bulletin*. 2015;60(6):591-97. <https://doi.org/10.1007/s11434-015-0729-8>
- Li Z, Guo F, Cornelis JT, Song Z, Wang X, Delvaux B. Combined silicon-phosphorus fertilization affects the biomass and phytolith stock of rice plants. *Frontiers in Plant Science*. 2020;11:67. <https://doi.org/10.3389/fpls.2020.00067>
- Song Z, Parr JF, Guo F. Potential of global cropland phytolith carbon sink from optimization of cropping system and fertilization. *PLoS One*. 2013;8(9):e73747. <https://doi.org/10.1371/journal.pone.0073747>
- Huang Z, Li Y, Chang SX, Jiang P, Meng C, Wu J, Zhang Y. Phytolith-occluded organic carbon in intensively managed Lei bamboo (*Phyllostachys praecox*) stands and implications for carbon sequestration. *Canadian Journal of Forest Research*. 2015;45(8):1019-25. <https://doi.org/10.1139/cjfr-2014-0495>
- Kumar V, Singh S, Kumar RM, Sharma S, Tripathi R, Nayak AK, Ladha JK. Growing Rice in eastern India: new paradigms of risk reduction and improving productivity. In: Mohanty S, et al, editors. *The future rice strategy for India*. Academic Press; 2017.p.221-58. <https://doi.org/10.1016/B978-0-12-805374-4.00008-7>
- Song A, Li P, Fan F, Li Z, Liang Y. The effect of silicon on photosynthesis and expression of its relevant genes in rice (*Oryza sativa* L.) under high-zinc stress. *PLoS One*. 2014;9(11):e113782. <https://doi.org/10.1371/journal.pone.0113782>
- Guan W, Ji F, Chen Q, Yan P, Zhang Q. Preparation and phosphorus recovery performance of porous calcium-silicate-hydrate. *Ceramics International*. 2013;39(2):1385-91. <https://doi.org/10.1016/j.ceramint.2012.07.079>
- Singh T, Singh P, Singh A. Silicon significance in crop production: special consideration to rice: an overview. *Pharma Innov*. 2021;10:223-29. <https://doi.org/10.22271/tpi.2021.v10.i3d.5776>
- Liang Y, Nikolic M, Bélanger R, Gong H, Song A. Silicon in agriculture: from theory to practice. *Springer Science*. 2015:123-42. <https://doi.org/10.1007/978-94-017-9978-2>
- Piperno DR. Paleoethnobotany in the neotropics from microfossils: new insights into ancient plant use and agricultural origins in the tropical forest. *Journal of World Prehistory*. 1998;12:393-449.
- Savant NK, Snyder GH, Datnoff LE. Silicon management and sustainable rice production. *Advances in Agronomy*. 1996;58:151-99. [https://doi.org/10.1016/S0065-2113\(08\)60255-2](https://doi.org/10.1016/S0065-2113(08)60255-2)
- Rastogi A, Yadav S, Hussain S, Kataria S, Hajjhashemi S, Kumari P, et al. Does silicon really matter for the photosynthetic machinery in plants?. *Plant Physiology and Biochemistry*. 2021;169:40-48. <https://doi.org/10.1016/j.plaphy.2021.11.004>
- Liang Y, Sun W, Zhu YG, Christie P. Mechanisms of silicon-mediated alleviation of abiotic stresses in higher plants: a review. *Environmental Pollution*. 2007;147(2):422-28. <https://doi.org/10.1016/j.envpol.2006.06.008>
- Gaur S, Kumar J, Kumar D, Chauhan DK, Prasad SM, Srivastava PK. Fascinating impact of silicon and silicon transporters in plants: a review. *Ecotoxicology and Environmental Safety*. 2020;202:110885. <https://doi.org/10.1016/j.ecoenv.2020.110885>

32. Rao GB, Susmitha P. Silicon uptake, transportation and accumulation in rice. *Journal of Pharmacognosy and Phytochemistry*. 2017;6(6):290-93.
33. Chaiwong N, Prom-U-Thai C. Significant roles of silicon for improving crop productivity and factors affecting silicon uptake and accumulation in rice: a review. *Journal of Soil Science and Plant Nutrition*. 2022;22(2):1970-82. <https://doi.org/10.1007/s42729-022-00787-y>
34. Anala R, Nambisan P. Study of morphology and chemical composition of phytoliths on the surface of paddy straw. *Springer Japan*; 2015 Oct. <https://doi.org/10.1007/s10333-014-0468-5>
35. Ma JF, Yamaji N. Silicon uptake and accumulation in higher plants. *Trends in Plant Science*. 2006;11(8):392-97.
36. Mitani N, Ma JF, Iwashita T. Identification of the silicon form in xylem sap of rice (*Oryza sativa* L.). *Plant and Cell Physiology*. 2005;46(2):279-83.
37. Haynes RJ. The nature of biogenic Si and its potential role in Si supply in agricultural soils. *Agriculture Ecosystems and Environment*. 2017;245:100-11. <https://doi.org/10.1016/j.agee.2017.04.021>
38. Frings PJ, Clymans W, Fontorbe G, Christina L, Conley DJ. The continental Si cycle and its impact on the ocean Si isotope budget. *Chemical Geology*. 2016;425:12-36.
39. Albert RM, Shahack-Gross R, Cabanes D, Gilboa A, Lev-Yadun S, Portillo M, et al. Phytolith-rich layers from the late Bronze and Iron Ages at Tel Dor (Israel): mode of formation and archaeological significance. *Journal of Archaeological Science*. 2008;35(1):57-75. <https://doi.org/10.1016/j.jas.2007.02.015>
40. Nguyen AT, Nguyen AM, Nguyen MT, Nguyen HT, Duong LT, Dinh VM, et al. The regulatory role of CO₂ on nutrient releases from ashed rice straw phytoliths. *Biogeochemistry*. 2022;160(1):35-47. <https://doi.org/10.1007/s10533-022-00938-4>
41. Nawaz MA, Zakharenko AM, Zemchenko IV, Haider MS, Ali MA, Imtiaz M, et al. Phytolith formation in plants: from soil to cell. *Plants*. 2019;8(8):249. <https://doi.org/10.3390/plants8080249>
42. Li Z, Song Z, Cornelis JT. Impact of rice cultivar and organ on elemental composition of phytoliths and the release of bio-available silicon. *Frontiers in Plant Science*. 2014;5:529. <https://doi.org/10.3389/fpls.2014.00529>
43. Gu Y, Zhao Z, Pearsall DM. Phytolith morphology research on wild and domesticated rice species in East Asia. *Quaternary International*. 2013;287:141-48. <https://doi.org/10.1016/j.quaint.2012.02.013>
44. Wüst RA, Bustin RM. Opaline and Al-Si phytoliths from a tropical mire system of West Malaysia: abundance, habit, elemental composition, preservation and significance. *Chemical Geology*. 2003;200(3-4):267-92. [https://doi.org/10.1016/S0009-2541\(03\)00196-7](https://doi.org/10.1016/S0009-2541(03)00196-7)
45. Parr JF, Sullivan LA. Phytolith occluded carbon and silica variability in wheat cultivars. *Plant and Soil*. 2011;342:165-71. <https://doi.org/10.1007/s11104-010-0680-z>
46. Prajapati K, Rajendiran S, Coumar V, Dotaniya ML, Kumar A, Kundu S. Carbon occlusion potential of rice phytoliths: implications for global carbon cycle and climate change mitigation. *Appl Ecol Environ Res*. 2016;14(2):265-81.
47. Zimin LI, Zhaoliang SO, Peikun JI. Biogeochemical sequestration of carbon within phytoliths of wetland plants: a case study of Xixi wetland, China. *Chinese Science Bulletin*. 2013;58(20):2480-87. <https://doi.org/10.1007/s11434-013-5785-3>
48. Wang Y, Hughes P, Niu H, Fan Y. A new method to improve the properties of recycled aggregate concrete: composite addition of basalt fiber and nano-silica. *Journal of Cleaner Production*. 2019;236:117602. <https://doi.org/10.1016/j.jclepro.2019.07.077>
49. Wang M, Lu Y, Lalevic B. Design considerations of silicon avalanche cathodes. *Journal of Vacuum Science and Technology B: Microelectronics and Nanometer Structures Processing Measurement and Phenomena*. 1993;11(2):426-28. <https://doi.org/10.1116/1.586876>
50. Cappers RT, Neef R. *Handbook of plant palaeoecology*. Barkhuis; 2021 Apr 28.
51. Rehman IU, Qader W, Dar RA, Rashid I, Shah RA. Phytolith-based paleoecological reconstruction from a loess-paleosol sequence in the Kashmir Himalaya, India. *Catena*. 2024;245:108318.
52. Kundu S, Rajendiran S, CoumarMV, Ajay. Effect of land use and management practices on quantifying changes of phytolith-occluded carbon in arable soils. *Carbon Management in Tropical and Sub-Tropical Terrestrial Systems*. 2020:37-55. https://doi.org/10.1007/978-981-13-9628-1_3
53. Sun X, Liu Q, Zhao G, Chen X, Tang T, Xiang Y. Comparison of phytolith-occluded carbon in 51 main cultivated rice (*Oryza sativa*) cultivars of China. *RSC Advances*. 2017;7(86):54726-33. <https://doi.org/10.1039/C7RA10685H>
54. Pise VH, Harlalka R, Thorat BN. *Drying of aromatic plant material for natural perfumes*. Boca Raton: CRC Press; 2023.
55. Dang QT, Nguyen AT, Nguyen AD, Nguyen NT, Dam TT, Tran TT, et al. Desilification of phytolith exacerbates the release of arsenic from rice straw. *Chemosphere*. 2024;349:140797. <https://doi.org/10.1016/j.chemosphere.2023.140797>
56. Nguyen MN, Dultz S, Guggenberger G. Effects of pretreatment and solution chemistry on solubility of rice-straw phytoliths. *Journal of Plant Nutrition and Soil Science*. 2014;177(3):349-59. <https://doi.org/10.1002/jpln.201300056>
57. Rajendiran S, Coumar MV, Kundu S, Dotaniya AM, Rao AS. Role of phytolith occluded carbon of crop plants for enhancing soil carbon sequestration in agro-ecosystems. *Current Science*. 2012:911-20.
58. Yuan F, Wang L, Sheng M. The application of crop phytoliths for reviewing occluded organic carbon. *Chinese Journal of Eco-Agriculture*. 2020;28(12):1932-40. <https://doi.org/10.13930/j.cnki.cjea.200307>
59. Prajapati K, Rajendiran S, Coumar MV, Dotaniya ML, Meena VD, Srivastava A, et al. Bio-sequestration of carbon in rice phytoliths. *National Academy Science Letters*. 2015;38(2):129-33. <https://doi.org/10.1007/s40009-014-0313-9>
60. Li Z, Song Z, Parr JF, Wang H. Occluded C in rice phytoliths: implications to biogeochemical carbon sequestration. *Plant and Soil*. 2013;370:615-23. <https://doi.org/10.1007/s11104-013-1661-9>
61. Yang X, Song Z, Qin Z, Wu L, Yin L, Van ZL, et al. Phytolith-rich straw application and groundwater table management over 36 years affect the soil-plant silicon cycle of a paddy field. *Plant and Soil*. 2020;454:343-58. <https://doi.org/10.1007/s11104-020-04656-4>
62. Kosten S, Roland F, Da Motta Marques DM, Van Nes EH, Mazzeo N, Sternberg LD, et al. Climate-dependent CO₂ emissions from lakes. *Global Biogeochemical Cycles*. 2010;24(2). <https://doi.org/10.1029/2009GB003618>
63. Prasad V, Stromberg CA, Alimohammadian H, Sahni A. Dinosaur coprolites and the early evolution of grasses and grazers. *Science*. 2005;310(5751):1177-80. <https://doi.org/10.1126/science.1118806>
64. Zheng Y, Matsui A, Fujiwara H. Phytoliths of rice detected in the Neolithic sites in the Valley of the Taihu Lake in China. *Environmental Archaeology*. 2003;8(2):177-83. <https://doi.org/10.1179/env.2003.8.2.177>
65. Zuo X, Lü H. Carbon sequestration within millet phytoliths from dry-farming of crops in China. *Chinese Science Bulletin*. 2011;56:3451-56. <https://doi.org/10.1007/s11434-011-4674-x>
66. Zhao Y, Song Z, Xu X, Liu H, Wu X, Li Z, et al. Nitrogen application increases phytolith carbon sequestration in degraded grasslands of North China. *Ecological Research*. 2016;31:117-23. <https://doi.org/10.1007/s11284-015-1320-0>
67. Wilding LP, Brown RE, Holowaychuk N. Accessibility and properties of occluded carbon in biogenetic opal. *Soil Science*. 1967;103(1):56-61.
68. Mulholland SC, Prior C. AMS radiocarbon dating of phytoliths. *MASCA Research Papers in Science and Archaeology*. 1993;10:21-23.

69. Song A, Ning D, Fan F, Li Z, Provance-Bowley M, Liang Y. The potential for carbon bio-sequestration in China's paddy rice (*Oryza sativa* L.) as impacted by slag-based silicate fertilizer. *Scientific Reports*. 2015;5(1):17354. <https://doi.org/10.1038/srep17354>
70. Song Z, Liu H, Si Y, Yin Y. The production of phytoliths in China's grasslands: implications to the biogeochemical sequestration of atmospheric CO₂. *Global Change Biology*. 2012;18(12):3647-53. <https://doi.org/10.1111/gcb.12017>
71. Parr J, Sullivan L, Quirk R. Sugarcane phytoliths: encapsulation and sequestration of a long-lived carbon fraction. *Sugar Tech*. 2009;11:17-21. <https://doi.org/10.1007/s12355-009-0003-y>
72. Parr J, Sullivan L, Chen B, Ye G, Zheng W. Carbon bio-sequestration within the phytoliths of economic bamboo species. *Global Change Biology*. 2010;16(10):2661-67. <https://doi.org/10.1111/j.1365-2486.2009.02118.x>
73. Rashid I, Mir SH, Zurro D, Dar RA, Reshi ZA. Phytoliths as proxies of the past. *Earth-Science Reviews*. 2019;194:234-50. <https://doi.org/10.1016/j.earscirev.2019.05.005>
74. Yang X, Song Z, Liu H, Bolan NS, Wang H, Li Z. Plant silicon content in forests of North China and its implications for phytolith carbon sequestration. *Ecological Research*. 2015;30:347-55. <https://doi.org/10.1007/s11284-014-1228-0>
75. Yang JL, Zhang GL. Silicon cycling by plant and its effects on soil Si translocation in a typical subtropical area. *Geoderma*. 2018;310:89-98.
76. Qi L, Li FY, Huang Z, Jiang P, Baoyin T, Wang H. Phytolith-occluded organic carbon as a mechanism for long-term carbon sequestration in a typical steppe: The predominant role of belowground productivity. *Science of the Total Environment*. 2017;577:413-17. <https://doi.org/10.1016/j.scitotenv.2016.10.206>
77. Song A, Ning D, Fan F, Li Z, Provance-Bowley M, Liang Y. The potential for carbon bio-sequestration in China's paddy rice (*Oryza sativa* L.) as impacted by slag-based silicate fertilizer. *Scientific Reports*. 2015;5(1):17354. <https://doi.org/10.1038/srep17354>
78. Kovács S, Kutasy E, Csajbók J. The multiple role of silicon nutrition in alleviating environmental stresses in sustainable crop production. *Plants*. 2022;11(9):1223. <https://doi.org/10.3390/plants11091223>
79. You H, Zheng L, You W, Tan F, Wang F, Cao Y, et al. The influence of exogenous nitrogen input on the characteristics of phytolith-occluded carbon in the *Kandelia obovata* Soil System. *Forests*. 2023;14(11):2202. <https://doi.org/10.3390/f14112202>
80. Anjum M, Nagabovanalli PB. Assessing production of phytolith and phytolith occluded carbon in aboveground biomass of intensively cultivated rice ecosystems in India. *Carbon Management*. 2021;12(5):509-19. <https://doi.org/10.1080/17583004.2021.1978552>
81. Li Z, Delvaux B. Phytolith-rich biochar: a potential Si fertilizer in desilicated soils. *GCB Bioenergy*. 2019;11(11):1264-82. <https://doi.org/10.1111/gcbb.12635>
82. Huang C, Wang L, Gong X, Huang Z, Zhou M, Li J, et al. Silicon fertilizer and biochar effects on plant and soil PhytOC concentration and soil PhytOC stability and fractionation in subtropical bamboo plantations. *Science of the Total Environment*. 2020;715:136846. <https://doi.org/10.1016/j.scitotenv.2020.136846>
83. Rehman IU, Qader W, Dar RA, Rashid I, Shah RA. Phytolith based paleoecological reconstruction from a loess-paleosol sequence in the Kashmir Himalaya, India. *Catena*. 2024;245:108318.
84. Ali MA, Lee CH, Lee YB, Kim PJ. Silicate fertilization in no-tillage rice farming for mitigation of methane emission and increasing rice productivity. *Agriculture Ecosystems and Environment*. 2009;132(1-2):16-22. <https://doi.org/10.1016/j.agee.2009.02.014>
85. Hodson MJ. The relative importance of cell wall and lumen phytoliths in carbon sequestration in soil: a hypothesis. *Frontiers in Earth Science*. 2019;7:167. <https://doi.org/10.3389/feart.2019.00167>
86. Bassham JA. The path of carbon in photosynthesis. *Scientific American*. 1962;206(6):88-104.
87. Bokor B, Santos CS, Kostoláni D, Machado J, da Silva MN, Carvalho SM, et al. Mitigation of climate change and environmental hazards in plants: potential role of the beneficial metalloids silicon. *Journal of Hazardous Materials*. 2021;416:126193. <https://doi.org/10.1016/j.jhazmat.2021.126193>
88. Hussain S, Peng S, Fahad S, Khaliq A, Huang J, Cui K, Nie L. Rice management interventions to mitigate greenhouse gas emissions: a review. *Environmental Science and Pollution Research*. 2015;22:3342-60. <https://doi.org/10.1007/s11356-014-3760-4>
89. Benbi DK. Greenhouse gas emissions from agricultural soils: sources and mitigation potential. *Journal of Crop Improvement*. 2013;27(6):752-72. <https://doi.org/10.1080/15427528.2013.845054>
90. Wang K, Sheng M, Wang L, He Y, Guo C. Response of soil phytolith occluded organic carbon accumulation to long-term vegetation restoration in Southwest China karst. *Land Degradation and Development*. 2022;33(16):3088-102. <https://doi.org/10.1002/ldr.4374>
91. Zhao E, Pang Z, Li W, Tan L, Peng H, Luo J, et al. Spatial variation in stability of wheat (*Triticum aestivum* L.) straw phytolith-occluded carbon in China. *Science of the Total Environment*. 2024;920:170909. <https://doi.org/10.1016/j.scitotenv.2024.170909>
92. Davamani V, John JE, Poornachandhra C, Gopalakrishnan B, Arulmani S, Parameswari E, et al. A critical review of climate change impacts on groundwater resources: a focus on the current status, future possibilities and role of simulation models. *Atmosphere*. 2024;15(1):122. <https://doi.org/10.3390/atmos15010122>
93. Lv H, Yang L, Zhou J, Zhang X, Wu W, Li Y, Jiang D. Water resource synergy management in response to climate change in China: from the perspective of urban metabolism. *Resources Conservation and Recycling*. 2020;163:105095. <https://doi.org/10.1016/j.resconrec.2020.105095>