



RESEARCH ARTICLE

# Effect of bioactive compounds on enzymatic regulation and shelf life of mango during post-harvest

Adarsh Balachandran<sup>1</sup>, V Sivakumar<sup>1\*</sup>, C Kavitha<sup>2</sup>, V Veeranan Arun Giridhari<sup>3</sup> & M Rajavel<sup>4</sup>

<sup>1</sup>Department of Fruit Science, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

<sup>2</sup>Department of Fruit Science, Tamil Nadu Agricultural University, Bhavanisagar 638 451, Tamil Nadu, India

<sup>3</sup>Center for Post harvest Technology, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

<sup>4</sup>Department of Crop Physiology, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

\*Email: [sivakumarv@tnau.ac.in](mailto:sivakumarv@tnau.ac.in)



## ARTICLE HISTORY

Received: 18 October 2024

Accepted: 26 November 2024

Available online

Version 1.0 : 27 December 2024



Check for updates

## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Balachandran A, Sivakumar V, Kavitha C, Giridhari VVA, Rajavel M. Effect of bioactive compounds on enzymatic regulation and shelf life of mango during post-harvest. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.5949>

## Abstract

The effects of melatonin, salicylic acid (SA) and brassinosteroids on the post-harvest quality and shelf life of two mango varieties, Alphonso and Sindhura, were investigated with respect to physiological, biochemical and enzymatic changes. The study assessed physiological loss in weight (PLW), Total Soluble Solids (TSS), titratable acidity, sugars, total phenolics, carotenoids, ripening related enzymes (pectin methylesterase, polygalacturonase, amylase and cellulase) and antioxidant enzymes (peroxidase, catalase and superoxide dismutase). According to the results, the shelf life of mango was considerably increased by 200 µM melatonin, 2.0 mM salicylic acid and 25 ppm brassinosteroids, which also decreased PLW to 9.81 %, 10.12 % and 9.79 % in Alphonso and 9.48 %, 9.51 % and 9.92 % in Sindhura. These treatments maintained a balance between TSS, sugars and acidity, while preserving higher concentration of total phenolics and carotenoids. Treated fruits exhibited lower ripening enzyme activity, particularly polygalacturonase, which declined by 16.7 %, 11.2 % and 17.0 % in Alphonso and 11.5 %, 10.9 % and 11.2 % in Sindhura with 200 µM melatonin, 2.0 mM salicylic acid and 25 ppm brassinosteroids treatments, respectively. Similarly, pectin methyl esterase (PME) was reduced by 12.2 %, 11.6 % and 10.5 % in Alphonso and 12.1 %, 10.3 % and 12.1 % in Sindhura corresponding to these treatments, delaying cell wall degradation and maintaining fruit firmness. Additionally, antioxidant enzymes were upregulated, mitigating oxidative stress by reducing reactive oxygen species (ROS) and preserving fruit quality. These treatments were effective in delaying ripening, enhancing stress resistance and extending the marketability of mangoes during storage. This study highlights the potential of melatonin, salicylic acid and brassinosteroids as eco-friendly alternatives to synthetic chemicals for improving post-harvest mango management.

## Keywords

brassinosteroids; enzymes; mango; melatonin; past-harvest; salicylic acid

## Introduction

Mango (*Mangifera indica* L.) is one of the most important tropical fruits globally, renowned for its rich flavour, nutritional value and economic significance (1). As a staple fruit in many countries, mangoes play a pivotal role in agricultural production, trade and dietary intake. Beyond their delightful taste, mangoes are a vital source of vitamins A and C, dietary fibre and antioxidants, making them highly valued in both fresh and processed forms (2).

Despite their importance, mangoes are highly perishable and face significant challenges in post-harvest handling. Issues such as rapid ripening, susceptibility to mechanical damage, microbial infection and post-harvest diseases drastically reduce fruit quality, shelf life and marketability. These issues result in substantial economic losses for producers and traders, particularly in tropical regions where proper storage and transportation infrastructure are limited (3).

In recent years, the use of bioactive compounds has emerged as a promising approach to mitigate post-harvest issues and extend the shelf life of fruits like mangoes. Compounds such as melatonin (4), salicylic acid (5) and brassinosteroids (6) have shown potential in enhancing fruit quality, delaying ripening and improving stress resistance. These natural compounds, often found in plants, offer an eco-friendly alternative to synthetic chemicals traditionally used in post-harvest treatments. The use of eco-friendly alternatives in post-harvest treatments is increasingly relevant in the context of sustainable agriculture and environmental conservation. Although traditional synthetic chemicals are effective, they pose risks such as environmental pollution, chemical residues on produce and potential harm to human health. (7). In contrast, natural bioactive compounds like melatonin, salicylic acid and brassinosteroids offer safe and sustainable solutions.

Melatonin (N-acetyl-5-methoxytryptamine) is a naturally occurring hormone in plants and animals, acting as an antioxidant and regulating plant growth, stress responses and defence mechanisms. Salicylic acid is a phenolic compound that plays a crucial role in plant defence against pathogens and in modulating stress tolerance. Brassinosteroids, a class of polyhydroxysteroids, are essential plant hormones that promote growth, development and enhance resistance to various environmental stresses. The exogenous application of these compounds has been proven to increase the shelf life and maintain quality in many fruits (8-10).

While there are numerous bioactive compounds with potential applications in post-harvest management, this study focuses on comparing the effectiveness of melatonin, salicylic acid and brassinosteroids. By evaluating their roles in improving the post-harvest quality and longevity of mangoes, this research aims to provide insights into the most effective treatments for minimizing post-harvest losses and maintaining fruit quality during storage.

## Materials and Methods

Mature green mangoes of the cultivars Alphonso and Sindhura, both known for their commercial importance, were collected from a farmer's field in Muthalamada, Palakkad district, Kerala. These fruits were carefully transported to the Postgraduate Analytical Laboratory, Horticultural College and Research Institute (HC&RI), Tamil Nadu Agricultural University (TNAU), Coimbatore. Upon arrival, the mangoes were cleaned with distilled water to remove any field debris before applying treatments. The experiment was laid out in a completely randomized design (CRD) with ten treatments, including a control group and three replications. The treatments encompass: T<sub>1</sub> - Control (distilled water), T<sub>2</sub> - melatonin (150 µM), T<sub>3</sub> - melatonin (200 µM), T<sub>4</sub> - melatonin (250 µM), T<sub>5</sub> - salicylic

acid (1.5 mM), T<sub>6</sub> - salicylic acid (2.0 mM), T<sub>7</sub> - salicylic acid (2.5 mM), T<sub>8</sub> - brassinosteroids (10 ppm), T<sub>9</sub> - brassinosteroids (25 ppm) and T<sub>10</sub> - brassinosteroids (50 ppm). Each treatment involved dipping the fruits for 30 seconds before allowing them to air-dry and storing them at an ambient temperature of 27±2 °C. Observations were made on various physiological and biochemical parameters at regular intervals.

### Shelf Life

Shelf life was recorded based on visual observations, noting the number of days the fruits remained marketable from the time of harvest. This parameter indicated the duration the fruits stayed marketable.

### Physiological loss in weight (PLW)

PLW was determined as the percentage reduction in the initial weight of the fruit compared to its weight at each measurement point. This calculation was made every three days and expressed as a percentage.

### Total Soluble Solids (TSS), Titratable acidity (TA), TSS/TA

Total Soluble Solids (TSS) were measured in °Brix using a hand refractometer (0-32 °Brix, Erma Inc., Tokyo) by placing 2-3 drops of mashed pulp juice on the prism. Titratable acidity (TA) was assessed by homogenizing 5 g of fruit with 30 ml of distilled water, filtering the mixture and titrating 5 ml of the filtrate against 0.1 N NaOH with phenolphthalein as an indicator until a pale pink colour appeared. TA was expressed as a percentage of citric acid using a standard formula (11). The TSS: acid ratio was calculated by dividing TSS by TA.

### Total sugars

Total sugars were estimated using the anthrone method and expressed in g/100 g. A 0.5 g sample was homogenized with 10 mL of 85 % ethanol, centrifuged at 10,000 rpm for 10 min and the supernatant was diluted to 100 mL. From this, 0.5 mL was mixed with 0.5 mL of distilled water and 4 mL anthrone reagent. After 10 min of incubation and heating in a boiling water bath for 8 min, the mixture was cooled to room temperature and absorbance was measured at 640 nm. Sugar content was calculated using a standard D-glucose calibration curve (12).

### Total phenol content

A 5 g sample was homogenized using 80 % methanol (or ethanol) for two to three grinding cycles. The homogenate was centrifuged at 10,000 rpm for 20 min, and the supernatant was collected and diluted to 50 mL with distilled water. A 0.5 mL aliquot of the supernatant was further diluted to 3 mL with distilled water, followed by the addition of 0.5 mL Folin-Ciocalteu reagent and a 3-min incubation. Subsequently, 2 mL of 20 % sodium carbonate solution was added, mixed thoroughly and incubated in a boiling water bath for 10 min. After cooling, the solution developed a blue colour and its absorbance was measured at 700 nm using a spectrophotometer (13).

### Carotenoids

Total carotenoid content was determined by homogenizing a 1 g sample with 10 mL of a 3:2 petroleum ether: acetone mixture, followed by centrifugation. The supernatant was diluted to 50 mL with the same solvent and the absorbance was measured at

450 nm using a UV spectrophotometer. Carotenoid concentration was calculated and expressed as micrograms per 100 g of sample (14).

### Enzyme Activity

Two grams of pulp were homogenized in a chilled mortar and pestle with 15 mL of sodium phosphate buffer (0.2 M, pH 7.0), supplemented with 0.2 mL EDTA and 0.2 mL cysteine-HCl. The homogenate was centrifuged at 15,000 rpm for 20 min at 4 °C and the clear supernatant was collected for enzyme activity assays, as per the method described by Srivastava and Dwivedi (15). Sodium phosphate buffer provides a stable pH environment to maintain enzyme activity during extraction. EDTA chelates divalent metal ions, preventing unwanted enzyme activation, while cysteine-HCl acts as a reducing agent to protect enzymes from oxidation. These components together stabilize the enzymes, ensuring accurate activity assays.

### Ripening Enzymes

**Pectin Methyl Esterase (PME) activity:** The activity of PME activity was conducted using a spectrophotometer set at 620 nm. A mixture of 0.1 mL bromothymol blue (0.01 %), 0.2 mL sodium chloride (0.15 M) and 1 mL pectin solution (0.01%) was prepared in a cuvette. The initial absorbance was recorded before adding 0.1 mL of enzyme extract. After the addition of the enzyme extract, absorbance was measured after three minutes. The change in absorbance was used to calculate enzyme activity, referencing a galacturonic acid standard curve (16). The results were expressed as milli equivalents of COOH released per minute per gram of fresh sample weight.

**Polygalacturonase activity:** To measure polygalacturonase activity, a reaction mixture was prepared by combining 0.2 mL of sodium acetate buffer (0.2 M), 0.1 mL of sodium chloride (2 M), 0.3 mL of polygalacturonic acid (1%) and 0.4 mL of enzyme extract, making a total volume of 1 mL. This mixture was incubated at room temperature for one hour. The reaction was then halted by adding 1 mL of 1% Di Nitro Salicylic acid (DNS) and placing the mixture in a boiling water bath. The absorbance of the solution was measured at 540 nm using a spectrophotometer against a blank. Enzyme activity was calculated using a galacturonic acid standard curve, as described in a study (17) and expressed in  $\mu\text{g}/\text{min/g}$  of the fresh weight of the sample.

**Amylase activity:** To determine amylase activity, a reaction mixture consisting of 0.5 mL phosphate buffer (0.2 M), 1 mL of 0.5% potato starch substrate and 1 mL of enzyme extract was incubated at 37 °C for one hour. The reaction was stopped by adding 1 mL of 1% Di Nitro Salicylic acid (DNS). Absorbance was then measured at 540 nm using a spectrophotometer, with a blank sample as the reference. The activity was calculated from a standard maltose curve, following the method of Bernfeld (18) and expressed in micrograms per minute per gram ( $\mu\text{g}/\text{min/g}$ ) of fresh weight of the sample.

**Cellulase activity:** The cellulase activity was determined by incubating a reaction mixture containing 1 mL of carboxymethyl cellulose, 1 mL of sodium acetate buffer (0.2 M) and 1 mL of enzyme extract at 37 °C for 60 minutes. To stop the reaction, 1 mL of 1% Di Nitro Salicylic acid (DNS) was added after placing the mixture in a boiling water bath for 5 minutes. Absorbance was recorded at 540 nm using a spectrophotometer, with a blank sample as the reference. The enzyme activity was calculated

from a standard glucose curve (19) and expressed as micrograms per minute per gram ( $\mu\text{g}/\text{min/g}$ ) of fresh weight of the sample.

### Antioxidant enzymes

**Peroxidase:** To initiate the reaction, 0.2 mL of hydrogen peroxide (0.1 M) was added to a cuvette containing 3.5 mL of phosphate buffer (0.2 M), 0.2 mL of enzyme extract and 0.1 mL of o-Dianisidine (1 mg/mL). The enzyme activity was monitored using a spectrophotometer at 430 nm for three min, with readings taken every 30 sec. The increase in absorbance over time was plotted, and the linear rise within one minute was used to calculate peroxidase activity (20). The activity was expressed as  $\Delta A/\text{min/g}$  of fresh sample weight.

**Catalase activity:** To initiate the reaction, 0.5 mL of hydrogen peroxide (12.5 mM) was added to a cuvette containing 1.5 mL phosphate buffer (0.2 M), 50  $\mu\text{L}$  of enzyme extract and 1.5 mL of distilled water. The cuvette was placed in a spectrophotometer and the enzyme activity was monitored at 240 nm for one min, with readings taken every 30 sec. The catalase activity, which caused a decrease in absorbance, was determined following the method of Aebi (21) and expressed as catalase activity/min/g of fresh sample weight.

**Superoxide dismutase activity:** To perform the reaction, 0.5 mL of enzyme extract, 0.1 mL of phenazine methosulphate (186  $\mu\text{M}$ ), 0.3 mL of nitroblue tetrazolium (300  $\mu\text{M}$ ), 0.2 mL of reduced nicotinamide adenine dinucleotide (780  $\mu\text{M}$ ) and 1.2 mL of sodium pyrophosphate buffer (0.052 M) were combined in a cuvette. The mixture was incubated at room temperature for 3 minutes. Following incubation, 4 mL of butanol and 1 mL of glacial acetic acid were quickly added while shaking vigorously. After allowing the layers to separate, the upper butanol phase was transferred to a clean cuvette and absorbance was measured at 520 nm with a spectrophotometer. Superoxide dismutase activity was calculated following the method of Kakkar et al. (22).

### Statistical Analysis

The experiment followed a completely randomized design (CRD). A one-way analysis of variance (ANOVA) was conducted to compare the treatment means, and the Least Significant Difference (LSD) test was applied to determine which treatments showed significant differences. Statistical significance was set at  $p \leq 0.05$ . All data analyses were performed using R Studio (version 4.3.1, released 2023-06-16) with relevant R packages for statistical computing.

## Results

### Shelf life

All treatments significantly influenced the storability of the fruits. Specifically, the treatments with 200  $\mu\text{M}$  melatonin ( $T_3$ ), 25 ppm brassinosteroid ( $T_9$ ) and 2.0 mM salicylic acid ( $T_6$ ) led to the longest shelf life for both varieties compared to the control and other treatments (Table 1). In the Alphonso variety,  $T_3$  resulted in the greatest increase in mean shelf life, extending it by 12.67 days, followed closely by  $T_6$  and  $T_9$ , each extending shelf life by 12.00 days. In the Sindhura variety,  $T_3$  also provided the longest shelf life extension of 12.00 days, with  $T_6$  and  $T_9$  closely following at 11.67 days at the end of the storage period.

**Table 1.** Effect of bioactive compounds on shelf life (days) in mango varieties under ambient storage

Treatments	Alphonso	Sindhura
T <sub>1</sub>	10.67 <sup>c</sup>	9.33 <sup>e</sup>
T <sub>2</sub>	11.33 <sup>bc</sup>	9.67 <sup>de</sup>
T <sub>3</sub>	12.67 <sup>a</sup>	12.00 <sup>a</sup>
T <sub>4</sub>	11.33 <sup>bc</sup>	10.33 <sup>cd</sup>
T <sub>5</sub>	11.00 <sup>bc</sup>	10.33 <sup>cd</sup>
T <sub>6</sub>	12.00 <sup>ab</sup>	11.67 <sup>ab</sup>
T <sub>7</sub>	11.33 <sup>bc</sup>	10.67 <sup>c</sup>
T <sub>8</sub>	11.00 <sup>bc</sup>	10.33 <sup>cd</sup>
T <sub>9</sub>	12.00 <sup>ab</sup>	11.67 <sup>ab</sup>
T <sub>10</sub>	11.33 <sup>bc</sup>	11.00 <sup>bc</sup>
SE(d)	0.58	0.42
CD	1.20	0.88

T<sub>1</sub> - Control (water dip); T<sub>2</sub> - Melatonin (150 µM); T<sub>3</sub> - Melatonin (200 µM); T<sub>4</sub> - Melatonin (250 µM); T<sub>5</sub> - Salicylic acid (1.5 mM); T<sub>6</sub> - Salicylic acid (2.0 mM); T<sub>7</sub> - Salicylic acid (2.5 mM); T<sub>8</sub> - Brassinosteroid (10ppm); T<sub>9</sub> - Brassinosteroid (25ppm); T<sub>10</sub> - Brassinosteroid (50 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).

**Table 2.** Effect of bioactive compounds on physiological loss in weight (%) in mango varieties under ambient storage

Treatments	Alphonso				Sindhura			
	Day 3	Day 6	Day 9	Day 12	Day 3	Day 6	Day 9	Day 12
T <sub>1</sub>	0.48 <sup>b</sup>	7.68 <sup>a</sup>	9.11 <sup>f</sup>	11.06 <sup>a</sup>	3.08 <sup>a</sup>	7.05 <sup>a</sup>	10.85 <sup>a</sup>	NA
T <sub>2</sub>	0.35 <sup>e</sup>	6.24 <sup>e</sup>	9.33 <sup>f</sup>	10.42 <sup>bc</sup>	2.62 <sup>c</sup>	5.42 <sup>d</sup>	8.62 <sup>d</sup>	NA
T <sub>3</sub>	0.26 <sup>g</sup>	5.29 <sup>g</sup>	8.12 <sup>g</sup>	9.81 <sup>d</sup>	1.75 <sup>g</sup>	5.13 <sup>e</sup>	7.09 <sup>f</sup>	9.48 <sup>b</sup>
T <sub>4</sub>	0.44 <sup>c</sup>	5.83 <sup>f</sup>	9.88 <sup>e</sup>	10.27 <sup>bc</sup>	2.29 <sup>e</sup>	5.69 <sup>bc</sup>	10.41 <sup>b</sup>	NA
T <sub>5</sub>	0.39 <sup>d</sup>	6.57 <sup>cd</sup>	10.30 <sup>d</sup>	10.17 <sup>c</sup>	2.49 <sup>d</sup>	5.63 <sup>c</sup>	9.57 <sup>c</sup>	NA
T <sub>6</sub>	0.26 <sup>g</sup>	5.85 <sup>f</sup>	10.34 <sup>d</sup>	10.12 <sup>cd</sup>	1.89 <sup>f</sup>	5.02 <sup>e</sup>	8.63 <sup>d</sup>	9.51 <sup>b</sup>
T <sub>7</sub>	0.35 <sup>e</sup>	6.29 <sup>de</sup>	10.61 <sup>cd</sup>	10.31 <sup>bc</sup>	2.26 <sup>e</sup>	5.84 <sup>b</sup>	9.69 <sup>c</sup>	NA
T <sub>8</sub>	0.39 <sup>d</sup>	7.12 <sup>b</sup>	10.88 <sup>bc</sup>	10.36 <sup>bc</sup>	3.06 <sup>ab</sup>	5.67 <sup>bc</sup>	8.55 <sup>d</sup>	NA
T <sub>9</sub>	0.31 <sup>f</sup>	5.77 <sup>f</sup>	11.15 <sup>b</sup>	9.79 <sup>d</sup>	2.01 <sup>f</sup>	5.11 <sup>e</sup>	7.59 <sup>e</sup>	9.92 <sup>a</sup>
T <sub>10</sub>	0.91 <sup>a</sup>	6.69 <sup>c</sup>	12.04 <sup>a</sup>	10.61 <sup>b</sup>	2.96 <sup>b</sup>	5.60 <sup>cd</sup>	10.62 <sup>ab</sup>	NA
SE(d)	0.01	0.15	0.20	0.17	0.058	0.09	0.17	0.09
CD	0.02	0.31	0.41	0.35	0.120	0.19	0.36	0.19

T<sub>1</sub> - Control (water dip); T<sub>2</sub> - Melatonin (150 µM); T<sub>3</sub> - Melatonin (200 µM); T<sub>4</sub> - Melatonin (250 µM); T<sub>5</sub> - Salicylic acid (1.5 mM); T<sub>6</sub> - Salicylic acid (2.0 mM); T<sub>7</sub> - Salicylic acid (2.5 mM); T<sub>8</sub> - Brassinosteroid (10ppm); T<sub>9</sub> - Brassinosteroid (25ppm); T<sub>10</sub> - Brassinosteroid (50 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).

**Table 3.** Effect of postharvest treatments on Total Soluble Solids (°Brix) in mango varieties under ambient storage

Treatments	Alphonso					Sindhura				
	Day 0	Day 3	Day 6	Day 9	Day 12	Day 0	Day 3	Day 6	Day 9	Day 12
T <sub>1</sub>	6.00 <sup>b</sup>	9.00 <sup>b</sup>	15.00 <sup>b</sup>	19.00 <sup>a</sup>	25.37 <sup>a</sup>	7.00 <sup>b</sup>	13.00 <sup>a</sup>	19.00 <sup>a</sup>	24.00 <sup>a</sup>	NA
T <sub>2</sub>	5.00 <sup>c</sup>	8.00 <sup>d</sup>	13.50 <sup>d</sup>	17.00 <sup>c</sup>	23.11 <sup>cd</sup>	6.00 <sup>c</sup>	10.50 <sup>ef</sup>	16.00 <sup>d</sup>	21.00 <sup>cd</sup>	NA
T <sub>3</sub>	5.00 <sup>c</sup>	7.00 <sup>e</sup>	12.50 <sup>e</sup>	15.00 <sup>e</sup>	20.07 <sup>h</sup>	7.00 <sup>b</sup>	10.00 <sup>f</sup>	15.00 <sup>e</sup>	20.50 <sup>de</sup>	22.00 <sup>b</sup>
T <sub>4</sub>	5.20 <sup>c</sup>	7.00 <sup>e</sup>	14.50 <sup>c</sup>	16.00 <sup>d</sup>	22.00 <sup>ef</sup>	6.00 <sup>c</sup>	9.00 <sup>g</sup>	15.00 <sup>e</sup>	21.80 <sup>bc</sup>	NA
T <sub>5</sub>	7.00 <sup>a</sup>	10.00 <sup>a</sup>	13.50 <sup>d</sup>	18.00 <sup>b</sup>	22.50 <sup>de</sup>	7.00 <sup>b</sup>	11.00 <sup>de</sup>	17.00 <sup>c</sup>	22.00 <sup>b</sup>	NA
T <sub>6</sub>	6.00 <sup>b</sup>	7.00 <sup>e</sup>	11.00 <sup>g</sup>	15.00 <sup>e</sup>	21.00 <sup>g</sup>	7.00 <sup>b</sup>	10.50 <sup>ef</sup>	15.50 <sup>de</sup>	20.00 <sup>e</sup>	23.00 <sup>a</sup>
T <sub>7</sub>	7.00 <sup>a</sup>	8.00 <sup>d</sup>	13.50 <sup>d</sup>	16.00 <sup>d</sup>	23.50 <sup>bc</sup>	8.00 <sup>a</sup>	12.00 <sup>bc</sup>	16.00 <sup>d</sup>	22.50 <sup>b</sup>	NA
T <sub>8</sub>	6.00 <sup>b</sup>	8.50 <sup>c</sup>	16.50 <sup>a</sup>	17.00 <sup>c</sup>	24.00 <sup>b</sup>	6.00 <sup>c</sup>	12.50 <sup>ab</sup>	18.00 <sup>b</sup>	23.50 <sup>a</sup>	NA
T <sub>9</sub>	5.00 <sup>c</sup>	7.00 <sup>e</sup>	12.00 <sup>f</sup>	15.00 <sup>e</sup>	21.50 <sup>fg</sup>	6.00 <sup>c</sup>	10.50 <sup>ef</sup>	14.00 <sup>f</sup>	21.00 <sup>cd</sup>	22.00 <sup>b</sup>
T <sub>10</sub>	5.00 <sup>c</sup>	8.00 <sup>d</sup>	12.40 <sup>ef</sup>	15.00 <sup>e</sup>	22.00 <sup>ef</sup>	7.00 <sup>b</sup>	11.50 <sup>cd</sup>	16.00 <sup>d</sup>	22.00 <sup>b</sup>	NA
SE(d)	0.12	0.18	0.23	0.36	0.42	0.16	0.26	0.29	0.46	0.20
CD	0.25	0.37	0.48	0.76	0.88	0.33	0.54	0.60	0.95	0.43

**Treatment details:** T<sub>1</sub> - Control (water dip); T<sub>2</sub> - Melatonin (150 µM); T<sub>3</sub> - Melatonin (200 µM); T<sub>4</sub> - Melatonin (250 µM); T<sub>5</sub> - Salicylic acid (1.5 mM); T<sub>6</sub> - Salicylic acid (2.0 mM); T<sub>7</sub> - Salicylic acid (2.5 mM); T<sub>8</sub> - Brassinosteroid (10 ppm); T<sub>9</sub> - Brassinosteroid (25 ppm); T<sub>10</sub> - Brassinosteroid (50 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).

### Physiological loss in weight

The percentage loss in weight (PLW) for Alphonso and Sindhura mangoes, as shown in Table 2, revealed that PLW increased progressively throughout the storage period. All treatments significantly reduced weight loss compared to the control, which exhibited the highest weight loss of 11.06% for Alphonso on the 12<sup>th</sup> day. In Alphonso, the treatment that was most effective in minimizing weight loss was T<sub>9</sub> (Brassinosteroid 25 ppm) with a loss of 9.79%, followed by T<sub>3</sub> (Melatonin 200 µM) with 9.81% and T<sub>6</sub> (Salicylic Acid 2.0 mM) with 10.12%. Similarly, for Sindhura, only these treatments (T<sub>3</sub>, T<sub>6</sub> and T<sub>9</sub>) maintained fruit quality up to the 12<sup>th</sup> day, while all other treatments led to deterioration by day 9. The 9<sup>th</sup> day, Sindhura exhibited the lowest weight losses in T<sub>3</sub> (7.09%), followed by T<sub>9</sub> (7.59%) and T<sub>6</sub> (8.63%).

### Total Soluble Solids (TSS)

Total Soluble Solids (TSS) increased during storage in both Alphonso and Sindhura (Table 3), but treatments T<sub>3</sub>, T<sub>6</sub> and T<sub>9</sub> effectively maintained lower Brix values. By day 12 in Alphonso, T<sub>3</sub>, T<sub>6</sub> and T<sub>9</sub> recorded 20.07, 21.00 and 21.50°Brix, respectively while control had 25.37°Brix. In Sindhura, on day 9, T<sub>3</sub>, T<sub>6</sub> and T<sub>9</sub> showed significantly lower readings (20.50, 20.00 and 21.00°Brix) than the control (24.00).



### Titrateable acidity (TA)

Titrateable acidity (TA) decreased throughout storage in both Alphonso and Sindhura, but treatments with melatonin, salicylic acid, and brassinosteroids significantly slowed the rate of decline as given in Table 4. Control fruits ( $T_1$ ) showed the lowest TA, with 0.23% on day 12 for Alphonso and day 9 for Sindhura. In Alphonso,  $T_3$  and  $T_6$  maintained higher TA levels (0.46% and 0.38%), while in Sindhura,  $T_3$  (0.84%),  $T_9$  (0.77%), and  $T_6$  (0.69 %) were most effective in preserving acidity.

### TSS:TA

The TSS: acid ratio increased steadily during storage in both Alphonso and Sindhura mangoes (Table 5). Treatments with melatonin (200  $\mu$ M), salicylic acid (2 mM) and brassinosteroids (25 ppm) significantly influenced this ratio, maintaining lower values compared to the control. By the end of storage, the highest ratios were in control fruits (110.11 for Alphonso and 95.48 for Sindhura).  $T_3$  and  $T_6$  maintained lower ratios in Alphonso (50.99 and 52.27), while in Sindhura,  $T_3$  (24.26),  $T_9$  (27.34) and  $T_6$  (28.93) were most effective on day 9.

### Total sugar

Total sugar content increased linearly from unripe to ripe stages across all treatments. The highest sugar content was observed in the control fruits on day 12 for Alphonso (20.05 g/100 g) and day 9 for Sindhura (13.33 g/100 g) as given in Table

6. The lowest sugar content was recorded in Alphonso on day 12 for  $T_9$  (16.10 g/100 g),  $T_3$  (16.22 g/100 g), and  $T_6$  (16.64 g/100 g). For Sindhura, the lowest content on day 9 was in  $T_3$  (10.19 g/100 g), followed by  $T_6$  (10.33 g/100 g) and  $T_9$  (10.51 g/100 g).

### Total phenol content

Total phenol content decreased during storage but remained higher in treated fruits compared to the control as given in Table 7. By day 12 in Alphonso,  $T_3$  (20.05),  $T_9$  (19.40) and  $T_4$  (19.25) showed significantly higher phenol levels than the control (13.05). In Sindhura, on day 9,  $T_3$  (16.00),  $T_6$  (15.65) and  $T_9$  (15.45) maintained higher phenol content, while the control recorded the lowest (10.15).

### Carotenoids

Carotenoid content increased in both treated and untreated Alphonso and Sindhura mangoes during storage, with treated fruits showing lower levels than the control (Table 8). By day 12 in Alphonso,  $T_3$  (21,133.43  $\mu$ g/100g),  $T_6$  (21,773.26  $\mu$ g/100g) and  $T_9$  (21,288.92  $\mu$ g/100g) recorded the lowest carotenoid levels. In Sindhura, viable treatments on day 12 included  $T_3$  (19,880.33  $\mu$ g/100g),  $T_6$  (20,468.76  $\mu$ g/100g) and  $T_9$  (20,666.03  $\mu$ g/100g), while the highest carotenoid content was observed in the control for both varieties.

**Table 4.** Effect of postharvest treatments on titrateable acidity (%) in mango varieties under ambient storage

Treatments	Alphonso					Sindhura				
	Day 0	Day 3	Day 6	Day 9	Day 12	Day 0	Day 3	Day 6	Day 9	Day 12
$T_1$	3.23 <sup>de</sup>	1.99 <sup>e</sup>	1.38 <sup>f</sup>	0.61 <sup>g</sup>	0.23 <sup>d</sup>	2.53 <sup>c</sup>	1.69 <sup>f</sup>	0.99 <sup>f</sup>	0.23 <sup>g</sup>	NA
$T_2$	3.45 <sup>c</sup>	2.61 <sup>cd</sup>	1.69 <sup>e</sup>	0.84 <sup>e</sup>	0.31 <sup>c</sup>	2.61 <sup>bc</sup>	1.99 <sup>cd</sup>	1.46 <sup>a</sup>	0.38 <sup>f</sup>	NA
$T_3$	3.69 <sup>b</sup>	2.99 <sup>a</sup>	2.15 <sup>a</sup>	1.22 <sup>a</sup>	0.46 <sup>a</sup>	2.69 <sup>ab</sup>	2.22 <sup>a</sup>	1.46 <sup>a</sup>	0.84 <sup>a</sup>	0.38 <sup>a</sup>
$T_4$	3.30 <sup>cde</sup>	2.69 <sup>c</sup>	1.92 <sup>cd</sup>	1.07 <sup>c</sup>	0.31 <sup>c</sup>	2.61 <sup>bc</sup>	2.07 <sup>bc</sup>	1.38 <sup>b</sup>	0.54 <sup>e</sup>	NA
$T_5$	3.15 <sup>e</sup>	2.30 <sup>f</sup>	1.69 <sup>e</sup>	0.92 <sup>d</sup>	0.31 <sup>c</sup>	2.53 <sup>c</sup>	1.92 <sup>de</sup>	0.92 <sup>g</sup>	0.61 <sup>d</sup>	NA
$T_6$	3.45 <sup>c</sup>	2.84 <sup>b</sup>	1.99 <sup>bc</sup>	1.15 <sup>b</sup>	0.38 <sup>b</sup>	2.76 <sup>a</sup>	2.15 <sup>ab</sup>	1.38 <sup>b</sup>	0.69 <sup>c</sup>	0.23 <sup>c</sup>
$T_7$	3.45 <sup>c</sup>	2.99 <sup>a</sup>	1.84 <sup>d</sup>	1.07 <sup>c</sup>	0.23 <sup>d</sup>	2.69 <sup>b</sup>	1.92 <sup>de</sup>	1.30 <sup>c</sup>	0.53 <sup>e</sup>	NA
$T_8$	3.38 <sup>cd</sup>	2.46 <sup>e</sup>	1.84 <sup>d</sup>	0.69 <sup>f</sup>	0.23 <sup>d</sup>	2.61 <sup>bc</sup>	1.84 <sup>e</sup>	1.07 <sup>e</sup>	0.38 <sup>f</sup>	NA
$T_9$	3.76 <sup>ab</sup>	2.84 <sup>b</sup>	2.07 <sup>ab</sup>	1.23 <sup>a</sup>	0.31 <sup>c</sup>	2.69 <sup>ab</sup>	2.07 <sup>bc</sup>	1.46 <sup>a</sup>	0.77 <sup>b</sup>	0.31 <sup>b</sup>
$T_{10}$	3.92 <sup>a</sup>	2.53 <sup>de</sup>	1.92 <sup>cd</sup>	1.07 <sup>c</sup>	0.23 <sup>d</sup>	2.61 <sup>bc</sup>	1.99 <sup>cd</sup>	1.15 <sup>d</sup>	0.61 <sup>d</sup>	NA
SE(d)	0.07	0.06	0.04	0.02	0.01	0.07	0.04	0.04	0.02	0.00
CD	0.16	0.12	0.08	0.05	0.02	0.14	0.09	0.07	0.03	0.01

**Treatment details:**  $T_1$  - Control (water dip);  $T_2$  - Melatonin (150  $\mu$ M);  $T_3$  - Melatonin (200  $\mu$ M);  $T_4$  - Melatonin (250  $\mu$ M);  $T_5$  - Salicylic acid (1.5 mM);  $T_6$  - Salicylic acid (2.0 mM);  $T_7$  - Salicylic acid (2.5 mM);  $T_8$  - Brassinosteroid (10 ppm);  $T_9$  - Brassinosteroid (25 ppm);  $T_{10}$  - Brassinosteroid (50 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).

**Table 5.** Effect of postharvest treatments on TSS: acid ratio in mango varieties under ambient storage

Treatments	Alphonso					Sindhura				
	Day 0	Day 3	Day 6	Day 9	Day 12	Day 0	Day 3	Day 6	Day 9	Day 12
$T_1$	1.86 <sup>c</sup>	4.50 <sup>a</sup>	10.85 <sup>a</sup>	30.92 <sup>a</sup>	110.11 <sup>a</sup>	2.76 <sup>b</sup>	7.69 <sup>a</sup>	19.03 <sup>a</sup>	95.48 <sup>a</sup>	NA
$T_2$	1.45 <sup>f</sup>	3.06 <sup>a</sup>	7.99 <sup>a</sup>	20.12 <sup>a</sup>	75.23 <sup>cd</sup>	2.29 <sup>e</sup>	5.26 <sup>e</sup>	10.96 <sup>f</sup>	54.69 <sup>c</sup>	NA
$T_3$	1.36 <sup>g</sup>	2.38 <sup>a</sup>	5.81 <sup>a</sup>	12.20 <sup>a</sup>	50.99 <sup>e</sup>	2.60 <sup>cd</sup>	4.49 <sup>h</sup>	10.28 <sup>g</sup>	24.26 <sup>h</sup>	57.29 <sup>c</sup>
$T_4$	1.57 <sup>e</sup>	2.60 <sup>a</sup>	7.55 <sup>a</sup>	14.88 <sup>a</sup>	71.61 <sup>d</sup>	2.29 <sup>e</sup>	4.34 <sup>h</sup>	10.85 <sup>f</sup>	40.55 <sup>de</sup>	NA
$T_5$	2.22 <sup>a</sup>	4.34 <sup>a</sup>	7.99 <sup>a</sup>	19.53 <sup>a</sup>	73.24 <sup>d</sup>	2.76 <sup>b</sup>	5.73 <sup>d</sup>	18.44 <sup>b</sup>	35.81 <sup>f</sup>	NA
$T_6$	1.74 <sup>d</sup>	2.46 <sup>a</sup>	5.51 <sup>a</sup>	13.02 <sup>a</sup>	52.27 <sup>e</sup>	2.53 <sup>d</sup>	4.88 <sup>g</sup>	11.21 <sup>f</sup>	28.93 <sup>g</sup>	91.15 <sup>a</sup>
$T_7$	2.03 <sup>b</sup>	2.67 <sup>a</sup>	7.32 <sup>a</sup>	14.88 <sup>a</sup>	91.15 <sup>b</sup>	2.98 <sup>a</sup>	6.25 <sup>c</sup>	12.25 <sup>e</sup>	41.85 <sup>d</sup>	NA
$T_8$	1.78 <sup>d</sup>	3.46 <sup>a</sup>	8.95 <sup>a</sup>	24.59 <sup>a</sup>	93.32 <sup>b</sup>	2.29 <sup>e</sup>	6.78 <sup>b</sup>	16.74 <sup>c</sup>	61.19 <sup>b</sup>	NA
$T_9$	1.33 <sup>gh</sup>	2.46 <sup>a</sup>	5.79 <sup>a</sup>	12.21 <sup>a</sup>	78.13 <sup>c</sup>	2.23 <sup>e</sup>	5.06 <sup>f</sup>	9.59 <sup>h</sup>	27.34 <sup>g</sup>	68.36 <sup>b</sup>
$T_{10}$	1.28 <sup>h</sup>	3.16 <sup>a</sup>	6.46 <sup>a</sup>	13.95 <sup>a</sup>	95.49 <sup>b</sup>	2.68 <sup>bc</sup>	5.76 <sup>d</sup>	13.89 <sup>d</sup>	39.06 <sup>e</sup>	NA
SE(d)	0.04	1.16	4.38	12.49	2.16	0.06	0.08	0.26	1.23	0.63
CD	0.08	2.42	9.13	26.06	4.51	0.12	0.17	0.54	2.56	1.30

**Treatment details:**  $T_1$  - Control (water dip);  $T_2$  - Melatonin (150  $\mu$ M);  $T_3$  - Melatonin (200  $\mu$ M);  $T_4$  - Melatonin (250  $\mu$ M);  $T_5$  - Salicylic acid (1.5 mM);  $T_6$  - Salicylic acid (2.0 mM);  $T_7$  - Salicylic acid (2.5 mM);  $T_8$  - Brassinosteroid (10 ppm);  $T_9$  - Brassinosteroid (25 ppm);  $T_{10}$  - Brassinosteroid (50 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).

**Table 6.** Effect of postharvest treatments on total sugar content (g/100g) in mango varieties under ambient storage

Treatments	Alphonso					Sindhura				
	Day 0	Day 3	Day 6	Day 9	Day 12	Day 0	Day 3	Day 6	Day 9	Day 12
<b>T<sub>1</sub></b>	4.85 <sup>a</sup>	6.29 <sup>a</sup>	8.83 <sup>a</sup>	14.17 <sup>a</sup>	20.05 <sup>a</sup>	4.25 <sup>abc</sup>	5.65 <sup>a</sup>	8.05 <sup>a</sup>	13.33 <sup>a</sup>	NA
<b>T<sub>2</sub></b>	4.78 <sup>ab</sup>	5.78 <sup>bc</sup>	8.42 <sup>b</sup>	11.21 <sup>d</sup>	18.92 <sup>b</sup>	4.43 <sup>a</sup>	5.42 <sup>b</sup>	7.39 <sup>bc</sup>	11.58 <sup>de</sup>	NA
<b>T<sub>3</sub></b>	4.45 <sup>c</sup>	5.54 <sup>d</sup>	6.83 <sup>d</sup>	10.12 <sup>e</sup>	16.22 <sup>g</sup>	4.10 <sup>cd</sup>	4.51 <sup>e</sup>	6.60 <sup>f</sup>	10.19 <sup>f</sup>	16.60 <sup>c</sup>
<b>T<sub>4</sub></b>	4.42 <sup>c</sup>	5.53 <sup>d</sup>	8.25 <sup>b</sup>	11.11 <sup>d</sup>	17.02 <sup>ef</sup>	4.21 <sup>bcd</sup>	5.07 <sup>c</sup>	7.16 <sup>cd</sup>	11.18 <sup>e</sup>	NA
<b>T<sub>5</sub></b>	4.60 <sup>bc</sup>	5.85 <sup>b</sup>	8.39 <sup>b</sup>	13.10 <sup>bc</sup>	18.38 <sup>bc</sup>	4.03 <sup>d</sup>	5.11 <sup>c</sup>	7.42 <sup>bc</sup>	11.86 <sup>cd</sup>	NA
<b>T<sub>6</sub></b>	4.76 <sup>ab</sup>	5.60 <sup>cd</sup>	7.46 <sup>c</sup>	10.48 <sup>e</sup>	16.64 <sup>fg</sup>	4.37 <sup>ab</sup>	5.02 <sup>c</sup>	6.78 <sup>ef</sup>	10.33 <sup>f</sup>	17.99 <sup>b</sup>
<b>T<sub>7</sub></b>	4.75 <sup>ab</sup>	5.83 <sup>b</sup>	8.26 <sup>b</sup>	13.46 <sup>b</sup>	18.05 <sup>cd</sup>	4.05 <sup>d</sup>	5.22 <sup>c</sup>	7.51 <sup>b</sup>	12.10 <sup>c</sup>	NA
<b>T<sub>8</sub></b>	4.74 <sup>ab</sup>	6.21 <sup>a</sup>	8.39 <sup>b</sup>	14.15 <sup>a</sup>	18.84 <sup>bc</sup>	4.42 <sup>a</sup>	5.49 <sup>ab</sup>	7.89 <sup>a</sup>	12.67 <sup>b</sup>	NA
<b>T<sub>9</sub></b>	4.44 <sup>c</sup>	5.41 <sup>d</sup>	7.69 <sup>c</sup>	9.25 <sup>f</sup>	16.10 <sup>g</sup>	4.12 <sup>cd</sup>	4.72 <sup>d</sup>	6.98 <sup>de</sup>	10.51 <sup>f</sup>	18.75 <sup>a</sup>
<b>T<sub>10</sub></b>	4.62 <sup>bc</sup>	5.60 <sup>cd</sup>	8.39 <sup>b</sup>	12.86 <sup>c</sup>	17.48 <sup>de</sup>	4.12 <sup>cd</sup>	5.48 <sup>ab</sup>	7.49 <sup>b</sup>	11.80 <sup>cd</sup>	NA
<b>SE(d)</b>	<b>0.11</b>	<b>0.10</b>	<b>0.16</b>	<b>0.25</b>	<b>0.38</b>	<b>0.09</b>	<b>0.10</b>	<b>0.15</b>	<b>0.23</b>	<b>0.18</b>
<b>CD</b>	<b>0.22</b>	<b>0.21</b>	<b>0.34</b>	<b>0.52</b>	<b>0.80</b>	<b>0.19</b>	<b>0.20</b>	<b>0.30</b>	<b>0.49</b>	<b>0.38</b>

**Treatment details:** T<sub>1</sub> - Control (water dip); T<sub>2</sub> - Melatonin (150 µM); T<sub>3</sub> - Melatonin (200 µM); T<sub>4</sub> - Melatonin (250 µM); T<sub>5</sub> - Salicylic acid (1.5 mM); T<sub>6</sub> - Salicylic acid (2.0 mM); T<sub>7</sub> - Salicylic acid (2.5 mM); T<sub>8</sub> - Brassinosteroid (10 ppm); T<sub>9</sub> - Brassinosteroid (25 ppm); T<sub>10</sub> - Brassinosteroid (50 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).

**Table 7.** Effect of postharvest treatments on total phenol content in mango varieties under ambient storage

Treatments	Alphonso					Sindhura				
	Day 0	Day 3	Day 6	Day 9	Day 12	Day 0	Day 3	Day 6	Day 9	Day 12
<b>T<sub>1</sub></b>	34.55 <sup>a</sup>	28.30 <sup>c</sup>	23.90 <sup>f</sup>	21.70 <sup>b</sup>	13.05 <sup>f</sup>	33.20 <sup>a</sup>	21.50 <sup>e</sup>	16.35 <sup>g</sup>	10.15 <sup>f</sup>	NA
<b>T<sub>2</sub></b>	34.70 <sup>a</sup>	29.05 <sup>bc</sup>	25.70 <sup>cd</sup>	22.00 <sup>b</sup>	18.45 <sup>b</sup>	33.55 <sup>a</sup>	24.50 <sup>cd</sup>	18.15 <sup>d</sup>	12.95 <sup>d</sup>	NA
<b>T<sub>3</sub></b>	35.45 <sup>a</sup>	30.45 <sup>a</sup>	27.55 <sup>ab</sup>	24.50 <sup>a</sup>	20.05 <sup>a</sup>	33.45 <sup>a</sup>	28.15 <sup>a</sup>	19.55 <sup>a</sup>	16.00 <sup>a</sup>	12.95 <sup>a</sup>
<b>T<sub>4</sub></b>	35.90 <sup>a</sup>	29.10 <sup>bc</sup>	26.85 <sup>bc</sup>	22.25 <sup>b</sup>	19.25 <sup>ab</sup>	32.65 <sup>a</sup>	23.70 <sup>d</sup>	17.85 <sup>de</sup>	11.20 <sup>e</sup>	NA
<b>T<sub>5</sub></b>	34.30 <sup>a</sup>	29.15 <sup>bc</sup>	25.30 <sup>de</sup>	20.10 <sup>c</sup>	16.55 <sup>c</sup>	33.00 <sup>a</sup>	24.45 <sup>d</sup>	18.85 <sup>bc</sup>	14.95 <sup>c</sup>	NA
<b>T<sub>6</sub></b>	35.30 <sup>a</sup>	29.85 <sup>ab</sup>	27.40 <sup>ab</sup>	24.00 <sup>a</sup>	18.45 <sup>b</sup>	32.80 <sup>a</sup>	26.10 <sup>b</sup>	19.30 <sup>ab</sup>	15.65 <sup>ab</sup>	12.40 <sup>b</sup>
<b>T<sub>7</sub></b>	35.50 <sup>a</sup>	23.15 <sup>e</sup>	25.75 <sup>cd</sup>	21.90 <sup>b</sup>	15.05 <sup>de</sup>	33.30 <sup>a</sup>	23.85 <sup>d</sup>	17.25 <sup>f</sup>	10.45 <sup>f</sup>	NA
<b>T<sub>8</sub></b>	35.05 <sup>a</sup>	25.45 <sup>d</sup>	24.15 <sup>ef</sup>	21.30 <sup>b</sup>	14.65 <sup>e</sup>	33.60 <sup>a</sup>	23.70 <sup>d</sup>	17.00 <sup>f</sup>	10.30 <sup>f</sup>	NA
<b>T<sub>9</sub></b>	34.95 <sup>a</sup>	29.15 <sup>bc</sup>	28.40 <sup>a</sup>	24.50 <sup>a</sup>	19.40 <sup>a</sup>	33.50 <sup>a</sup>	25.35 <sup>bc</sup>	18.75 <sup>c</sup>	15.45 <sup>abc</sup>	11.95 <sup>c</sup>
<b>T<sub>10</sub></b>	35.15 <sup>a</sup>	28.75 <sup>bc</sup>	25.05 <sup>def</sup>	22.00 <sup>b</sup>	15.75 <sup>cd</sup>	33.65 <sup>a</sup>	24.20 <sup>d</sup>	17.45 <sup>ef</sup>	15.10 <sup>bc</sup>	NA
<b>SE(d)</b>	<b>0.95</b>	<b>0.54</b>	<b>0.57</b>	<b>0.55</b>	<b>0.39</b>	<b>0.85</b>	<b>0.43</b>	<b>0.23</b>	<b>0.29</b>	<b>0.16</b>
<b>CD</b>	<b>1.98</b>	<b>1.13</b>	<b>1.18</b>	<b>1.15</b>	<b>0.82</b>	<b>1.77</b>	<b>0.90</b>	<b>0.49</b>	<b>0.61</b>	<b>0.33</b>

**Treatment details:** T<sub>1</sub> - Control (water dip); T<sub>2</sub> - Melatonin (150 µM); T<sub>3</sub> - Melatonin (200 µM); T<sub>4</sub> - Melatonin (250 µM); T<sub>5</sub> - Salicylic acid (1.5 mM); T<sub>6</sub> - Salicylic acid (2.0 mM); T<sub>7</sub> - Salicylic acid (2.5 mM); T<sub>8</sub> - Brassinosteroid (10 ppm); T<sub>9</sub> - Brassinosteroid (25 ppm); T<sub>10</sub> - Brassinosteroid (50 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).

**Table 8.** Effect of postharvest treatments on carotenoid content (µg/100g) in mango varieties under ambient storage

Treatments	Alphonso					Sindhura				
	Day0	Day 3	Day 6	Day 9	Day12	Day0	Day 3	Day 6	Day 9	Day12
<b>T<sub>1</sub></b>	4674.62 <sup>a</sup>	8578.9 <sup>a</sup>	13575.37 <sup>a</sup>	17671.91 <sup>a</sup>	23004.91 <sup>a</sup>	3425.30 <sup>a</sup>	7595.81 <sup>a</sup>	13533.72 <sup>a</sup>	18142.12 <sup>a</sup>	NA
<b>T<sub>2</sub></b>	4688.64 <sup>a</sup>	8138.12 <sup>bc</sup>	12137.32 <sup>cde</sup>	16780.31 <sup>bcd</sup>	22375.34 <sup>ab</sup>	3429.31 <sup>a</sup>	6555.19 <sup>c</sup>	12781.19 <sup>b</sup>	16706.73 <sup>b</sup>	NA
<b>T<sub>3</sub></b>	4686.83 <sup>a</sup>	7790.40 <sup>d</sup>	11615.73 <sup>e</sup>	16020.03 <sup>e</sup>	21133.43 <sup>c</sup>	3426.44 <sup>a</sup>	6057.02 <sup>d</sup>	11669.70 <sup>c</sup>	15990.03 <sup>c</sup>	19880.33 <sup>a</sup>
<b>T<sub>4</sub></b>	4686.36 <sup>a</sup>	8057.30 <sup>bcd</sup>	11803.40 <sup>de</sup>	16324.98 <sup>cde</sup>	22432.47 <sup>ab</sup>	3422.97 <sup>a</sup>	6188.53 <sup>d</sup>	11928.96 <sup>c</sup>	16352.16 <sup>bc</sup>	NA
<b>T<sub>5</sub></b>	4688.95 <sup>a</sup>	8117.86 <sup>bc</sup>	12517.01 <sup>bc</sup>	16970.80 <sup>bc</sup>	22436.25 <sup>ab</sup>	3422.76 <sup>a</sup>	6663.12 <sup>bc</sup>	12809.60 <sup>b</sup>	16722.68 <sup>b</sup>	NA
<b>T<sub>6</sub></b>	4687.27 <sup>a</sup>	7945.39 <sup>cd</sup>	11762.15 <sup>de</sup>	16185.76 <sup>de</sup>	21773.26 <sup>bc</sup>	3423.81 <sup>a</sup>	6164.52 <sup>d</sup>	11898.13 <sup>c</sup>	16272.69 <sup>bc</sup>	20468.76 <sup>a</sup>
<b>T<sub>7</sub></b>	4672.43 <sup>a</sup>	8072.05 <sup>bcd</sup>	12258.09 <sup>cd</sup>	16470.01 <sup>bcd</sup>	22233.27 <sup>ab</sup>	3425.20 <sup>a</sup>	6218.64 <sup>d</sup>	12149.47 <sup>c</sup>	16693.11 <sup>bc</sup>	NA
<b>T<sub>8</sub></b>	4688.81 <sup>a</sup>	8362.71 <sup>ab</sup>	12938.41 <sup>b</sup>	17047.00 <sup>ab</sup>	22984.11 <sup>a</sup>	3424.57 <sup>a</sup>	6953.95 <sup>b</sup>	12829.26 <sup>b</sup>	17766.35 <sup>a</sup>	NA
<b>T<sub>9</sub></b>	4687.30 <sup>a</sup>	7977.36 <sup>cd</sup>	11791.61 <sup>de</sup>	16072.06 <sup>e</sup>	21288.92 <sup>c</sup>	3427.91 <sup>a</sup>	6151.29 <sup>d</sup>	11885.42 <sup>c</sup>	16191.93 <sup>bc</sup>	20666.03 <sup>a</sup>
<b>T<sub>10</sub></b>	4683.28 <sup>a</sup>	8067.18 <sup>bcd</sup>	12683.63 <sup>bc</sup>	16446.96 <sup>bcd</sup>	22586.90 <sup>ab</sup>	3426.54 <sup>a</sup>	6207.52 <sup>d</sup>	12022.41 <sup>c</sup>	16418.94 <sup>bc</sup>	NA
<b>SE(d)</b>	<b>78.86</b>	<b>151.94</b>	<b>280.5015</b>	<b>333.11</b>	<b>423.44</b>	<b>62.77</b>	<b>140.95</b>	<b>232.11</b>	<b>341.76</b>	<b>495.16</b>
<b>CD</b>	<b>164.50</b>	<b>316.93</b>	<b>585.1158</b>	<b>694.85</b>	<b>883.28</b>	<b>130.93</b>	<b>294.01</b>	<b>484.18</b>	<b>712.90</b>	<b>1211.62</b>

**Treatment details:** T<sub>1</sub> - Control (water dip); T<sub>2</sub> - Melatonin (150 µM); T<sub>3</sub> - Melatonin (200 µM); T<sub>4</sub> - Melatonin (250 µM); T<sub>5</sub> - Salicylic acid (1.5 mM); T<sub>6</sub> - Salicylic acid (2.0 mM); T<sub>7</sub> - Salicylic acid (2.5 mM); T<sub>8</sub> - Brassinosteroid (10 ppm); T<sub>9</sub> - Brassinosteroid (25 ppm); T<sub>10</sub> - Brassinosteroid (50 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).


## Enzyme Activity

### Ripening enzymes


**Pectin methyl esterase activity:** Pectin methyl esterase (PME) activity in mangoes increased as the fruits ripened, as represented in Fig. 1. The highest PME activity was observed on the 12<sup>th</sup> day for Alphonso (466.00 m.eq. COOH/min/g) and on the 9<sup>th</sup> day for Sindhura (107.00 m.eq. COOH/min/g). All treatments significantly reduced PME activity compared to the control, with 200  $\mu$ M Melatonin ( $T_3$ ), 2.0 mM Salicylic Acid ( $T_6$ ) and 25 ppm Brassinosteroids ( $T_9$ ) showing the lowest levels by the end of the storage period. Specifically, PME activities for Alphonso were 409.00, 412.00 and 417.00 m.eq. COOH/min/g, while for Sindhura, they were 94.00, 96.00 and 94.00 m.eq. COOH/min/g on the 12<sup>th</sup> and 9<sup>th</sup> days, respectively.

**Polygalacturonase activity:** Polygalacturonase (PG) activity in the mango varieties Alphonso and Sindhura increased with storage duration. The highest enzyme activity was recorded in the control fruits on the last day of storage, at 79.43  $\mu$ g/min/g for Alphonso and 77.24  $\mu$ g/min/g for Sindhura. In contrast, the lowest PG activity for Alphonso was found in  $T_9$  at 65.91  $\mu$ g/min/g, followed by  $T_3$  at 66.15  $\mu$ g/min/g and  $T_8$  at 66.27  $\mu$ g/min/g. For Sindhura, the lowest values were similar across treatments, with  $T_3$  at 68.34  $\mu$ g/min/g,  $T_6$  at 68.83  $\mu$ g/min/g and  $T_9$  at 69.27  $\mu$ g/min/g (as shown in Fig. 2.).


**Amylase activity:** Amylase activity, as shown in Fig. 3, increased significantly during ripening before declining toward the end of storage in both mango varieties. For Alphonso, peak activity occurred on the 9<sup>th</sup> day in the control group (45.35  $\mu$ g/min/g), while the lowest activity was found in  $T_3$  (33.79  $\mu$ g/min/g),  $T_6$



**Fig. 1.** Effect of bioactive compounds on PME activity (%) in mango varieties A) Alphonso and B) Sindhura under ambient storage.  $T_1$ - Control (water dip);  $T_3$ - Melatonin (200  $\mu$ M);  $T_6$ - Salicylic acid (2.0 mM);  $T_9$ - Brassinosteroid (25 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).



**Fig. 2.** Effect of bioactive compounds on polygalacturonase activity ( $\mu$ g/min/g) in mango varieties A) Alphonso and B) Sindhura under ambient storage.  $T_1$ - Control (water dip);  $T_3$ - Melatonin (200  $\mu$ M);  $T_6$ - Salicylic acid (2.0 mM);  $T_9$ - Brassinosteroid (25 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).



**Fig. 3.** Effect of bioactive compounds on amylase activity ( $\mu\text{g}/\text{min}/\text{g}$ ) in mango varieties A) Alphonso and B) Sindhura under ambient storage. T<sub>1</sub>- Control (water dip); T<sub>3</sub>- Melatonin (200  $\mu\text{M}$ ); T<sub>6</sub>- Salicylic acid (2.0 mM); T<sub>9</sub>- Brassinosteroid (25 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).


(34.03  $\mu\text{g}/\text{min}/\text{g}$ ), and T<sub>9</sub> (35.50  $\mu\text{g}/\text{min}/\text{g}$ ) on the same day. Enzyme activity declined by the final storage day. In Sindhura, peak activity was noted on the 6<sup>th</sup> day for the control fruits, with treated groups experiencing a delay of nearly three days. The control recorded a peak of 47.25  $\mu\text{g}/\text{min}/\text{g}$ , while the lowest activity among treatments was in T<sub>3</sub> (34.03  $\mu\text{g}/\text{min}/\text{g}$ ), followed by T<sub>10</sub> (35.56  $\mu\text{g}/\text{min}/\text{g}$ ) and T<sub>6</sub> and T<sub>9</sub>, which were comparable at 36.15  $\mu\text{g}/\text{min}/\text{g}$  and 36.68  $\mu\text{g}/\text{min}/\text{g}$ , respectively.

**Cellulase activity:** Cellulase activity for Alphonso and Sindhura is presented in Fig. 4. In Alphonso, enzyme activity increased steadily, peaking on the last day of storage at 101.20  $\mu\text{g}/\text{min}/\text{g}$  in the control group. In contrast, Sindhura's peak activity in treatments T<sub>3</sub>, T<sub>6</sub> and T<sub>9</sub> was delayed by three days, while other treatments reached their peak on the 9<sup>th</sup> day. The lowest

cellulase activities for Alphonso were noted in T<sub>3</sub> (72.39  $\mu\text{g}/\text{min}/\text{g}$ ), T<sub>9</sub> (75.23  $\mu\text{g}/\text{min}/\text{g}$ ) and T<sub>6</sub> (75.37  $\mu\text{g}/\text{min}/\text{g}$ ). For Sindhura, the control recorded a peak of 81.90  $\mu\text{g}/\text{min}/\text{g}$ , with the lowest values seen in T<sub>3</sub> and T<sub>6</sub> on the 9<sup>th</sup> day.


#### Antioxidant enzymes

**Peroxidase activity:** Fig. 5 shows a gradual increase in peroxidase activity during storage for both mango varieties, peaking before declining at the end. Alphonso reached its peak on the 9<sup>th</sup> day, while Sindhura peaked on the 6<sup>th</sup> day. The control group displayed significantly lower peroxidase activity at 0.30  $\Delta\text{A}/\text{min}/\text{g}$  on their respective peak days compared to the treated fruits. Notably, the highest peroxidase activity was found in fruits treated with 200  $\mu\text{M}$  Melatonin (T<sub>3</sub>), 2.0 mM Salicylic Acid (T<sub>6</sub>) and 25 ppm Brassinosteroid (T<sub>9</sub>).



**Fig. 4.** Effect of bioactive compounds on cellulase activity ( $\mu\text{g}/\text{min}/\text{g}$ ) in mango varieties A) Alphonso and B) Sindhura under ambient storage. T<sub>1</sub>- Control (water dip); T<sub>3</sub>- Melatonin (200  $\mu\text{M}$ ); T<sub>6</sub>- Salicylic acid (2.0 mM); T<sub>9</sub>- Brassinosteroid (25 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).







**Fig. 5.** Effect of bioactive compounds on peroxidase activity ( $\Delta A/\text{min/g}$ ) in mango varieties A) Alphonso and B) Sindhura under ambient storage. T<sub>1</sub>- Control (water dip); T<sub>3</sub>- Melatonin (200  $\mu\text{M}$ ); T<sub>6</sub>- Salicylic acid (2.0 mM); T<sub>9</sub>- Brassinosteroid (25 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).

**Catalase activity:** Fig. 6 presents the changes in catalase activity during storage for Alphonso and Sindhura mangoes. Catalase activity initially increased, peaking on the 9<sup>th</sup> day before declining through the 12<sup>th</sup> day across all treatments. Only three treatments, 200  $\mu\text{M}$  Melatonin (T<sub>3</sub>), 2.0 mM Salicylic Acid (T<sub>6</sub>) and 25 ppm Brassinosteroid (T<sub>9</sub>), remained effective in Sindhura until day 12. These treatments exhibited significantly higher catalase activity, with values of 4.87, 3.80 and 4.39 activity/min/g in Alphonso and 5.96, 5.79 and 5.92 activity/min/g in Sindhura. In contrast, the control group displayed lower levels of catalase activity at 2.68 and 4.34 activity/min/g for Alphonso and Sindhura, respectively.

**Superoxide dismutase activity:** Fig. 7 shows the superoxide dismutase (SOD) activity for Alphonso and Sindhura mangoes. SOD activity steadily increased throughout the storage period, peaking on the last day. Control fruits had lower SOD activity, measuring 69.85 units/min/g for Alphonso and 58.49 units/min/g for Sindhura. In contrast, fruits treated with 200  $\mu\text{M}$  melatonin (T<sub>3</sub>), 25 ppm brassinosteroid (T<sub>9</sub>) and 2.0 mM salicylic acid (T<sub>6</sub>) exhibited significantly higher SOD activity. In Alphonso, the recorded values on the 12<sup>th</sup> day were 81.56, 81.02 and 79.36 units/min/g, respectively, while in Sindhura, the corresponding values on the 9<sup>th</sup> day were 69.03, 67.58 and 67.02 units/min/g.



**Fig. 6.** Effect of bioactive compounds on catalase activity (activity/min/g) in mango varieties A) Alphonso and B) Sindhura under ambient storage. T<sub>1</sub>- Control (water dip); T<sub>3</sub>- Melatonin (200  $\mu\text{M}$ ); T<sub>6</sub>- Salicylic acid (2.0 mM); T<sub>9</sub>- Brassinosteroid (25 ppm). Each value is a mean for three replicates ( $p = 0.05$ ).



**Fig. 7. Effect of bioactive compounds on superoxide dismutase activity (units/min/g) in mango varieties A) Alphonso and B) Sindhura under ambient storage.** T<sub>1</sub>- Control (water dip); T<sub>3</sub>- Melatonin (200 µM); T<sub>6</sub>- Salicylic acid (2.0 mM); T<sub>9</sub>- Brassinosteroid (25 ppm). Each value is a mean for three replicates ( $p=0.05$ ).

## Discussion


The increase in shelf life and quality of Alphonso and Sindhura mangoes treated with melatonin, salicylic acid and brassinosteroids can be attributed to several key factors related to the modulation of ripening, enzymatic activity and antioxidant capacity. Melatonin (200 µM), salicylic acid (2.0 mM) and brassinosteroids (25 ppm) demonstrated the ability to delay fruit ripening by reducing ethylene biosynthesis and respiration rates (23). Ethylene accelerates ripening and triggers various ripening-related processes such as enzymatic activities, softening and colour change (Fig. 8). By lowering ethylene levels, these treatments were found to slow down these processes, extending the shelf life of fruit. Exogenous application of melatonin, salicylic acid and brassinosteroids has been reported to increase storability in fruits (9, 10, 24) by regulating quality parameters and ethylene production.

The physiological loss in weight (PLW) was significantly lower in treated fruits, which retained firmness over a long duration. These findings align with those of Ali et al. (25), Kaur et al. (26) and others (27). These results are likely due to the reduced activity of ripening enzymes in treated fruits. Enzymes such as pectin methylesterase (PME), polygalacturonase (PG), cellulase (CEL) and amylase are involved in the breakdown of cell wall components, leading to fruit softening and textural changes (28). In treated fruits, these enzymes exhibited lower activity, especially in the T<sub>3</sub> (Melatonin), T<sub>6</sub> (Salicylic acid) and T<sub>9</sub> (Brassinosteroid) treatments. This lower activity reduces the degradation of pectin and cellulose, thereby maintaining fruit firmness and weight loss for a longer period (Fig. 8). This is likely due to the ability of melatonin, salicylic acid and brassinosteroids to reduce water loss and slow down the physiological processes that lead to fruit dehydration (29). Reduced PLW also means less susceptibility to shrivelling and spoilage.

The interplay between TSS, acidity and total sugars is critical for the maintenance of quality and shelf life. A balanced increase in TSS during storage indicates ripening and controlled TSS levels help maintain fruit quality over extended periods (30). Low sugar levels slow the conversion of starches to sugars,

reducing the metabolic activity associated with ripening. It reduces the activity of cell wall-degrading enzymes (31). While high acidity helps maintain firmness, preventing excessive softening of the fruit over time and it balances flavour during storage (2). It suppresses the growth of spoilage-causing microorganisms, effectively extending its shelf life.

Higher total phenol content contributes significantly to shelf life by delaying deterioration, maintaining quality, and reducing vulnerability to spoilage. Phenols neutralize free radicals, reducing oxidative stress in fruit tissues. This preserves cellular integrity, slowing down ripening and senescence. Additionally, phenols inhibit enzymes like polygalacturonase and cellulase, which degrade cell walls, thereby reducing fruit softening and maintaining texture (32). Phenols also exhibit antimicrobial properties that inhibit spoilage causing microorganisms. High carotenoid content further enhances mangoes shelf life by boosting antioxidant capacity and improving fruit quality. As potent antioxidants, carotenoids help neutralize free radicals, reducing oxidative stress and delaying senescence in fruit tissues. It also provides a vibrant yellow to orange colour to the fruits, which enhances marketability (33).



**Fig. 8. Probable mode of action of bioactive compounds in postharvest fruits.**

Reduced activity of cell wall degradation enzymes might be due to the lower ethylene levels and respiration rates, as ethylene acts as a cascade and is important for the activation of genes responsible for the synthesis of key enzymes (34). PME hydrolyses pectin, a major component of the cell wall, causing fruit softening. The activation of PG is a direct response to increased ethylene levels. Cellulase and hemicellulase break down cellulose and hemicellulose in the cell wall, while PME modifies pectin by demethylating it, making it more susceptible to degradation by PG. Ethylene regulates the activity of PG, PME and cellulase (34). In our study on melatonin, salicylic acid and brassinosteroid application the activity of these enzymes was found to be reduced. The results are in line with previous studies of Wang et al. (35), Amiri et al. (36) and Meena et al. (37).

Fruit ripening is a complex oxidative process involving reactive oxygen species (ROS), which can accelerate fruit deterioration if not controlled. Ethylene production during ripening elevates respiration, increasing ROS levels. Antioxidant enzymes like superoxide dismutase (SOD), catalase (CAT) and peroxidase (POD) play crucial roles in scavenging ROS and minimizing oxidative damage (38). Melatonin, salicylic acid (SA) and brassinosteroids (BRs) have been shown to enhance the activity of these enzymes, reducing oxidative stress and extending fruit shelf life. Melatonin boosts the expression of ROS-related genes, increasing the activity of SOD, CAT and ascorbate peroxidase (APX), thereby lowering lipid peroxidation and preserving fruit quality (39). Similarly, SA and BRs regulate antioxidant defences by increasing the levels of SOD, CAT and PAL (phenylalanine ammonia-lyase), reducing the accumulation of ROS and oxidative stress markers like malondialdehyde (MDA) and hydrogen peroxide ( $H_2O_2$ ) (7, 40). Collectively, these treatments delay ripening, maintain fruit quality and extend shelf life by bolstering the fruit's natural antioxidant defence systems.

The use of bioactive compounds like melatonin, salicylic acid and brassinosteroids in post-harvest management offers significant environmental and economic benefits. These natural compounds provide a sustainable alternative to synthetic chemicals, reducing pollution, soil degradation and chemical residues on fruits while supporting environmental conservation. Economically, they minimize post-harvest losses, extend shelf life, and maintain fruit quality, enabling access to distant markets and reducing costs associated with refrigeration and chemical inputs. By enhancing fruit appearance, taste and nutritional value, these treatments meet consumer demands for premium-quality produce, increasing profitability while promoting sustainable and eco-friendly agricultural practices (41, 42).

The combined effects of melatonin, salicylic acid and brassinosteroids in regulating ripening enzymes, enhancing antioxidant defences, maintaining firmness and controlling chemical changes are key reasons for the observed increase in shelf life and improved quality in mangoes. These treatments effectively delay the ripening process, slow down enzymatic activities related to softening and protect the fruit from oxidative damage, thereby extending its marketability and storage potential.

Future research should focus on optimizing the dosage, timing and application methods of melatonin, salicylic acid and brassinosteroids for different mango varieties and storage conditions. Integrating these treatments with existing post-

harvest technologies such as cold storage and modified atmosphere packaging, could enhance scalability. Additionally, developing cost-effective formulations and evaluating their feasibility for large-scale adoption will be critical for bridging the gap between laboratory research and commercial application in post-harvest management systems.

## Conclusion

The present study demonstrates that, among the various treatments tested at different concentrations, melatonin (200  $\mu$ M), salicylic acid (2.0 mM) and brassinosteroids (25 ppm) proved to be the most effective in extending the shelf life and improving the quality of Alphonso and Sindhura mango varieties. These treatments successfully delayed the ripening process by regulating key ripening enzymes, such as pectin methylesterase, polygalacturonase, cellulase and amylase, which are responsible for softening and starch degradation. Additionally, they enhanced the activity of antioxidant enzymes, including superoxide dismutase, catalase and peroxidase, thereby reducing oxidative damage during storage. This resulted in better retention of fruit firmness, reduced weight loss and favorable chemical changes like reduced TSS and sugar content and high acidity. Moreover, these treatments maintained non-enzymatic antioxidants, such as phenols and carotenoids. Overall, melatonin (200  $\mu$ M), salicylic acid (2.0 mM) and brassinosteroids (25 ppm) provided superior post-harvest benefits, making them the best treatments for maximizing mango shelf life and quality under ambient storage conditions. The beneficial effects of these bioactive compounds demonstrate significant potential for scalability in commercial mango production. The use of these natural bioactive compounds offers an eco-friendly and cost-effective solution suitable for integration into existing post-harvest management practices. With minimal infrastructure requirements and broad applicability, these treatments could be adopted by both small-scale and large-scale producers, enhancing the marketability and export potential of mangoes while reducing post-harvest losses on a commercial scale.

## Acknowledgements

The authors are thankful to the Tamil Nadu Agricultural University for supporting their research work.

## Authors' contributions

AB planned the formal analysis and carried out the investigation, formulated and adopted the standard methodology, software and wrote the original draft. SV contributed to the conceptualisation, framing the methodology, obtaining resources, funding acquisition, supervision and writing, reviewing and editing the manuscript. KC helped in framing the methodology, employing software and writing, reviewing and editing the manuscript. VAG contributed to writing, reviewing and editing the manuscript. RM also participated in writing, reviewing and editing the manuscript. All authors have read and agreed to the published version of the manuscript.

## Compliance with ethical standards

**Conflict of interest:** The authors declare no conflicts of interest.

**Ethical issues:** None

## References

- Tharanathan R, Yashoda H, Prabha T. Mango (*Mangifera indica* L.), "The king of fruits"—An overview. *Food Reviews International*. 2006;22(2):95-123. <https://doi.org/10.1080/87559120600574493>
- Bhardwaj R, Pareek S, Mani S, Domínguez-Avila JA, González-Aguilar GA. A melatonin treatment delays postharvest senescence, maintains quality, reduces chilling injury and regulates antioxidant metabolism in mango fruit. *Journal of Food Quality*. 2022;2022(1):2379556. <https://doi.org/10.1155/2022/2379556>
- Kebbeh M, Dong J-X, Chen H, Yan L, Zheng X-L. Melatonin treatment alleviates chilling injury in mango fruit 'Keitt' by modulating proline metabolism under chilling stress. *Journal of Integrative Agriculture*. 2023;22(3):935-44. <https://doi.org/10.1016/j.jia.2023.02.008>
- Cortés-Montaña D, Bernalte-García MJ, Velardo-Micharet B, Serrano M, Serradilla MJ. Impact of pre-storage melatonin application on the standard, sensory and bioactive quality of early sweet cherry. *Foods*. 2023;12(8):1723. <https://doi.org/10.3390/foods12081723>
- Zhang W, Jiang Y, Zhang Z. The role of different natural organic acids in postharvest fruit quality management and its mechanism. *Food Frontiers*. 2023;4(3):1127-43. <https://doi.org/10.1002/fft2.245>
- Zhu Z, Zhang Z, Qin G, Tian S. Effects of brassinosteroids on postharvest disease and senescence of jujube fruit in storage. *Postharvest Biology and Technology*. 2010;56(1):50-55. <https://doi.org/10.1016/j.postharvbio.2009.11.014>
- Meng X, Fang J, Fu M, Jiao W, Ren P, Yang X. The role of 1-methylcyclopropylene (1-MCP) and salicylic acid (SA) in induced resistance of postharvest fruits. *Horticulturae*. 2023;9(1):108. <https://doi.org/10.3390/horticulturae9010108>
- Jayarajan S, Sharma R. Melatonin: A blooming biomolecule for postharvest management of perishable fruits and vegetables. *Trends in Food Science and Technology*. 2021;116:318-28. <https://doi.org/10.1016/j.tifs.2021.07.034>
- Chen C, Sun C, Wang Y, Gong H, Zhang A, Yang Y, et al. The preharvest and postharvest application of salicylic acid and its derivatives on storage of fruit and vegetables: A review. *Scientia Horticulturae*. 2023;312:111858. <https://doi.org/10.1016/j.scienta.2023.111858>
- Ali S, Anjum MA, Nawaz A, Naz S, Hussain S, Ejaz S. Effects of brassinosteroids on postharvest physiology of horticultural crops: a concise review. *J Hortic Sci Technol*. 2019;2(3):62-68. <https://doi.org/10.46653/jhst190203062>
- Ranganna S. Manual of analysis of fruit and vegetable products. New Delhi: McGraw-Hill; 1977.
- Hedge J, Hofreiter B, Whistler R. Carbohydrate chemistry. Academic Press, New York. 1962;17:371-80.
- Bray H, Thorpe W. Analysis of phenolic compounds of interest in metabolism. *Methods of Biochemical Analysis*. 1954;27-52. <https://doi.org/10.1002/9780470110171.ch2>
- Roy SK. Simple and rapid method for estimation of total carotenoid pigments in mango. 1973;45.
- Srivastava MK, Dwivedi UN. Delayed ripening of banana fruit by salicylic acid. *Plant Science*. 2000;158(1-2):87-96. [https://doi.org/10.1016/S0168-9452\(00\)00304-6](https://doi.org/10.1016/S0168-9452(00)00304-6)
- Hagerman AE, Austin PJ. Continuous spectrophotometric assay for plant pectin methyl esterase. *Journal of Agricultural and Food Chemistry*. 1986;34(3):440-44. <https://doi.org/10.1021/jf00069a015>
- Gayathri T, Nair AS. Biochemical analysis and activity profiling of fruit ripening enzymes in banana cultivars from Kerala. *Journal of Food Measurement and Characterization*. 2017;11:1274-83. <https://doi.org/10.1007/s11694-017-9505-6>
- Bernfeld P. Amylases,  $\alpha$  and  $\beta$ . *Methods in Enzymology*. 1955;1:149-58. [https://doi.org/10.1016/0076-6879\(55\)01021-5](https://doi.org/10.1016/0076-6879(55)01021-5)
- Li R, Wang Y, Li W, Shao Y. Comparative analyses of ripening, texture properties and cell wall composition in three tropical fruits treated with 1-methylcyclopropene during cold storage. *Horticulturae*. 2023;9(2):126. <https://doi.org/10.3390/horticulturae9020126>
- Pütter J. Peroxidases. Elsevier. 1974;685-90. <https://doi.org/10.1016/B978-0-12-091302-2.50033-5>
- Aebi H. [13] Catalase *in vitro*. Elsevier. 1984;121-26. [https://doi.org/10.1016/S0076-6879\(84\)05016-3](https://doi.org/10.1016/S0076-6879(84)05016-3)
- Kakkar P, Das B, Viswanathan PN. A modified spectrophotometric assay of superoxide dismutase. *Indian Journal of Biochemistry and Biophysics*. 1984;21(2):130-32.
- Ma Q, Lin X, Wei Q, Yang X, Zhang Yn, Chen J. Melatonin treatment delays postharvest senescence and maintains the organoleptic quality of 'Newhall' navel orange (*Citrus sinensis* (L.) Osbeck) by inhibiting respiration and enhancing antioxidant capacity. *Scientia Horticulturae*. 2021;286:110236. <https://doi.org/10.1016/j.scienta.2021.110236>
- Ze Y, Gao H, Li T, Yang B, Jiang Y. Insights into the roles of melatonin in maintaining quality and extending shelf life of postharvest fruits. *Trends in Food Science and Technology*. 2021;109:569-78. <https://doi.org/10.1016/j.tifs.2021.01.051>
- Ali S, Nawaz A, Naz S, Ali M, Ejaz S, Azam M, et al. Exogenous melatonin mitigates chilling injury in zucchini fruit by enhancing antioxidant system activity, promoting endogenous proline and GABA accumulation and preserving cell wall stability. *Postharvest Biology and Technology*. 2023;204:112445. <https://doi.org/10.1016/j.postharvbio.2023.112445>
- Kaur P, Brar J, Kaur G. Salicylic acid and ascorbic acid extend the shelf life of guava (*Psidium guajava* L.) fruits under cold and ambient storage conditions. *Agricultural Research Journal*. 2022;59(4). <https://doi.org/10.5958/2395-146x.2022.00102.8>
- Gutiérrez-Villamil DA, Balaguera-López HE, Álvarez-Herrera JG. Brassinosteroids improve postharvest quality, antioxidant compounds and reduce chilling injury in 'Arrayana' mandarin fruits under cold storage. *Horticulturae*. 2023;9(6):622. <https://doi.org/10.3390/horticulturae9060622>
- Batool M, Bashir O, Amin T, Wani SM, Masoodi F, Jan N, et al. Effect of oxalic acid and salicylic acid treatments on the post-harvest life of temperate grown apricot varieties (*Prunus armeniaca*) during controlled atmosphere storage. *Food Science and Technology International*. 2022;28(7):557-69. <https://doi.org/10.1177/10820132211032074>
- Sun Q, Zhang N, Wang J, Zhang H, Li D, Shi J, et al. Melatonin promotes ripening and improves quality of tomato fruit during postharvest life. *Journal of Experimental Botany*. 2015;66(3):657-68. <https://doi.org/10.1093/jxb/eru332>
- Makroo H, Prabhakar PK, Rastogi N, Srivastava B. Characterization of mango puree based on total soluble solids and acid content: Effect on physico-chemical, rheological, thermal and ohmic heating behavior. *Lwt*. 2019;103:316-24. <https://doi.org/10.1016/j.lwt.2019.01.003>
- Ali A, Rasool K, Ganai N, Lone I, Parveen S, Anayat R, et al. Postharvest salicylic acid application affects the biochemical quality and stability of strawberry (*Fragaria x ananassa* Duch.) cv. Chandler fruits under ambient storage conditions. *Journal of Experimental Agriculture International*. 2022;44(9):195-202. <https://doi.org/10.9734/jeai/2022/v44i930866>
- Liu X, Wei L, Miao C, Zhang Q, Yan J, Li S, et al. Application of exogenous phenolic compounds in improving postharvest fruits quality: Classification, potential biochemical mechanisms and synergistic treatment. *Food Reviews International*. 2024;40(6):1776-95. <https://doi.org/10.1080/87559129.2023.2233599>



33. Lenucci MS, Tornese R, Mita G, Durante M. Bioactive compounds and antioxidant activities in different fractions of mango fruits (*Mangifera indica* L., cultivar Tommy Atkins and Keitt). *Antioxidants*. 2022;11(3):484. <https://doi.org/10.3390/antiox11030484>
34. Lohani S, Trivedi PK, Nath P. Changes in activities of cell wall hydrolases during ethylene-induced ripening in banana: effect of 1-MCP, ABA and IAA. *Postharvest Biology and Technology*. 2004;31(2):119-26. <https://doi.org/10.1016/j.postharvbio.2003.08.001>
35. Wang J, Hong M, Feng Y, He M-y, Wang R-k, Yu Z-x, et al. Effects of exogenous melatonin treatment on fruit quality and ethanol metabolism of 'Aiyuan 38' citrus. 2019;147-54. <https://doi.org/10.13995/j.cnki.11-1802/ts.021282>
36. Amiri S, Nicknam Z, Radi M, Sayadi M, Bagheri F, Karimi Khorrami N, et al. Postharvest quality of orange fruit as influenced by salicylic acid, acetic acid and carboxymethyl cellulose coating. *Journal of Food Measurement and Characterization*. 2021;15(5):3912-30. <https://doi.org/10.1007/s11694-021-00966-y>
37. Meena NK, Asrey R, Singh S, Singh D, Paul V. Exogenous application of brassinolide affects ripening, phenotypic traits, jelly seed and sensory properties of 'Dashehari' mango at ambient storage. *Annals of Plant and Soil Research*. 2023;25(2):205-10. <https://doi.org/10.47815/apsr.2023.10258>
38. Meitha K, Pramesti Y, Suhandono S. Reactive oxygen species and antioxidants in postharvest vegetables and fruits. *International Journal of Food Science*. 2020;2020(1):8817778. <https://doi.org/10.1155/2020/8817778>
39. Yun Z, Gao H, Chen X, Chen Z, Zhang Z, Li T, et al. Effects of hydrogen water treatment on antioxidant system of litchi fruit during the pericarp browning. *Food Chemistry*. 2021;336:127618. <https://doi.org/10.1016/j.foodchem.2020.127618>
40. Rashidi H, Amiri J, Shirzad H. Effect of postharvest treatment with 24-epibrassinolide and fennel (*Foeniculum vulgare*) essential oil on quality attributes and storage life of orange (*Citrus sinensis* Cv. 'Valencia'). *Erwerbs-Obstbau*. 2023;65(4):927-39. <https://doi.org/10.1007/s10341-022-00791-7>
41. Liu Y, Xu J, Lu X, Huang M, Yu W, Li C. The role of melatonin in delaying senescence and maintaining quality in postharvest horticultural products. *Plant Biology*. 2024. <https://doi.org/10.1111/plb.13706>
42. Mandal D, Ennio N, Lalhrualtuangi N, Fanai AV. Response of melatonin on postharvest qualities and shelf life of pineapple cv. Kew at ambient storage. *Journal of Applied and Natural Science*. 2024;16(2):794-804. <https://doi.org/10.31018/jans.v16i2.5562>