



RESEARCH ARTICLE

# Climatic and edaphic influences on productivity and carbon sequestration of farm grown teak (*Tectona grandis*, linn.f) in Tamil Nadu, India

S Navaneetha Krishnan<sup>1</sup>, A Balasubramanian<sup>1\*</sup>, M Sivaprakash<sup>1</sup>, R Ravi<sup>2</sup>, B Sivakumar<sup>1</sup>, C N Hari Prasath<sup>1</sup>, G Swathiga<sup>1</sup>, V Manimaran<sup>1</sup> & K S Anjali<sup>1</sup>

<sup>1</sup>Department of Silviculture and NRM, Forest College and Research Institute, Tamil Nadu Agricultural University, Mettupalayam, Coimbatore 641 301, Tamil Nadu, India

<sup>2</sup>Department of Forest Products and Wildlife, Forest College and Research Institute, Tamil Nadu Agricultural University, Mettupalayam, Coimbatore 641 301, Tamil Nadu, India

\*Email: [balasubramanian.a@tnau.ac.in](mailto:balasubramanian.a@tnau.ac.in)



## ARTICLE HISTORY

Received: 23 October 2024  
Accepted: 03 December 2024

Available online  
Version 1.0 : 28 December 2024



## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Krishnan NS, Balasubramanian A, Sivaprakash M, Ravi R, Sivakumar B, CN Hari Prasath CNH, Swathiga G, Manimaran V, Anjali KS. Climatic and edaphic influences on productivity and carbon sequestration of farm grown teak (*Tectona grandis*, linn.f) in Tamil Nadu, India. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.6090>

## Abstract

Teak (*Tectona grandis*, Linn.f) is one of the most sought-after tree crops due to the exceptional quality and high market value of its timber. Currently, the Indian subcontinent imports a significant quantity of teak to meet its growing demand. The growth and wood quality of teak are influenced by edaphic and climatic variations across sites, a factor that has been largely underexplored in research. The primary objective of the study is to analyse teak growth, wood quality, and carbon sequestration potential under farmland condition. To address this, the study was conducted across three agroclimatic zones of Tamil Nadu: North Eastern Zone (NEZ), North Western Zone (NWZ), and Cauvery Delta Zone (CDZ), focusing specifically on farmland conditions. The analysis encompassed biometric attributes, volume, and carbon sequestration potential of farm-grown teak across different age classes. The findings of the study reveal that among the three agroclimatic zones, the NEZ (15-20 years age class) recorded the highest biometric attributes, including mid diameter (MD) (0.193 m), volume (0.379 m<sup>3</sup>), heartwood proportion (0.239 m<sup>3</sup>), and carbon sequestration (0.683 Mg/acre).. Based on the overall performance, trees in the 15-20 years age class exhibited superior heartwood formation under farmland conditions. Therefore, this age class is recommended for teak cultivation in farmland conditions across Tamil Nadu to maximize growth, wood quality, and carbon sequestration potential.

## Keywords

agroclimatic zones; carbon sequestration; farmland; heartwood volume; teak; wood quality

## Introduction

Teak (*Tectona grandis* Linn.f) is one of the most predominant and prominent hardwood tree species which grows extensively in tropical conditions (1–3). Its natural growing range is predominantly concentrated in South and Southeast Asia, which together contribute more than 90 percent of the world's teak resources (4, 5). India accounts for approximately 1.68 M ha of teak plantations and 6.8 M ha of natural teak forests (6), contributing around 45% of global teak resources. Paradoxically, despite this significant share, India remains a net importer of teak, with an annual domestic production of only 0.25 M cu.m (7) compared to an annual demand of 100 M cu.m (8–10). Globally, India ranks first in teak consumption, meeting nearly

70 to 100 percent of its requirement through imports, predominantly from Africa and Latin America (11).

Given this significant demand supply gap, the promotion of teak cultivation outside forested areas, particularly on private patta lands, has gained considerable importance. Such initiatives aim to bridge the gap between teak wood demand and supply while also providing a viable revenue source for farmers.

Farm-grown teak in Tamil Nadu exhibits notable growth variations across different agroclimatic zones (12). These variations are influenced by the climatic and edaphic factors unique to each zone, resulting in significant differences in volume production (13). Additionally, long-rotation species such as teak possess the unique advantage of long-term carbon sequestration, as the wood is commonly used for furniture and construction, thereby locking carbon either for extended periods or permanently (14, 15).

However, there is a lack of systematic studies to understand the growth biometry, yield variations, and carbon sequestration potential of farm-grown teak across different agroclimatic zones. This study aims to fill this gap by providing scientific insights into variations in growth patterns, including parameters such as height, merchantable volume, girth, heart content, sap-to-heartwood ratio, and carbon sequestration potential of teak cultivated on farmland across Tamil Nadu's agroclimatic zones. The findings will enable a comprehensive evaluation of teak productivity and quality, ultimately aiding in the substantial promotion of teak cultivation in the state.

## Materials and Methods

### Study area

Farm-grown teak plantations were surveyed across three agroclimatic zones of Tamil Nadu, namely the NEZ, NWZ, and the CDZ, as detailed in Table 1.

Climatic data for these three agroclimatic zones over the past ten years were collected using NASA's Mod-

**Table 1.** Study zones and district wise teak resources

	Zone	District covered
Zone 1	North eastern zone	Thiruvannamalai, Vellore, Ranipet,
Zone 2	North western zone	Kallakurichi, Krishnagiri, Salem,
Zone 3	Cauvery delta zone (CDZ)	Trichy, Thanjavur, Perambalur

**NEZ** - North Eastern Zone, **NWZ** - North Western Zone, **CDZ** - Cauvery Delta Zone.

**Table 2.** Climatic attributes of three agroclimatic zones

Zones	Mean Maximum Temperature (°C)	Mean Minimum Temperature (°C)	Annual rainfall (mm)	Relative Humidity (%)	Elevation (m)
NEZ	40.43	13.83	1109	68.92	666
NWZ	41.53	13.12	849	68.67	453
CDZ	38.49	19.38	956.3	71.66	629

**NEZ** - North Eastern Zone, **NWZ** - North Western Zone, **CDZ** - Cauvery Delta Zone.

ern-Era Retrospective analysis for Research and Applications (MERRA 2) satellite data source. Key climatic parameters, including the average mean maximum and minimum temperature (°C), average annual rainfall (mm), average relative humidity (%), and elevation (m) (16) were documented and are presented in Table 2.

### Field survey

A total of 39 block plantations were surveyed to estimate growth biometry across the NEZ, NWZ, and CDZ agroclimatic zones, with a sampling intensity of 5%. Among these, 21 plantations were located in NEZ, 6 in NWZ, and the remaining 12 in CDZ. In all agroclimatic zones, three age classes of teak namely 5-10 years, 10-15 years, and 15-20 years were studied.

### Estimation of growth biometry in teak plantation

Growth biometric parameters, namely Mid Diameter (MD in m), Height (Ht in m), and Clear bole height (CBH in m), were measured (17, 18) using a Laser distance meter (Leica DISTOTM D810). A tree telescope was employed in the study to facilitate the measurement of tree diameters at various heights on standing tree. Using this advantage, the Mid Diameter was measured at 2-m intervals along the tree's height up to the clear bole height (CBH).

For each 2-meter height interval, both the bottom and top diameters were measured, and their average was calculated. By taking the arithmetic mean of all mid-diameters across the 2-meter height intervals for each tree, the average diameter of the standing tree up to the clear bole level was determined. The clear bole volume was then estimated without applying a form factor (19). This methodology ensures that the volume calculated in the study accurately reflects the actual volume of the standing tree, achieving a precision level of approximately 99%.

### Estimation of volume

The volume of standing tree was estimated using the formula proposed by Chaturvedi and Khanna in 1982 (Eqn. 1)

$$V = \pi r^2 h \quad \dots\dots\dots(\text{Eqn. 1})$$

Where, V = Volume, r = Radius at the mid-point, h = Total height of the tree

### Estimation of sap wood (SW), heart wood (HW) and bark content (BC)

In all three agroclimatic zones, for trees across all diameter classes, a sample tree was felled to measure bark thickness, SW thickness, and HW thickness (20). Based on these measurements, the quantities of SW, bark, and HW were calculated, and the SW-to-HW ratio was derived. The growth biometry, in relation to age and climate data, was

statistically analyzed using IBM SPSS and the Agricolae package in the R environment (21).

### Carbon sequestration potential of farm grown Teak

#### Estimation of above ground biomass (AGB)

The above-ground biomass of the plantation was calculated using the formula suggested by pandya (Eqn. 2) in 2013 as:

$$\text{Above ground biomass} = \text{Volume (m}^3\text{)} \times \text{Wood density (g/cm}^3\text{)} \dots\dots\dots(\text{Eqn. 2})$$

Wood density was determined using the water displacement method.

#### Estimation of below ground biomass (BGB)

BGB, which includes live roots biomass but excludes fine roots, was calculated using a root: shoot ratio of 0.26 (Eqn. 3).

$$\text{Below ground biomass (BGB)} = 0.26 \times \text{AGB (ton)} \dots\dots\dots(\text{Eqn. 3})$$

#### Estimation of total biomass (TB)

Total biomass was calculated using the formula suggested by sheikh (Eqn. 4) in 2011 as:

$$\text{Total biomass (TB)} = \text{Above ground biomass (AGB)} + \text{Below ground biomass (BGB)} \dots\dots\dots(\text{Eqn. 4})$$

#### Estimation of weight of carbon (C)

The carbon weight was estimated by the formula given by pearson in 2005 (Eqn. 5) as:

$$\text{Wt. of C} = \text{Biomass} \times 0.50 \dots\dots\dots(\text{Eqn. 5})$$

#### Estimation of total quantity of carbon dioxide (CO<sub>2</sub>)

The total CO<sub>2</sub> equivalent was calculated using the ratio of the molecular weights of carbon dioxide (44 g) to carbon (12 g), as expressed in Equation 6:

$$\text{Total CO}_2 \text{ Equivalent} = \text{Weight of C} \times 3.67 \dots\dots\dots(\text{Eqn. 6})$$

#### Statistical analysis

The experimental databases were statistically analysed under a randomized block design (RBD). Relationships between various parameters were evaluated using Karl Pearson's correlation analysis, conducted via IBM SPSS Statistics 21 Software.

#### Results

Farm-grown teak plantations from the NEZ, NWZ, and CDZ of Tamil Nadu were analysed for their biometric attributes across 3 different age classes. The total height and average mid diameter of *Tectona grandis* were observed to increase progressively with age. Similarly, wood composition, including heartwood, sapwood, and bark content, showed a consistent increase with advancing age classes.

#### Biometric estimation of *Tectona grandis*

The biometric attributes, namely mid diameter, height, and volume, exhibited significant variations across different agroclimatic zones and age classes of farm-grown teak. Among the three agroclimatic zones, teak plantations in the NEZ demonstrated the highest biometric value (Table 3) viz., mid diameter (0.193 m), height (12.92 m), and volume (0.379 m<sup>3</sup>). Conversely, plantation in the cauvery delta zone recorded the lowest biometric values: mid diameter (0.101 m), height (5.80 m), and volume (0.0033 m<sup>3</sup>).


**Table 3.** Growth biometry of farm-grown teak in block plantations

Zones	Age class (Years)	Mid Diameter (m)	Height (m)	Volume (m <sup>3</sup> )
NEZ	5-10	0.118	7.54	0.087
	10-15	0.139	9.04	0.137
	15-20	0.193	12.92	0.379
	<b>Mean</b>	0.150	9.83	0.201
	<b>S.Ed</b>	0.013	0.53	0.052
	<b>CD</b>	0.029	1.18	0.114
	NWZ	5-10	0.128	9.05
10-15		0.135	9.30	0.160
15-20		0.182	9.64	0.304
<b>Mean</b>		0.148	9.33	0.194
<b>S.Ed</b>		0.009	0.10	0.033
<b>CD</b>		0.020	0.22	0.073
CDZ		5-10	0.101	5.80
	10-15	0.141	6.77	0.131
	15-20	0.153	7.20	0.154
	<b>Mean</b>	0.132	6.59	0.114
	<b>S.Ed</b>	0.009	0.24	0.017
	<b>CD</b>	0.020	0.53	0.037

**NEZ** - North Eastern Zone, **NWZ** - North Western Zone, **CDZ** - Cauvery Delta Zone, **S.Ed** - Standard Error of Difference, **CD** - Critical Difference.

### Wood composition of *Tectona grandis*

In addition to biometric attributes, the wood parameters (Fig.1) also showed significant variations across agroclimatic zones and age classes. The teak plantations in the NEZ exhibited the highest values for HW, SW, and bark content volume, with HW volume (0.239 m<sup>3</sup>), SW volume (0.114 m<sup>3</sup>), and bark content volume (0.027 m<sup>3</sup>). The favourable climatic condition, including an average annual rainfall of 1109 mm and medium alluvial soils, in the NEZ likely contributed to these superior values. On the other hand, the CDZ recorded the lowest values for these parameters: HW volume (0.0122 m<sup>3</sup>), SW volume (0.040 m<sup>3</sup>), and bark content volume (0.003 m<sup>3</sup>) (Table 4).



**Fig 1.** Comparison of wood parameters of farm grown teak under three agroclimatic zones. \*significant at the 0.01 level (2-tailed), **NEZ** - North Eastern Zone, **NWZ** - North Western Zone, **CDZ** - Cauvery Delta Zone, **HW** - Heart-

### Carbon sequestration potential of *Tectona grandis*

The carbon sequestration potential of farm-grown teak varied significantly across the agroclimatic zones and age classes. Among the three zones, the NEZ recorded the highest total carbon equivalent (0.683 Mg/acre), followed by NWZ(0.556 Mg/acre), and the CDZ (0.282 Mg/acre).

**Table 4.** Wood composition parameters of farm-grown teak plantations in three agroclimatic zones

Zones	Age Class (Years)	HW Volume (m <sup>3</sup> )	SW Volume (m <sup>3</sup> )	BC Volume (m <sup>3</sup> )
NEZ	5-10	0.031	0.050	0.007
	10-15	0.071	0.054	0.012
	15-20	0.239	0.114	0.027
	<b>Mean</b>	0.114	0.072	0.015
	<b>S.Ed</b>	0.037	0.012	0.003
	<b>CD</b>	0.081	0.026	0.007
NWZ	5-10	0.050	0.055	0.012
	10-15	0.088	0.059	0.013
	15-20	0.204	0.079	0.021
	<b>Mean</b>	0.114	0.064	0.015
	<b>S.Ed</b>	0.027	0.004	0.002
	<b>CD</b>	0.059	0.009	0.004
CDZ	5-10	0.012	0.040	0.003
	10-15	0.046	0.075	0.011
	15-20	0.086	0.054	0.014
	<b>Mean</b>	0.048	0.056	0.009
	<b>S.Ed</b>	0.012	0.006	0.002
	<b>CD</b>	0.026	0.013	0.004

**NEZ** - North Eastern Zone, **NWZ** - North Western Zone, **CDZ** - Cauvery Delta Zone, **HW** - Heartwood, **SW** - Sapwood, **BC** - Bark Content, **S.Ed** - Standard Error of Difference, **CD** - Critical Difference.

When compared across age classes, teak plantations aged 15-20 years exhibited the highest carbon sequestration in all zones (Table 5).

Karl Pearson's correlation analysis was performed to evaluate the relationship between growth biometrics and carbon sequestration potential of farm-grown teak across the three agroclimatic zones in Tamil Nadu. The result indicated a significant positive correlation between biometric parameters and total carbon sequestered by teak (Table 6). There is a positive correlation between mid-diameter ( $k=0.939$ ,  $p<0.01$ ), height ( $k=0.834$ ,  $p<0.01$ ), volume ( $k=1.000$ ,  $p<0.01$ ), above ground biomass ( $k=1.000$ ,  $p<0.01$ ), below ground biomass ( $k=1.000$ ,  $p<0.01$ ), and total carbon ( $k=1.000$ ,  $p<0.01$ ).

These results suggest a strong positive relationship between the growth parameters of teak and its carbon sequestration potential, highlighting the species' significant role in carbon storage.

### Discussion

When comparing the growth biometrics of farm-grown teak across age classes and agroclimatic zones, the highest values were consistently observed in the 15-20 years age class ( $p<0.05$ ). However, for the same age classes, the trees exhibited uniform growth biometrics across all three zones ( $p > 0.05$ ). The interaction effect between zone and age class was not statistically significant for the observed growth biometrics. The variations in growth biometrics were attributed to differences in edaphic and climatic conditions across the agroclimatic zones, as supported by earlier studies (22,23). Similar findings were reported by Reddy and Madiwalar (24), who observed that teak growth in Karnataka was positively influenced by increasing rainfall and favorable soil profiles in different agroclimatic

**Table 5.** Carbon sequestration potential of farm grown Teak

Zones	Age class (Years)	AGB (Mg/acre)	BGB (Mg/acre)	TB (Mg/acre)	Wt. of C (Mg/acre)	Total CO <sub>2</sub> (Mg/acre)
NEZ	5-10	0.069	0.018	0.087	0.044	0.160
	10-15	0.108	0.028	0.137	0.068	0.251
	15-20	0.299	0.078	0.372	0.186	0.683
	<b>Mean</b>	0.159	0.041	0.199	0.099	0.364
	<b>S.Ed</b>	0.024	0.006	0.029	0.015	0.054
	<b>CD</b>	0.052	0.013	0.064	0.033	0.119
NWZ	5-10	0.092	0.024	0.116	0.058	0.214
	10-15	0.127	0.033	0.159	0.080	0.293
	15-20	0.240	0.063	0.303	0.151	0.556
	<b>Mean</b>	0.153	0.040	0.193	0.096	0.354
	<b>S.Ed</b>	0.015	0.004	0.019	0.009	0.035
	<b>CD</b>	0.033	0.009	0.042	0.020	0.077
CDZ	5-10	0.044	0.011	0.055	0.028	0.102
	10-15	0.104	0.027	0.131	0.065	0.240
	15-20	0.121	0.032	0.154	0.077	0.282
	<b>Mean</b>	0.090	0.023	0.113	0.057	0.208
	<b>S.Ed</b>	0.008	0.002	0.010	0.005	0.018
	<b>CD</b>	0.018	0.004	0.022	0.011	0.040

**NEZ** - North Eastern Zone, **NWZ** - North Western Zone, **CDZ** - Cauvery Delta Zone, **AGB** - Above Ground Biomass, **BGB** - Below Ground Biomass, **TB** - Total Biomass, **Wt. of C** - Weight of Carbon, **Total CO<sub>2</sub>** - Total Carbon dioxide Equivalent, **S.Ed** - Standard Error of Difference, **CD** - Critical Difference.

**Table 6.** Correlation between growth biometrics and carbon sequestration potential of farm grown teak

Pearson's Correlation	Mid Diameter (m)	Height (m)	Volume (m <sup>3</sup> )	Above Ground Biomass (Mg/acre)	Below Ground Biomass (Mg/acre)	Total Biomass (Mg/acre)	Weight of Carbon (Mg/acre)	Total Carbon dioxide Equivalent (Mg/acre)
Mid Diameter (m)	1							
Height (m)	.655**	1						
Volume (m <sup>3</sup> )	.939**	.834**	1					
Above Ground Biomass (Mg/acre)	.939**	.834**	1.000**	1				
Below Ground Biomass (Mg/acre)	.939**	.833**	1.000**	1.000**	1			
Total Biomass (Mg/acre)	.939**	.834**	1.000**	1.000**	1.000**	1		
Weight of Carbon (Mg/acre)	.939**	.834**	1.000**	1.000**	1.000**	1.000**	1	
Total Carbon Equivalent (Mg/acre)	.939**	.834**	1.000**	1.000**	1.000**	1.000**	1.000**	1

\*\* . Correlation is significant at the 0.01 level (2-tailed).

zones.

In terms of wood parameters, the maximum values were again recorded for the 15-20 age class (Fig. 1). The study also found that the heartwood ratio increased with age and mid-diameter (MD), while bark content decreased with increasing age and MD. These observations align with the findings of Tewari and Mariswamy (22), who reported similar trends in teak growth under varying agroclimatic zones in Karnataka. Teak grown in the Northeastern Zone exhibited the highest total carbon equivalent compared to the other two zones. This can be attributed to the higher wood density and favorable edaphic and climatic conditions in this zone, which support long-term carbon sequestration under farm-grown conditions. Carbon sequestration in trees is directly linked to biomass, as highlighted by studies conducted by Odusote et al. (25) and Balasooriya et al. (26). Priya and Bhat (27) also reported comparable carbon sequestration potential in teak plantations across

different agroclimatic zones of Southern India, emphasizing the significant influence of climatic and edaphic factors on carbon storage.

Similar studies analyzing the carbon sequestration potential of various teak clones were conducted by Subba and Honnurappa (28), and Rahmawati et al. (29), further corroborating the findings of this study. These studies underscore the importance of zone-specific climatic and soil conditions in enhancing the carbon sequestration capacity and overall growth performance of teak plantations.

## Conclusion

The critical analysis of wood parameters and growth biometrics with respect to age across different agroclimatic zones was conducted. The study concludes that the NEZ, particularly in 15-20 year age class, exhibited the highest growth and wood parameter values. Furthermore, a posi-



tive correlation was established between growth biometrics and the carbon sequestration potential of farm-grown teak, highlighting the Northeastern Zone as having the highest carbon sequestration potential.

These findings suggest that the age of the plantation, along with variations in average climatic parameters across zones, can serve as key indicators for wood quality, carbon sequestration, and growth performance. This information can assist farmers in determining the optimal harvest period for teak plantations, ensuring both economic viability and environmental sustainability.

The scope of this study was limited to three agroclimatic zones in Tamil Nadu. Future research should extend to all agroclimatic zones of the state to provide a more comprehensive comparison of the growth, wood quality, and carbon sequestration potential of farm-grown teak. Such studies would enhance the precision and applicability of these findings for broader agricultural and forestry practices.

## Acknowledgements

Authors would like to thank the Indian Council of Forestry Research and Education (ICFRE), Dehradun for providing financial support through All India Coordinated Research Project on Silvicultural interventions for productivity enhancement and carbon sequestration in plantations of important tree species

## Authors' contributions

SN, AB, MS and RR contributed in the conceptualization, investigation, and preparation of original draft. BS, CN, GS, VM and KS contribute in review, editing, and preparation of the final draft.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interests to declare.

**Ethical issues:** None

## References

- Balakrishnan S, Dev SA, Sakthi AR, Vikashini B, Bhasker T R, Magesh NS, Ramasamy Y. Gene-ecological zonation and population genetic structure of *Tectona grandis* Lf in India revealed by genome-wide SSR markers. *Tree Genetics & Genomes*. 2021; 17:1–4. <https://doi.org/10.1007/s11295-021-01514-x>
- Kidanu S, Mamo T, Stroosnijder L. Biomass production of Eucalyptus boundary plantations and their effect on crop productivity on Ethiopian highland Vertisols. *Agroforestry systems*. 2005;63:281–290. <https://doi.org/10.1007/s10457-005-5169-z>
- Vongkhamho S, Imaya A, Yamamoto K, Takenaka C, Yamamoto H. Influence of topographic conditions on teak growth performance in mountainous landscapes of Lao PDR. *Forests*. 2022; 13(1):118. <https://doi.org/10.3390/f13010118>
- Deb JC, Phinn S, Butt N, McAlpine CA. Climatic-induced shifts in the distribution of teak (*Tectona grandis*) in tropical Asia: Implications for forest management and planning. *Environmental management today*. 2017 Sep; 60:422–435. <https://doi.org/10.1007/s00267-017-0884-6>
- Ridder RM. Global forest resources assessment 2010: options and recommendations for a global remote sensing survey of forests. *FAO For. Resour. Assess. Programme Work. Pap.* 2007 Mar;141.
- Kollert W, Cherubini L. Teak resources and market assessment 2010. *FAO Planted Forests and Trees Working Paper FP/47/E*, Rome. 2012 Apr; 206. <http://www.fao.org/forestry/plantedforests/67508@170537/en/>
- Shrivastava S, Saxena AK. Wood is Good: But, is India doing enough to meet its present and future needs. *Centre for Science and Environment, New Delhi*. 2017:13.
- Pandey D, Brown C. Teak: a global overview. *UNASYLVA-FAO*. 2000 Jan 1:3–13. <http://www.apps.fao.org>
- Sani Lookose SL. Traditional teak wood articles used in households of Nilambur and Malapuram areas of Kerala. <http://nopr.niscpr.res.in/handle/123456789/581>
- Miranda I, Sousa V, Pereira H. Wood properties of teak (*Tectona grandis*) from a mature unmanaged stand in East Timor. *Journal of wood science*. 2011 Jun; 57:171–178. <https://doi.org/10.1007/s10086-010-1164-8>
- Palanisamy K, Hegde M, Yi JS. Teak (*Tectona grandis* Linn. f.): A renowned commercial timber species. *Journal of Environmental Science International*. 2009; 25(1):1–24.
- Arunachalam B, Natarajan HP, Satheesan A, Ganesan S, Suthandhirajan R, Muthuswamy S, Vellan M, Selvaraj NK. Yield model and yield table construction in Albizia (*Albizia lebbek* L.) under the western agro-climatic zone of Tamil Nadu, Southern India. *Environmental science and pollution research international*. 2024 Jun; 31(27):38781–38787. <https://doi.org/10.1007/s11356-023-29716-x>
- Buvaneshwaran C, George M, Perez D, Kanninen M. Biomass of teak plantations in Tamil Nadu, India and Costa Rica compared. *Journal of Tropical Forest Science*. 2006 Jul 1;18(3):195–197.
- Shukla SR, Viswanath S. Comparative study on growth, wood quality and financial returns of teak (*Tectona grandis* Lf) managed under three different agroforestry practices. *Agroforestry Systems*. 2014 Apr;88:331–341. <https://doi.org/10.1007/s10457-014-9686-5>
- Sreejesh KK, Thomas TP, Rugmini P, Prasanth KM, Kripa PK. Carbon sequestration Potential of Teak (*Tectona grandis*) plantations in Kerala. *Research journal of recent sciences*. 2013;2 (ISC-2012):167–170.
- Wiemann MC, Williamson GB. Geographic variation in wood specific gravity: effects of latitude, temperature, and precipitation. *Wood and Fiber Science*. 2002:96–107.
- Ravi J, Radhakrishnan S, Balasubramanian A. Construction of yield model for *Ailanthus excelsa* grown in Cauvery delta agro climatic zone of Tamil Nadu, India. *International journal of chemical studies*. 2018;7:3852–3854.
- Raviperumal A, Balasubramanian A, Manivasakam S, Prasath CH, Krishnan MS. Yield model construction in neem (*Azadirachta indica*) under western agro climatic zone of Tamil Nadu, Southern India. *Res Sq*. 2018:1–4.
- Zahabu E, Raphael T, Chamshama SA, Iddi S, Malimbwi RE. Effect of spacing regimes on growth, yield, and wood properties of *Tectona grandis* at Longuza forest plantation, Tanzania. *International journal of research*. 2015; 2015(1):469760. <https://doi.org/10.1155/2015/469760>
- Rajan LJ, Ramanan SS, Anoop EV. Physical and Anatomical-wood properties variation among the provenances of Red Sanders (*Pterocarpus santalinus*). *Indian Journal of Hill Farming*.

- 2019;32(1).
21. Mendiburu FD, Simon R. Agricola-Ten years of an open source statistical tool for experiments in breeding, agriculture and biology. Peer J PrePrints. 2015. <https://dx.doi.org/10.7287/peerj.preprints.1404v1>
  22. Tewari VP, Mariswamy KM. Heartwood, sapwood and bark content of teak trees grown in Karnataka, India. Journal of Forestry Research. 2013 Dec; 24:721–725. <https://doi.org/10.1007/s11676-013-0410-5>
  23. Naidu LG, Ramamurthy V, Srinivas S, Natarajan A, Thayalan S. Delineation of Agro-ecological Zones of Tamil Nadu. Agropedology. 2013; 23(2):83–92.
  24. Reddy MC, Madiwalar SL. Productivity assessment and economic analysis of teak plantations in different agro climatic zones of Karnataka. Indian For. 2014;140(3):287–290. <http://www.indianforester.co.in/>
  25. Odusote JK, Adeleke AA, Lasode OA, Malathi M, Paswan D. Thermal and compositional properties of treated *Tectona grandis*. Biomass Conversion and Biorefinery. 2019;9:511–519. <https://doi.org/10.1007/s13399-019-00398-1>
  26. Balasooriya B, Amarasinghe W, Weerakkody W, Muthumala C. Development of Bucking Techniques to Improve Timber Grading of Teak (*Tectona grandis*). In Proceedings of 19th Agricultural Research Symposium 2020;525:p. 529).
  27. Priya PB, Bhat KM. False ring formation in teak (*Tectona grandis* Lf) and the influence of environmental factors. Forest Ecology and Management. 1998; 108(3):215–222. [https://doi.org/10.1016/S0378-1127\(98\)00227-8](https://doi.org/10.1016/S0378-1127(98)00227-8)
  28. Subba S, Honnurappa S. Exploring Carbon Sequestration Potential Across Various Clones in Teak Plantation. International Journal of Economic Plants. 2024;11(1):056–059. <https://doi.org/10.23910/2/2024.5121>
  29. Rahmawati RB, Hardiwinoto S, Wibowo A. Productivity of Clonal Teak Plantation Under Different Spacing and Thinning Intensity in Java Monsoon Forest. InIOP Conference Series: Earth and Environmental Science 2024;1299(1): 012004.IOP Publishing. <https://doi.org/10.1088/1755-1315/1299/1/012004> .