



RESEARCH ARTICLE

Influence of naphthalene acetic acid, gibberellic acid and triacontanol on fruit retention, yield and quality of mango (*Mangifera indica* L.) cv. Banganpalli

Deepa Samant*, Kundan Kishore, Suchitra Behera & Gobinda Chandra Acharya

ICAR-IIHR-Central Horticultural Experiment Station, Bhubaneswar 751 019, Odisha, India

*Email: horti.deepa@gmail.com

 OPEN ACCESS

ARTICLE HISTORY

Received: 24 October 2024
Accepted: 10 December 2024
Available online
Version 1.0 : 27 January 2025
Version 2.0 : 30 January 2025



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Samant D, Kishore K, Behera S, Acharya GC. Influence of naphthalene acetic acid, gibberellic acid and triacontanol on fruit retention, yield and quality of mango (*Mangifera indica* L.) cv. Banganpalli. Plant Science Today. 2025; 12(1): 1-7. <https://doi.org/10.14719/pst.6112>

Abstract

A field investigation was carried out on fifteen-year-old mango cv. Banganpalli during 2021-2023 in the coastal region of India at the research farm of ICAR-IIHR-Central Horticultural Experiment Station, Bhubaneswar, Odisha to evaluate the efficacy of some plant growth regulators, namely naphthalene acetic acid (NAA), gibberellic acid (GA₃) and triacontanol (TRIA) for improving fruit retention, yield and quality of harvest. The experiment was laid out in randomized block design with 10 treatments consisting of NAA (10, 20 and 30 ppm), GA₃ (25, 50 and 75 ppm), TRIA (1, 3 and 5 ppm) and water spray as control. Each treatment was replicated thrice with four plants per replication. Treatments were applied on the plant canopy thrice at panicle initiation, pea and marble stages of fruit growth. Observations were recorded on flowering, fruiting, yield and fruit quality indices. Application of triacontanol at 3-5 ppm brought out a significant improvement over control in terms of panicle size, fruit retention and yield. Plants sprayed with 5 ppm triacontanol produced the largest panicle (length: 29.17 cm, width: 18.07 cm) and recorded the maximum value for fruit retention (68.93, 53.38, 39.41, 32.88, 26.54, 20.46 and 16.58% at 15, 30, 45, 60, 75, 90 and 105 days after pea stage, respectively), number of fruits/tree (104.72) and yield (38.95 kg/tree). With respect to fruit quality, GA₃ and TRIA exhibited significant influence on fractions of fruit, dry matter and on most of the chemical attributes (TSS, TSS/acid ratio, total sugar, reducing sugar, non-reducing sugar and vitamin C) when applied at the concentration of 50-75 and 3-5 ppm, respectively.

Keywords

chlorophyll; fruit retention; fruit quality; gibberellic acid; mango yield; triacontanol

Introduction

Mango (*Mangifera indica* L.; Anacardiaceae; 2n=2x=40) is one of the most popular and widely grown fruits in the tropics and sub-tropics of the world on account of its varietal wealth, production volume, versatile uses, enchanting shades of colour, delightful taste, unique flavour, captivating aroma, excellent nutritional properties and high marketability. It is believed to have originated in the Indo-Myanmar region (1) during the early Cretaceous era (2) and gradually spread throughout Asia and tropical and subtropical regions worldwide over the past 2500 years by travellers, traders and rulers (3). Presently, it is grown in approximately 140 countries (4), however, the mass-scale popularity, economic significance and extensivity of cultivation it enjoys in India is unmatched. Mango has a rich cultivation history of 4,000-6,000 years in India and is deeply intertwined with Indian art, scriptures, history, culture and traditions (5). India is the global leader in mango production with 2.26 million hectares of area and 21.82 million tonnes of

production (6). Despite its rich legacy in mango cultivation and unparalleled global production, the country falls short in tapping the full horticultural potential of mango in terms of productivity. The average productivity of mango in India is 9.66 tonnes/hectare, which is way lower than that realized in other countries, viz., 30 tonnes/hectare in Israel (7). One of the most common and critical constraints in realizing the higher yield potential of mango despite adequate flowering and initial fruit set, is the high magnitude of fruit drop.

Mango fruit drop is a highly complex and coordinated physiological process that involves the formation of a separation layer in the abscission zone, located at the junction of the pedicel and peduncle. Induction of separation layer is regulated by a variety of stress stimuli of biotic and abiotic nature, viz., inadequate pollination, self-incompatibility, incidence of insect pests and diseases and unfavourable soil and climatic conditions responsible for embryo abortion, hormonal imbalance (low levels of auxin, gibberellins and cytokinin coupled with high levels of ethylene and abscisic acid), expression of cell wall degrading enzymes (cellulase and pectinase) and sink rivalry in plant (8, 9). Studies carried out by various investigators suggest the use of plant growth regulators (PGRs), particularly naphthalene acetic acid (NAA), gibberellic acid (GA_3) and triacontanol (TRIA) in mango during flowering and fruiting for controlling the premature abscission of fruits and improving the quantum and quality of harvest (10-15). However, the response of crops to the PGRs may differ with the species, variety, climatic conditions and PGR application timing and concentration. Keeping all these aspects in view, the present investigation was, therefore, undertaken to assess the effects of foliar application of NAA, GA_3 and TRIA on fruit retention, yield and quality in mango cv. Banganpalii in the tropical eastern region of India.

Materials and Methods

The current field trial was conducted at ICAR-IIHR-Central Horticultural Experiment Station, Bhubaneswar, Odisha, India (20°15' N, 85°15' E, 25.5 m amsl) during 2021-2023 on a high-density mango orchard (5 m × 5 m) of cv. Banganpalli, which was planted in 2005 and maintained under uniform cultural practices. The site experiences a hot and humid climate, receiving an annual rainfall of 1400-1500 mm, primarily from June to September. The soil in experimental orchard was red lateritic with sandy loam texture (80.72% sand, 10.65% silt and 8.63% clay), acidic reaction (pH 4.8) and low levels of organic carbon (0.25%) and available nitrogen (191.74 kg/ha), phosphorus (16.45 kg/ha) and potassium (119.86 kg/ha). The trial was carried out employing a randomized block design with 10 treatments and 3 replications. The treatments comprised of varying concentrations of three PGRs which are reported to reduce fruit drop in mango, viz., NAA (T1: 10, T2: 20 and T3: 30 ppm), GA_3 (T4: 25, T5: 50 and T6: 75 ppm) and TRIA (T7: 1, T8: 3 and T9: 5 ppm) and water spray as control (T10). Treatments were applied on the plant canopy thrice during panicle initiation, pea and marble stages of fruit development. Observations were recorded on leaf chlorophyll content; flowering and fruiting characteristics, viz., flowering intensity (%), number of flowers per panicle, hermaphrodite flower intensity (%), panicle size in terms of length and width (cm) and fruit retention (%); yield (kg/

tree) and yield contributing attributes, viz., average fruit weight (g/fruit) and no. of fruits/tree; and on fruit quality indices, viz., fractions of fruit (%), dry matter content (%), TSS (°B), acidity (% equivalent of citric acid), TSS/acid ratio, sugar content (%), vitamin C (mg/100g of pulp) and total carotenoid ($\mu\text{g}/100\text{g}$ of pulp).

To measure leaf chlorophyll, 20 vegetative shoots that emerged after the first spray of treatments were randomly tagged on each tree covering all four directions of the canopy. Once shoots reached six months of maturity, two leaves were selected from the central portion and the chlorophyll content was measured at 10 random points while avoiding the midrib and direct sunlight, using a portable chlorophyll meter, called atLEAF. The results were expressed as atLEAF units (9, 16). For determining flowering intensity, all three types of shoots (floral, vegetative and dormant) were counted per square meter of tree canopy in all four directions with the help of quadrat and the following formula was used (17)-

$$\text{Flowering intensity (\%)} = \frac{\text{No. of floral shoots}}{\text{Total number of shoots}} \times 100$$

Total number of shoots = dormant shoots + vegetative shoots + floral shoots

To record flowering and fruiting parameters, 20 panicles were randomly tagged across the plant canopy at full bloom stage. A measuring tape was used to record panicle size. The length was measured from base to tip, while the width was at its broadest part. The intensity of the hermaphrodite flower was worked with the help of the following formula (18)-

$$\text{Hermaphrodite flower intensity (\%)} = \frac{\text{No. of hermaphrodite flowers per panicle}}{\text{Total number of flowers per panicle}} \times 100$$

Once the mango tree entered fruit setting phase, fruits were counted periodically on tagged panicles at fortnightly interval starting from pea stage to harvesting stage, subsequently fruit retention at 15, 30, 45, 60, 75, 90 and 105 days after pea stage (DAPS) was computed using following formula (9, 19)-

$$\text{Fruit retention (\%)} = \frac{\text{No. of fruits retained}}{\text{No. of fruits at pea stage}} \times 100$$

On reaching optimum maturity, fruits were harvested manually and the produce was weighed and counted for the purpose of yield data. The average fruit weight was computed using following formula (19).

$$\text{Average fruit weight (g/fruit)} = \frac{\text{Yield (kg/tree)}}{\text{Number of fruits per tree}} \times 1000$$

A random sample of 10 fruits was drawn from the harvest lot of each tree and used for the determination of physico-chemical attributes of fruit quality. Fractions of fruit, i.e., peel,

pulp and stone, were weighed and expressed as percentages. Ten grams of fruit pulp was oven dried at 60 °C until it reaches to a constant dry weight, thereafter following formula was used for the calculation of dry matter (19) -

$$\text{Dry matter content (\%)} = \frac{\text{Dry weight of pulp (g)}}{\text{Fresh weight of pulp (g)}} \times 100$$

The total soluble solids (TSS) in fruit pulp were measured with the help of portable digital refractometer (Hanna HI 96801, 0-85%). Estimation of acidity, sugar (reducing, non-reducing and total sugar), vitamin C and total carotenoid was performed as per the standard methods (20). The field and laboratory data collected over three consecutive years (2021, 2022 and 2023) were pooled and subjected to statistical analysis using OPSTAT, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, India.

Results and Discussion

Flowering

The observations recorded on some important characteristics of flowering are presented in Table 1. No significant improvements were observed in flowering intensity, no. of flowers per panicle and hermaphrodite flower percentage on application of NAA, GA₃ and TRIA, however, a significant improvement in panicle size was recorded with the application of 3-5 ppm TRIA. In PGR treatments, flowering intensity, no. of flowers per panicle and hermaphrodite flower percentage ranged between 67.92-71.57%, 1070.64-1287.49 and 12.72-14.66%, respectively, as against their corresponding values of 68.23%, 1096.90 and 12.15% noted under control. The T9-treated (5 ppm TRIA) mango trees recorded the largest panicles (29.17 cm × 18.07 cm), followed by 3 ppm TRIA (28.65 cm × 17.86 cm). The panicle size in the rest of the PGR treatments (21.66-24.06 cm × 12.07-14.34 cm) did not exhibit notable differences and remained comparable to the control (21.43 cm × 12.41 cm).

The stimulatory effect of triacontanol on mango panicle could be explained by its role in activation of growth-promoting second messenger called L(+)-adenosine and in enhancing the photosynthetic efficiency of plant via upregulation of genes

(rbcS) and enzymes (ribulose-1,5-bisphosphate carboxylase, phosphoenolpyruvate carboxylase, carbonic anhydrase, malate dehydrogenase, nitrate reductase) involved in the process and by improving the functioning and expression of various components of photosynthetic apparatus, viz., pigments, membrane permeability, stomatal conductance and PSII (21-27). Our study revealed significant enhancement in leaf chlorophyll due to the application of 3-5 ppm TRIA (Fig. 1) which could be attributed to its positive influence on the size and number of the chloroplast (28). Significant effects of TRIA on growth and leaf chlorophyll content have also been reported earlier in various horticultural crops (9, 26, 29).

Fruit retention and yield

The data pertaining to fruit retention and yield are presented in Table 2. All the PGR treatments and control exhibited a decreasing trend of fruit retention with the advancement of fruit development period. The maximum fruit retention was observed during initial phase of fruit development, i.e., at 15 DAPS (50.97 to 68.93%), whereas, at the final stage of fruit development or maturity, i.e., 105 DAPS, it was minimum (12.17 to 16.58%). The perusal of data in Table 2 further revealed the effectiveness of TRIA in exerting significant influence on fruit retention over NAA and GA₃, when applied at the concentration of 3-5 ppm. The treatment T9 (5 ppm TRIA) recorded the highest fruit retention, which was at par with the T8 (3 ppm TRIA). Fruit retention recorded in 1 ppm TRIA (T7) and all tried concentrations of other PGRs, namely NAA and GA₃ (T1, T2, T3, T4, T5 and T6) were statistically at par with the control.

Triacontanol-induced improvement in fruit retention as observed in the present study might be due to its inhibitory effect on hydrolytic enzymes, viz., cellulase and pectinase responsible for cell wall dissolution (30) and on abscission-inducing plant hormone, called ABA (31). Additionally, TRIA is suggested to enhance the source capacity and sink strength in plants on account of its positive influence on leaf area, water and nutrient uptake, photosynthetic pigments and enzymes and translocation and accumulation of photosynthates, whereas antagonistic effect on respiration and transpiration, which, in turn, could have reduced the fruit abscission and enhanced the fruit retention (27, 32-34). Our study supports TRIA's role in strengthening the source capacity, since mango trees sprayed with 3-5 ppm TRIA had notably high contents of leaf chlorophyll,

Table 1. Effect of plant growth regulators on flowering in mango cv. Banganpalli

Treatment	Flowering intensity (%)	No. of flowers/panicle	Hermaphrodite flower (%)	Panicle dimension (cm)	
				Length	Width
T1	68.65	1105.38	12.72	21.92	12.71
T2	69.29	1070.64	13.19	21.66	12.07
T3	71.57	1145.76	13.32	22.47	13.10
T4	70.24	1156.85	13.33	22.82	12.43
T5	67.92	1180.72	13.04	23.85	13.67
T6	71.52	1196.37	13.98	24.06	14.34
T7	70.88	1164.28	13.51	23.49	14.04
T8	69.96	1268.15	14.13	28.65	17.86
T9	69.71	1287.49	14.66	29.17	18.07
T10	68.23	1096.90	12.15	21.43	12.41
SE(m)±	2.10	130.47	1.40	1.36	1.02
CD (P=0.05)	ns	ns	ns	4.12	3.10


T1: 10 ppm NAA, T2: 20 ppm NAA, T3: 30 ppm NAA, T4: 25 ppm GA₃, T5: 50 ppm GA₃, T6: 75 ppm GA₃, T7: 1 ppm TRIA, T8: 3 ppm TRIA, T9: 5 ppm TRIA, T10: Control, ns: non-significant

Table 2. Effect of plant growth regulators on fruit retention and yield in mango cv. Banganpalli

Treatment	Fruit retention (%)							Fruit yield		
	15 DAPS	30 DAPS	45 DAPS	60 DAPS	75 DAPS	90 DAPS	105 DAPS	No. of fruits/tree	Fruit weight (g/fruit)	Yield (kg/tree)
T1	52.08	39.67	29.81	24.86	20.05	15.44	12.45	79.54	371.70	29.56
T2	53.18	41.15	30.45	25.37	20.48	15.78	12.76	80.91	372.56	30.12
T3	55.12	42.67	31.52	26.35	21.23	16.35	13.20	84.50	373.15	31.54
T4	53.80	41.65	30.76	25.67	20.72	15.96	12.85	81.75	369.81	30.21
T5	56.14	43.58	32.13	26.78	21.61	16.65	13.45	84.14	370.50	31.16
T6	58.32	45.12	33.06	27.73	22.35	17.31	13.82	88.12	371.34	32.71
T7	57.45	44.46	32.85	27.42	22.12	17.04	13.76	87.12	371.65	32.35
T8	66.51	51.50	38.10	31.76	25.62	19.78	15.95	101.21	372.80	37.72
T9	68.93	53.38	39.41	32.88	26.54	20.46	16.58	104.72	372.14	38.95
T10	50.97	39.45	29.17	24.48	19.67	15.12	12.17	77.46	371.35	28.76
SE(m)±	2.52	1.95	1.40	1.14	0.90	0.89	0.61	3.70	3.72	1.41
CD (P=0.05)	7.60	5.81	4.17	3.42	2.73	2.61	1.78	11.15	ns	4.17

T1: 10 ppm NAA, T2: 20 ppm NAA, T3: 30 ppm NAA, T4: 25 ppm GA₃, T5: 50 ppm GA₃, T6: 75 ppm GA₃, T7: 1 ppm TRIA, T8: 3 ppm TRIA, T9: 5 ppm TRIA, T10: Control, ns: non-significant

which is considered an important indicator of photosynthetic efficiency (Fig. 1). The beneficial effect of triacontanol on fruit retention has also been reported earlier in a variety of fruit crops, such as mango (10, 35), olive (36), pomegranate, (37), mandarin (38) and apple (39).

**Fig. 1.** Effect of plant growth regulators on leaf chlorophyll content in mango cv. Banganpalli.

T1: 10 ppm NAA, T2: 20 ppm NAA, T3: 30 ppm NAA, T4: 25 ppm GA₃, T5: 50 ppm GA₃, T6: 75 ppm GA₃, T7: 1 ppm TRIA, T8: 3 ppm TRIA, T9: 5 ppm TRIA, T10: Control.

Foliar application of TRIA (3-5 ppm) not only enhanced the fruit retention but also yielded significant improvement in the quantum of harvest over control. Plants sprayed with 5 ppm TRIA (T9) recorded the highest quantum of fruit yield (38.95 kg/tree, 104.72 fruits/tree), which remained statistically comparable to the yield noted in case of T8 (3 ppm TRIA). On the other hand, the lowest quantum of produce was observed control plants (28.76 kg/tree, 77.46 fruits/tree). Concerning fruit weight, it ranged between 369.81 to 373.15 g/fruit with a mean of 371.74 g/fruit under PGR treatments. However, none of the PGR treatments registered significant improvement over control (371.35 g/fruit). The pronounced effect of TRIA (3-5 ppm) on fruit yield in the present investigation could be resultant of better fruit retention, as other variables influencing the quantum of fruit harvest, viz., flowering intensity, no. of flowers per panicle, hermaphrodite flower % and fruit weight remained unaffected (Table 1 and Table 2). Similar effects of TRIA on quantum of mango harvest have been documented previously (15, 33, 35, 40).

Fruit quality

The data recorded on various physical and chemical fruit quality attributes are presented in Table 3. Of three PGRs, GA₃ and TRIA exhibited significant influence on fractions of fruit, dry matter and on most of the chemical parameters studied except acidity and total carotenoid, when applied at the concentration of 50-75 and

Table 3. Effect of plant growth regulators on fruit quality in mango cv. Banganpalli

Treatment	Pulp (%)	Peel (%)	Seed (%)	Dry matter (%)	TSS (°B)	Acidity (%)	TSS/ Acid ratio	Total Sugar	Reducing sugar (%)	Non-reducing	Vit. C (mg/100g)	Total carotenoid
T1	69.89	16.69	13.42	16.12	17.56	0.43	40.84	15.48	9.35	6.13	32.46	795.45
T2	68.72	17.20	14.08	15.67	17.75	0.41	43.29	14.96	9.12	5.84	31.74	835.16
T3	70.10	16.75	13.15	16.17	18.12	0.44	41.18	15.31	9.15	6.16	32.68	819.42
T4	69.41	16.96	13.63	15.89	18.37	0.42	43.74	15.25	9.21	6.04	33.25	876.97
T5	74.63	14.12	11.25	18.64	19.58	0.39	50.21	17.74	10.72	7.02	38.89	841.56
T6	75.54	13.70	10.76	19.35	19.73	0.37	53.32	18.12	10.97	7.15	40.97	902.28
T7	69.60	17.05	13.35	16.25	18.20	0.43	42.33	15.63	9.56	6.07	33.18	798.95
T8	74.92	13.90	11.18	18.97	19.65	0.40	49.13	17.90	10.83	7.07	39.63	837.14
T9	75.17	13.86	10.97	19.78	19.81	0.38	52.13	18.31	11.06	7.25	41.21	865.71
T10	69.16	17.30	13.54	15.72	17.68	0.43	41.12	14.75	9.03	5.72	31.25	810.37
SE(m)±	1.38	0.72	0.46	0.72	0.27	0.03	1.72	0.58	0.26	0.21	1.71	67.65
CD @5%	4.12	2.17	1.35	2.14	0.86	ns	5.21	1.72	0.78	0.65	5.16	ns

T1: 10 ppm NAA, T2: 20 ppm NAA, T3: 30 ppm NAA, T4: 25 ppm GA₃, T5: 50 ppm GA₃, T6: 75 ppm GA₃, T7: 1 ppm TRIA, T8: 3 ppm TRIA, T9: 5 ppm TRIA, T10: Control, ns: non-significant

3-5 ppm, respectively. Among four effective PGR treatments (T5, T6, T8 and T9), the treatment T6 (75 ppm GA₃) registered the highest pulp content (75.54%) and the lowest contents of peel (13.70%) and seed (10.76%), followed by T9 (5 ppm TRIA), T8 (3 ppm TRIA) and T5 (50 ppm GA₃), however, the variations were non-significant. Fruits with the highest dry matter content (19.78%) were harvested from T9-treated plants, whereas the lowest with T2 treatment (15.67%), which was comparable with the content recorded under control and other non-effective PGR treatments (NAA: T1 and T3; GA₃: T4; TRIA: T7). Similar effects of GA₃ and TRIA on fruit fractions and dry matter have been reported previously in mango, mandarin and sugar apple (9, 10, 41-43).

Perusal of fruit quality data further revealed that effective PGR treatments (T5: 50 ppm GA₃, T6: 75 ppm GA₃, T8: 3 ppm TRIA and T9: 5 ppm TRIA) did not vary significantly with each other for various chemical attributes. Of these, 5 ppm TRIA scored the highest value for TSS (19.81°B), total sugar (18.31%), reducing sugar (11.06%), non-reducing sugar (7.25%) and vitamin C (41.21 mg/100 g pulp), whereas the treatment 75 ppm GA₃ recorded the maximum value for TSS acid ratio (53.32). Control plants recorded the lowest value for total sugar (14.75%), reducing sugar (9.03%), non-reducing sugar (5.72%) and vitamin C (31.25 mg/100 g pulp), on the other hand, the lowest value for TSS and TSS/acid ratio were observed under 10 ppm NAA (17.56 °B and 40.48, respectively). Triacetonol and GA₃-mediated improvement in various chemical attributes of fruit quality have been noticed previously by several researchers (33, 40, 44-47).

The positive influence of GA₃ on various parameters of fruit quality could be ascribed to its role in cell division, cell enlargement, cell elongation, water absorption, nutrient uptake, metabolite synthesis, conversion of complex polysaccharides into simple and soluble sugar molecules, assimilate partitioning, dry matter accumulation, and synthesis of vitamin C precursor called glucose-6-phosphate (44, 48-50). Likewise, TRIA also regulates an array of metabolic, physiological, and biochemical processes in plants including cell division, elongation, expansion, and differentiation (33, 51). Exogenous application of TRIA has stimulatory effects on photosynthesis, enzymatic activity, water uptake, nutrient acquisition, carbohydrate metabolism, translocation, and accumulation of metabolites and photosynthates to the sink, all of which in turn could have resulted in enhancement in physical and chemical attributes of fruit quality (34, 42, 52).

Conclusion

Based on experimental results, it could be concluded that in mango cv. Banganpalli, three foliar sprays of triacetonol @ 3-5 ppm during panicle initiation, pea and marble phases of fruit development are efficacious for controlling premature fruit shedding and improving the yield and quality of harvest.

Acknowledgements

Authors are thankful to the Director, ICAR-IIHR, Bengaluru, India and Head, ICAR-IIHR-CHES, Bhubaneswar, India for providing facilities during the study period.

Authors' contributions

DS was responsible for conceptualization, investigation, methodology, data recording, statistical analysis, writing the original draft and reviewing and editing the manuscript. KK contributed to methodology and data recording. SB focused on data recording, while GCA provided methodological input, technical guidance and monitoring of the experiment.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

References

- Mukherjee SK. Origin of mango. *Indian J Genet Plant Breed.* 1951;11 (1):49-56.
- Yonemori K., Honsho C, Kanzaki S., Eiadthong W, Sugiura A. Phylogenetic relationships of *Mangifera* species revealed by ITS sequences of nuclear ribosomal DNA and a possibility of their hybrid origin. *Plant Syst Evol.* 2002; 231:59-75.
- Singh NK, Mahato AK, Jayaswal PK, Singh A, Singh S, Singh N et al. Origin, diversity and genome sequence of mango (*Mangifera indica* L.). *Indian J Hist Sci.* 2016;51.2.2:355-68. <https://doi.org/10.16943/ijhs/2016/v51i2.2/48449>
- Rajan S. Mango- The king of fruits. *Indian Hort.* 2021;66(4):3-5.
- Pandey SN, Dinesh MR. Mango. New Delhi (India): Directorate of Information and Publications of Agriculture, ICAR; 2010.
- Fruit Production Database, National Horticulture Board, New Delhi, India, 2017-18. <https://www.nhb.gov.in/>
- Parthiban S, Santhi VP, Snehapriya MS, Indumathi K, Masilamani P. Recent advances in enhancing the productivity of mango (*Mangifera indica* L.) through hi-tech practices. *Int J Curr Microbiol App Sci.* 2020;9(8):1850-64. <https://doi.org/10.20546/ijcmas.2020.908.212>
- Singh Z, Mallik AU, Davenport TL. Fruit drop in mango. In: Janick J, editor. *Horticultural Reviews.* USA: John Wiley & Sons; 2005. p.111-53.
- Dash A, Samant D, Dash DK, Dash SN, Mishra KN. Influence of *Ascophyllum nodosum* extract, homobrassinolide and triacetonol on fruit retention, yield and quality of mango. *J Environ Biol.* 2021;42(4):1085-91. <http://doi.org/10.22438/jeb/42/4/MRN-1541>
- Patil AS, Tidke SN, Tike MA, Shinde BN, Gore AK. Effect of chemicals and growth regulators on fruit set and fruit retention of Parbhani Bhushan mango. *J Soils and Crops.* 2005;15(1):64-6.
- Vejendla V, Maity PK, Banik B. Effect of chemicals and growth regulators on fruit retention, yield and quality of mango cv. Amrapali. *J Crop and Weed.* 2008;4(2):45-6.
- Ahmed W, Tahir FM, Rajwana IA, Raza SA, Asad HU. Comparative evaluation of plant growth regulators for preventing premature fruit drop and improving fruit quality parameters in 'Dusehri' mango. *Int J Fruit Sci.* 2012;12:372-89. <https://doi.org/10.1080/15538362.2012.679175>
- Ghosh SN. Effect of growth regulators on fruit retention and physico-chemical properties of mango cv. Amrapali grown in lateritic soil at close spacing. *J Crops and Weed.* 2016;12(3):83-5.
- Sahu C, Sahu DK. Effect of plant growth regulators on yield and yield attributing characteristics of mango (*Mangifera indica* L.) cv. Dashehari. *Int J Hort Food Sci.* 2020;2(1):52-4.
- Gaikwad RB, Nainwad RV, Gaikwad AR, Jadhav SD, Naikwadi PB, Gaikwad GB, Karakai SP and Gite AS. Effect of chemicals and biomix on fruit setting & yield of mango cv. Kesar. *Pharma Inno J.* 2023;12

- (11):2351-52.
16. Samant D and Kishore K. Standardization of pruning for high density Sardar guava orchards under hot and humid climate of Eastern India. *Indian J Hort.* 2019;76(1):70-4. <https://doi.org/10.5958/0974-0112.2019.00010.0>
 17. Kishore K, Singh HS, Kurian RM, Srinivas P and Samant D. Performance of certain mango varieties and hybrids in East Coast of India. *Indian J Plant Genet Resour.* 2015; 28: 296-302. <https://doi.org/10.5958/0976-1926.2015.00038.8>
 18. Samant D, Kishore K, Singh HS. Assessing effectiveness of 'Arka Mango Special' for improving yield and quality of mango variety 'Banganpalli' in lateritic soils of Odisha, India. *Int J Curr Microbiol App Sci.* 2018;7:168-73. <https://doi.org/10.20546/ijcmas.2018.701.018>
 19. Samant D, Mandal S, Nath V and Kurian RM. Effect of deficit drip irrigation and mulching on mango hybrid 'Amrapali'. *J Indian Soc Coastal Agric Res.* 2017; 35(2):53-8.
 20. Ranganna S. Handbook of analysis and quality control for fruit and vegetable products. New Delhi (India): McGraw-Hill Education (India) Private Limited; 2019.
 21. Houtz RL, Ries SK, Tolbert NE. Effect of triacontanol on *Chlamydomonas* II. Specific activity of ribulose-bisphosphate carboxylase/oxygenase, ribulose-bisphosphate concentration and characteristics of photorespiration. *Plant Physiol.* 1985;79(2):365-70. <https://doi.org/10.1104/pp.79.2.365>
 22. Savithiry S, Wert V, Ries S, Influence of 9-b-L (+)- adenosine on malate dehydrogenase activity in rice. *Physiol Plantarum.* 1992;84(3):460-66. <https://doi.org/10.1111/j.1399-3054.1992.tb04691.x>
 23. Chen X, Yuan H, Chen R, Zhu L, Du B, Weng Q, He G. Isolation and characterization of triacontanol-regulated genes in rice (*Oryza sativa* L.): Possible role of triacontanol as a plant growth stimulator. *Plant Cell Physiol.* 2002;43(8):869-76. <https://doi.org/10.1093/pcp/pcf100>
 24. Chen X, Yuan H, Chen R, Zhu L, He G. Biochemical and photochemical changes in response to triacontanol in rice (*Oryza sativa* L.). *Plant Growth Regul.* 2003; 40:249-56. <https://doi.org/10.1023/A:1025039027270>
 25. Aftab T, Khan MMA, Idrees M, Naeem M, Singh M, Ram M. Stimulation of crop productivity, photosynthesis and artemisinin production in *Artemisia annua* L. by triacontanol and gibberellic acid application. *J Plant Interact.* 2010;5(4):273-81. <https://doi.org/10.1080/17429141003647137>
 26. Idrees M, Khan MMA, Aftab T, Naeem M. Synergistic effects of gibberellic acid and triacontanol on growth, physiology, enzyme activities and essential oil content of *Coriandrum sativum* L. *Asian Aust J Plant Sci Biotechnol.* 2010;4(1):24-29.
 27. Naeem M, Khan MMA, Moinuddin. Triacontanol: a potent plant growth regulator in agriculture. *J Plant Interact.* 2012;7(2):129-42. <https://doi.org/10.1080/17429145.2011.619281>
 28. Islam S, Zaid A, Mohammad F. Role of triacontanol in counteracting the ill effects of salinity in plants: A review. *J Plant Growth Regul.* 2020; 40:1-20. <https://doi.org/10.1007/s00344-020-10064-w>
 29. Khan MMA, Bhardwaj G, Naeem M, Moinuddin, Mohammad F, Singh M, Nasir S, Idrees M. Response of tomato (*Lycopersicon esculentum* Mill.) to application of potassium and triacontanol. *Acta Hort.* 2009; 823:199-208. <https://doi.org/10.17660/ActaHortic.2009.823.29>
 30. Hu A, Lin B, Ren D, Chen J. Abscission of citrus fruits. I. The effects of triacontanol on cellulose and pectinase activities in the abscission of young fruit explants of Bendizao mandarin (*Citrus succosa* Hort. Ex Tanaka). *Acta Hortic.* 1985; 12:77-82.
 31. Shukla A, Farooqi AA, Shukla YN, Sharma S. Effect of triacontanol and chlormequat on growth, plant hormones and artemisinin yield in *Artemisia annua* L. *Plant Growth Regul.* 1992; 11:165-71. <https://doi.org/10.1007/BF00024071>
 32. Chaudhary BR, Sharma MD, Shakya SM, Gautam DM. Effect of plant growth regulators on growth, yield and quality of chilli (*Capsicum annuum* L.) at Rampur, Chitwan. *J Inst Agric Anim Sci.* 2006; 27:65-8. <https://doi.org/10.3126/jiaas.v27i0.697>
 33. El-Beltagi HS, Ismail SA, Ibrahim NM, Shehata WF, Alkhateeb AA, Ghazzawy HS, El-Mogy MM, Sayed EG. Unravelling the effect of triacontanol in combating drought stress by improving growth, productivity and physiological performance in strawberry plants. *Plants.* 2022; 11:1913. <https://doi.org/10.3390/plants11151913>
 34. Verma T, Bhardwaj S, Singh J, Kapoor D, Prasad R. Triacontanol as a versatile plant growth regulator in overcoming negative effects of salt stress. *J Ag Food Res.* 2022; 10:100351. <https://doi.org/10.1016/j.jafr.2022.100351>
 35. Momin SK, Gaikwad SS, Patel RJ, Amarcholi JJ, Sharma KM. Effect of foliar application of chemicals on fruiting parameters of mango (*Mangifera indica* L.) cv. Kesar. *Res J Agric Sci.* 2016;7(1):143-44.
 36. Sharma N, Singh K, Thakur A. Growth, fruit set, yield and fruit quality of olives (*Olea europaea* L.) as influenced by nutrients and bio-stimulants under rainfed condition. *Acta Hortic.* 2011; 890:385-92. <https://doi.org/10.17660/ActaHortic.2011.890.55>
 37. Abubakar AR, Ashraf N, Ashraf M. Effect of plant biostimulants on flowering, fruit drop, yield and return bloom of pomegranate cv. Kandhari Kabuli. *Asian J Hort.* 2012;7(2):473-77.
 38. Choudhary HD, Jain MC, Sharma MK, Bhatnagar P. Effect of plant growth regulators on growth and yield of nagpur mandarin (*Citrus reticulata* Blanc). *Asian J Hort.* 2013;8(2):746-50.
 39. Muhi HM, Abd AM. Effect of foliar spray with triacontanol, boron and low biuret urea on flowering and fruiting of apple trees cv. Sharaby. *Biochem. Cell Arch.* 2022;22(1):2963-68. <https://connectjournals.com/03896.2022.22.2963>
 40. Paralkar PD, Patil SG, Naglot UM. Effect of chemicals on fruit retention of mango (*Mangifera indica* L.) cv. Kesar. *Int J Chem Stud.* 2020;8(2):2951-54. <https://doi.org/10.22271/chemi.2020.v8.i2as.9199>
 41. Mahorkar KD, Naglot UM, Yadav SV, Hake PD. Effect of plant growth regulators on yield and quality of custard apple (*Annona squamosa* L.). *Int J Curr Microbiol App Sci.* 2020;9(3):587-93. <https://doi.org/10.20546/ijcmas.2020.903.069>
 42. Patel MK, Panda C, Senapati S. Influence of new generation PGRs on physical parameter of mango (*Mangifera indica* L.) cv. Dashehari. *Pharma Innov J* 2021;10(10):299-302.
 43. Hung NQ, Ha LTM, Hieu NQ, Tu PTT, Lam, VP. Gibberellin (GA₃) and copper sulphate pentahydrate (CuSO₄.5H₂O) reduce seeds per fruit and increase fruit quality in Bac Son mandarin fruit. *Seeds.* 2023;2(3):318-30. <https://doi.org/10.3390/seeds2030024>
 44. Lal N, Das R.P. Effect of growth regulators on yield and quality of guava (*Psidium guajava* L.) cv. Allahabad Safeda. *Int J Curr Microbiol App Sci.* 2017;6(5):857-63. <https://doi.org/10.20546/ijcmas.2017.605.096>
 45. Akram M. Effects of triacontanol on some qualitative and quantitative traits in kiwifruit. *Biol Forum.* 2018;10(1):96-100.
 46. Gaikwad GB, Nainwad RV, Hinge AM, Khote AV, Girase LA, Wadhekar NB. Effect of plant growth regulators and chemicals on quality of mango (*Mangifera indica* L.) var. Kesar. *Pharma Innov J.* 2022;11(9):2892-93.
 47. Dabhi DM, Patel MJ, Chaudhary HL. Effect of novel organic liquid fertilizer and plant growth regulators on quality of guava cv. Lucknow 49. *Biol Forum.* 2023;15(9):292-96.
 48. Iqbal N, Nazar R, Khan MIR, Masood A, Khan NA. Role of gibberellins in regulation of source- sink relations under optimal and limiting environmental conditions. *Curr Sci.* 2011; 100(7):998-1007.
 49. Gangadhar, Chaurasiya R, Sharma A, Tiwari S, Goyal G, Bhadauria A S, Singh AP, Yadav A. Effect of foliar application of GA₃ with and without NAA on fruit drop, growth, yield and quality of ber (*Zizyphus mauritiana* Lamk.) cv. Banarasi Karaka. *Int J Curr Microbiol App Sci.* 2019;Special issue-8: 45-56.

50. Bisen A. Response of growth promoting substances on flowering, fruiting and yield behavior of ber (*Zizyphus mauritiana* Lamk) cv. Apple ber. *Pharma Innov J.* 2022; 11(11): 2161-67.
51. Kumar A, Rajan R, Pandey K, Ramprasad RR, Kaur G, Vamshi T, Singh T. Impact of new generation plant growth regulators on fruit crops - A review. *Hortic Sci.* 2024;51(1):1-22. <https://doi.org/10.17221/166/2022-HORTSCI>
52. Islam S, Mohammad F. Triacantanol as a dynamic growth regulator for plants under diverse environmental conditions. *Physiol Mol Biol Plants.* 2020;26(5):871-83. <https://doi.org/10.1007/s12298-020-00815-0>