

RESEARCH ARTICLE

Laboratory aquaculture of *Lecane inermis* (Bryce, 1842) isolated from the Hilla River in Babylon province-Iraq

Wameedh AK AL-Yasari* & Adi Jassim Abd Al-Rezzaq

Department of Biology, College of Science, University of Babylon, Babylon 00 964, Iraq

*Email: sci.wameedh.adil@uobabylon.edu.iq

 OPEN ACCESS

ARTICLE HISTORY

Received: 24 November 2024

Accepted: 02 February 2025

Available online

Version 1.0 : 10 April 2025

Version 2.0 : 23 April 2025



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Wameedh AKAY, Adi Jassim AAR. Laboratory aquaculture of *Lecane inermis* (Bryce, 1842) isolated from the Hilla River in Babylon province-Iraq. Plant Science Today. 2025; 12 (2): 1-5. <https://doi.org/10.14719/pst.6305>

Abstract

The current study aimed to evaluate the environmental conditions necessary for cultivating the rotifer species (*Lecane inermis*) initially recorded in the Hilla River and their impact on its density. Using a zooplankton net, rotifer samples were collected in the early morning of winter 2024 from the Hilla River in Al-Mahawil district, Babylon province. The physical and chemical properties of the site were analyzed, including air temperature (6 °C), water temperature (9.5 °C), electrical conductivity (EC) (1090 µs/cm), total dissolved solids (TDS) (772 ppm), salinity (636 ppm), pH (8.6), dissolved oxygen (DO) (12.3 ppm) and turbidity (11.5 NTU). The cultured species *L. inermis* was isolated, identified and subsequently cultivated. Various factors, including temperature, salinity, dissolved oxygen and light versus dark, were then investigated to determine the effect on species density. The findings demonstrate that *L. inermis* exhibited higher density at low temperatures and revealed broad salinity tolerance, though it could only withstand lower salinity concentrations at high temperatures. Additionally, its density was greater under dark conditions than under light conditions. The highest density recorded was 585 ± 20 ind/L during the winter season.

Keywords

aquaculture; Hilla river; *Lecane inermis*; zooplankton density

Introduction

Zooplanktons are essential to the survival of a vast range of freshwater species, including fish. Gut content studies reveal that zooplankton constitutes a significant portion of the staple diet for several fish species in their natural habitats. Recognizing their importance in commercial aquaculture, several zooplankton species have been introduced and artificially cultivated. Zooplankton are considered excellent resources for live feed because of their high lipid and protein contents, which are crucial for the development and survival of fish larvae (1, 2). The availability of farmed zooplankton often facilitates a large culture of fish larvae. Even though commercially farmed fish species benefit significantly from zooplankton, their culture, and sufficient production must be financially viable. Regardless of the fish species, zooplankton culture is promoted because, in most cases, fish larvae obtain their entire or at least a significant portion of their nutrition from zooplankton food sources (3). Recent advances in zooplankton culture techniques have provided viable options for both small- and large-scale production. This study aims to evaluate the potential growth and survival of one zooplankton species (*L. inermis*) under artificial laboratory conditions, considering its nutritional benefits and ease of cultivation as it is known.

Aquaculture is one of the fastest-growing industries today. Fish require live food to survive. Although live food can be expensive, it is available in marketplaces. Therefore, culturing zooplankton is a cost-effective option for reducing production costs (4). Today, wastewater treatment facilities face significant problems due to the overgrowth of bacteria. *L. inermis* can effectively control the excessive growth of various bacteria (5). *L. inermis* is one of the most effective zooplankton species for treating wastewater. They are highly prolific rotifers, with populations dominated by females that lay an average of 20 eggs during their 9 days of lifespan. Successful growth of *Lecane* species depends upon precise feeding. Algae and bacteria serve as their primary source of food. *L. inermis* has already been identified as a possible control agent (6).

Many recent studies have highlighted the industrial importance of gradually replacing common live feed, such as brine shrimp and rotifer, with different zooplankton species, including Cladocera's like water fleas, either entirely or partially (7). Environmental factors play an important role in controlling zooplankton. Fish mostly regulate "top-down" processes for larger species, whereas "bottom-up" mechanisms primarily govern small species like rotifers and small Cladocera's (8).

Materials and Methods

Collection of samples

Water and zooplankton samples were collected in February 2024 from the Hilla River in Al-Mahaweel district in Babylon province. Zooplankton sampling was conducted in the early morning to minimize heat stress, using a zooplankton net with 50 µm mesh size. Samples were collected from depths up to 2 m from the water's surface and then emptied into a zooplankton net to pass through 70 L of water. The net was then cleaned externally with a sprayer and the samples were transferred to the laboratory for isolation, diagnosis and culture (9). Water samples were collected in 5 L polyethylene containers for physical and chemical analysis.

Physical and chemical parameters

Air and water temperatures were measured using a thermometer. Electrical conductivity, pH, TDS and salinity were measured using a HANNA multi-parameters device, DO was measured using a DO meter and Turbidity NTU (Nephelometric Turbidity Unit) was measured using a Turbidimeter manufactured by the Lovibond company.

Enumeration of the zooplankton

To count the zooplankton per liter, the volume of water filtered through the net was calculated using the cylinder volume formula (10). Moreover, the method described by APHA was employed to calculate the number of individuals per liter, with all the results expressed as the number of individuals/liters (ind/L) (11). A Sedgewick-Rafter slide chamber was used for zooplankton counting.

Identification of rotifer *Lecane inermis*

For taxonomic identification, a live sample was placed on the slide, observed and then fixed using a formaldehyde solution

of 4%. The *Lecane* specimens were examined under a light microscope and scanning electron microscope. The specimen was identified as *L. inermis*, a rotifer based on its size and morphological characteristics. Its identification was performed using the standard taxonomic keys (12-16).

The culture of algae and rotifer (*Lecane inermis*)

The culture of algae: The manufacturer (Reef Nutrition) provided a pure culture of *Nannochloropsis* sp., which was cultivated using the batch-culture technique. The cultivation conditions for the algae were established as per standard protocol (17). For rotifer growth under optimal laboratory conditions, a synthetic medium containing 96 mg NaHCO₃, 60 mg CaSO₄·2H₂O, 60 mg MgSO₄ and 4 mg KCl in 1 L of sterile deionized water was utilized and the right amount of algae was fed (17). The algae were harvested during the exponential phase, centrifuged at 2000 rpm for 8 min and concentrated to 1×10⁸ cells/mL at 4 °C in a refrigerator.

The rotifer stock culture: The samples were screened in the laboratory through a 500 µm mesh net to remove fish and decapod larvae. Then, they were filtered through a 120 µm mesh to remove copepods, barnacles and nauplii. A Pasteur pipette and capillary tubes separated *L. inermis* from the remaining zooplankton to create a stock culture using a batch culture approach, beginning with a 25 mL conical flask. During cultivation, a photoperiod of 12 h of light and 12 h of darkness was maintained. The *Nannochloropsis* sp. was fed to the experimental rotifer (18). Aeration was necessary to prevent algae aggregation and enhance its rotifers accessibility (19). Algal cells were then supplied to the culture containers to maintain the target cell concentrations.

Results and Discussion

Table 1 shows the values of the physicochemical parameters during the sampling time. The air temperature was 6 °C and the water temperature was 9.5 °C, variations in air temperature reflected the general attitude of the geographical location and water temperature followed a similar trend, indicating a close relationship between the 2 (20, 21). This study's findings were consistent with those of previous Iraqi environmental studies. The current study observed a narrow pH value (8.6) attributed to the buffering capacity of carbonic acid and bicarbonate, which resist significant pH fluctuations (22). Additionally, numerous investigations have noticed a small pH range (15, 16, 23).

Dissolved salts influence EC in water, while salinity represents the total ion concentrations in the water. Hence, there is a strong correlation between EC, TDS and salinity.

Table 1. Physico-chemical parameters measured at the study site during sampling

Parameters	Mean value
Air temperature (°C)	6
Water temperature (°C)	9.5
pH	8.6
Electrical conductivity (EC) (µs/cm)	1090
Total Dissolved Solids (TDS) (mg/L)	772
Salinity (mg/L)	636
Dissolved Oxygen (DO) (mg/L)	12.3
Turbidity (NTU)	11.5

This study's EC, TDS and salinity values were 1090 $\mu\text{S}/\text{cm}$, 772 ppm and 636 ppm, respectively. DO is a critical factor for the survival of aquatic life (24). Dissolved oxygen levels were favorable during winter (12.3 ppm). This may be due to a decline in the ratios of consumption and decomposition associated with an increase in gas solubility. This might have changed due to rising temperatures, making gasses less water-soluble (25). Finally, the average density of rotifers recorded at the study station was 23.4 individuals per liter during the sampling time. Environmental factors also impact rotifer growth, but they are not independent of one another. For instance, a change in temperature influences other environmental factors like salinity and dissolved oxygen, which in turn impacts rotifer growth and density. Generally, the most crucial factor affecting the change is temperature.

Description of *Lecane inermis*

In the present study, one species of rotifers (*L. inermis*) belonging to the family Lecanidae was recorded for the first time in the Hilla River. The diagnostic features include the length of lorica (60-80 μm), width of lorica (36-48 μm), length of ventral plate (60-62 μm), width of ventral plate (39-40 μm), length of toe with claw (30-44 μm), length of claw only (10-12 μm), length of trophi (20 μm), width of trophi (25 μm), length of fulcrum (7.06 μm), length of rami (5.88 μm), length of incus (7.53 μm) and length of manubria (19.18-20.82 μm) as clear in Fig. 1-4.

Laboratory conditions of *L. inermis* aquaculture

The basic environmental factors recorded in the current study (Table 2) are crucial for the culture of *L. inermis*. The water temperature ranged from 7 $^{\circ}\text{C}$ to 26 $^{\circ}\text{C}$ during winter

and summer, respectively, while the air temperature ranged from 10 $^{\circ}\text{C}$ to 30 $^{\circ}\text{C}$. The difference in temperature between the water and air is because the water's high specific heat helps it retain its temperature and protect aquatic life from sudden temperature fluctuations (20, 21).

Temperature significantly influences zooplankton's distribution and determines the water's oxygen level (26). Numerous studies have demonstrated that rotifers population densities typically increase during the summer and decrease in the winter (27-30). However, the current study indicated that *L. inermis* reached its highest density (585 ± 20 ind/L) in the winter (Table 2). The species also exhibited a broad tolerance range for EC, salinity and TDS, with its highest density recorded at an elevated salinity level (Table 2).

A study reported that most rotifers require more than 1 ppm DO concentration (31). While some can survive briefly in anaerobic conditions or close to them, others can live for extended periods in low-oxygen environments. In our current study, *L. inermis* achieved its highest density at a DO level of 7.8 ppm (Table 2). However, this species was first observed at an oxygen concentration of less than 4 ppm. Research suggests that *L. inermis* may reproduce periodically in aerobic and anaerobic environments, tolerating DO concentrations as low as 1 ppm (32).

Zooplankton nutrition is influenced by light and darkness. Vertical migration plays a crucial role, with zooplankton ascending to the water's surface at night to feed and retreating to deeper layers during the day to evade predators (33). This behavior aligns with our findings, where *L. inermis* cultivated in the dark exhibited a higher density than those exposed to 200 Lux (Fig.5). The increased density in dark-cultured conditions may be attributed to more effective feeding than in light-cultured conditions.




Fig. 1. Trophi of *L. inermis* observed under Scanning Electron Microscope (SEM).




Fig. 3. Diagrammatic illustration of *L. inermis* (a - Front view, b - Lateral




Fig. 2. *L. inermis* observed under a light microscope (a-front view, b-lateral view).





Fig. 4. Toe and claw of *L. inermis* observed under the light microscope.

Table 2. Laboratory conditions of *L. inermis* aquaculture

Parameters	Winter mean ± SD	Summer mean ± SD	Sig.
Water Temperature °C	7 ± 1	26 ± 2	0.00
Air Temperature °C	10 ± 1	30 ± 2	0.00
EC (mS)	9.6 ± 0.2	4 ± 0.1	0.00
Salinity ppt	6.1 ± 0.1	2.5 ± 0.1	0.00
TDS ppt	6.8 ± 0.1	2.4 ± 0.1	0.00
Do mg/L	7.8 ± 0.05	7.6 ± 0.1	0.055
pH	9.2 ± 0.01	9 ± 0.01	NS
Growth density (ind/L)	585 ± 20	137 ± 17	0.00

**Fig. 5.** Density of *L. inermis* in light and dark conditions.

Conclusion

The results of the current study demonstrate that *L. inermis* can be successfully cultivated in a high-density laboratory environment. The species exhibits a broad tolerance for salinity at low temperatures but is limited to lower salt concentrations at higher temperatures. Furthermore, its density is significantly higher in dark than light conditions.

Acknowledgements

The authors express their gratitude for the support from the Department of Biology, College of Science at the University of Babylon/ Iraq.

Authors' contributions

WAKAY and AJAAR designed and performed the experiment. AYWAK analyzed data and wrote the first draft of the manuscript; AARAJ performed statistical analysis and revised the manuscript. Both authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None

References

- Ogello EO, Wullur S, Yoshitaka S, Hagiwara A. Dietary value of waste-fed rotifer *Brachionus rotundiformis* on the larval rearing of Japanese

whiting *Sillago japonica*. E3S Web of Conference. 2020;147(4):01005. <https://doi.org/10.1051/e3sconf/202014701005>

- Lomartire S, Marques JC, Goncalves AMM. The key role of zooplankton in ecosystem services: A perspective of interaction between zooplankton and fish recruitment. Ecol Indic. 2021;129:107867. <https://doi.org/10.1016/j.ecolind.2021.107867>
- Kar S, Das P, Das U, Bimola M, Kar D, Aditya G. Culture of zooplankton as fish food: observations on three freshwater species from Assam, India. Aquac Aquar Conserv Legis. 2017;10(5):1210–20.
- Cotonnec G, Brunet C, Sautour B, Thoumelin G. Nutritive value and selection of food particles by copepods during a spring bloom of *Phaeocystis* sp. in the English Channel, as determined by pigment and fatty acid analyses. J Plankton Res. 2001;23(7):693–703. <https://doi.org/10.1093/plankt/23.7.693>
- Drzewicki A, Kowalska E, Pajdak-Stós A, Fiałkowska E, Kocerba-Soroka W, Sobczyk Ł, et al. Experimental attempt at using *Lecane inermis* rotifers to control filamentous bacteria Eikelboom type 0092 in activated sludge. Water Environ Res. 2015;87(3):205–10. <https://doi.org/10.2175/106143015x14212658613037>
- Attar MM, Patil P, Joshi R, Patil V. Laboratory culture of selected freshwater zooplankton species. J Adv Zool. 2024;45(3):506–20. <https://doi.org/10.53555/jaz.v45i3.4360>
- Samat NA, Yusoff FM, Rasdi NW, Karim M. The efficacy of *Moina micrura* enriched with probiotic *Bacillus pocheonensis* in enhancing survival and disease resistance of red hybrid Tilapia (*Oreochromis* spp.) Larvae. Antibiotics. 2021;10(8):989. <https://doi.org/10.3390/antibiotics10080989>
- Rasdi N, Suhaimi H, Yuslan A, Sung YY, Ikhwanuddin M, Omar SS, et al. Effect of mono and binary diets on growth and reproduction of cyclopoid copepod. Aquac Aquar Conserv Legis. 2018;11(5):1658–71.
- Harris RP, Wiebe PH, Lenz J, Skjoldal HR, Huntley M. Zooplankton methodology manual. London: Academic Press; 2000. 684 p.
- Lind O. A handbook of limnological methods. St. Louis: C.V. Mosby Co.; 1979. 199 p.
- APHA. Standard methods for the examination of water and wastewater. 21st ed. Washington, DC: American Public Health Association, American Water Works Association, Water Environment Federation; 2005
- Davidson TA, Wallace RL, Snell TW, Ricci C, Nogrady T. Rotifera: Volume 1-Biology, Ecology and Systematics. 2nd ed. J Paleolimnol. 2012;47:159. <https://doi.org/10.1007/s10933-011-9539-4>
- Segers H. Annotated checklist of the rotifers (Phylum Rotifera), with notes on nomenclature, taxo-nomy and distribution. Zootaxa. 2007;1564(1):1–104. <https://doi.org/10.11646/zootaxa.1564.1.1>
- Jersabek CD, Leitner MF, editors. Rotifer World Catalogue [Internet]. Catalogue of Life. [cited 2025 Feb 4]. Available from: <https://www.catalogueoflife.org/data/dataset/298081>. <https://doi.org/10.48580/dg9ld-g8gp>
- Abd Al-Rezzaq AJ. A diagnostic and ecological study of the planktonic species of *Rotifera* in the Hilla River, Iraq [thesis]. Babylon, Iraq: University of Babylon; 2014. 218 p.
- Al-Yasari WAK. Biodiversity of Rotifera and Cladocera in Al-Hilla River, Babylon Province, Iraq [thesis]. Babylon, Iraq: University of Babylon; 2023. 264 p.
- Lavens P, Sorgeloos P, editors. Manual on the production and use of live food for aquaculture. FAO Fisheries Technical Paper No. 361. Laboratory of Aquaculture and Artemia Reference Center, University of Gent, Belgium; 1996
- El-Tohamy WS, Abou Eleila RHM. The impacts of different algal diets, temperature and salinity on locally isolated euryhaline rotifer's (*Brachionus plicatilis*) growth and egg production. Egypt J Aquat Biol Fish. 2024;28(3):911–30. <https://doi.org/10.21608/ejabf.2024.361011>

19. Rahman AR, Cob ZC, Jamari Z, Mohamed AM, Toda T, Ross OH. The effects of microalgae as live food for *Brachionus plicatilis* (rotifer) in intensive culture system. Trop Life Sci Res. 2018;29(1):127. <https://doi.org/10.21315/tlsr2018.29.1.9>
20. Tiwari S, Dixit S, Gupta SK. An evaluation of various physico-chemical parameters in surface waters of Shahpura lake. Bhopal Pollut Res. 2004;23:829–32.
21. Manickam N, Bhavan PS, Santhanam P, Muralisankar T, Srinivasan V, Radhakrishnan S, et al. Seasonal variations of zooplankton diversity in a perennial reservoir at Thoppaiyar, Dharmapuri district, South India. Austin J Aquac Mar Biol. 2014;1(1):1–7.
22. Wetzel RG. Limnology: Lake and river ecosystems. 3rd ed. San Diego: Academic Press; 2001. 1006 p.
23. Al-Fanhrawi AA. Distribution and diversity of large benthic invertebrates in sediments of Shatt Al-Hilla, Iraq [thesis]. Babylon, Iraq: University of Babylon; 2010.
24. Bulbul A, Anushka, Mishra A. Effects of dissolved oxygen concentration on freshwater fish: A review. Int J Fish Aquat Stud. 2022;10(4):113–27. <https://doi.org/10.22271/fish.2022.v10.i4b.2693>
25. Stevens MR. Water quality and trend analysis of Colorado-Big Thompson system reservoirs and related conveyances, 1969 through 2000 [Internet]. Water-Resources Investigations Report 2003-4044. United States Geological Survey. 2003. <https://doi.org/10.3133/wri034044>
26. Chen G. Effects of physical and chemical factors on zooplankton in tropical shallow urban lakes. In: 2nd International Conference on Advances in Civil Engineering, Energy Resources and Environment Engineering; 2020 May 22–24; Nanning, China. IOP Conf Ser Earth Environ Sci; 2020. 526:012028. <https://doi.org/10.1088/1755-1315/526/1/012028>
27. Arora J, Mehra NK. Seasonal dynamics of the rotifers in relation to physical and chemical conditions of the river Yamuna (Delhi), India. Hydrobiologia. 2003;491:101–09. <https://doi.org/10.1023/A:1024490805310>
28. Chittapun S, Pholpunthin P, Segers H. Diversity of rotifer fauna from five coastal peat swamps on Phuket Island, Southern Thailand. Sci Asia. 2007;33:383–87.
29. Bhat NA, Wanganeo A, Rania R. The composition and diversity of net zooplankton species in a tropical water body (Bhoj Wetland) of Bhopal, India. Int J Biodivers Conserv. 2014;6(5):373–81. <https://doi.org/10.5897/IJBC2014.0702>
30. Roman MR, Brandt SB, Houde ED, Pierson JJ. Interactive effects of hypoxia and temperature on coastal pelagic zooplankton and fish. Front Mar Sci. 2019;6:139. <https://doi.org/10.3389/fmars.2019.00139>
31. Wallace RL, Snell TW. Phylum Rotifera. In: Thorp JH, Covich AP, editors. Ecology and classification of North American freshwater invertebrates. 2nd ed. San Diego: Academic Press; 2001. p. 195–254. <https://doi.org/10.1016/B978-012690647-9/50009-0>
32. Kocerba-Soroka W, Fialkowska E, Pajdak-Stos A, Sobczyk M, Plawecka M, Fyda J. Effect of the rotifer *Lecane inermis*, a potential sludge bulking control agent, on process parameters in a laboratory-scale SBR system. Water Sci Technol. 2013;68(9):2012–18. <https://doi.org/10.2166/wst.2013.453>
33. Melo TX, Dias JD, Simões NR, Bonecker CC. Effects of nutrient enrichment on primary and secondary productivity in a subtropical floodplain system: an experimental approach. Hydrobiologia. 2019;827:171–81. <https://doi.org/10.1007/s10750-018-3763-0>