



RESEARCH ARTICLE

Melatonin application extends banana shelf life by delayed ripening

Madhavan Devi¹, Prabhakaran Jeyakumar^{1*}, Veerasamy Ravichandran¹, Loganathan Arul², Murugesan Balakrishnan³ & Mathiyazhagan Kavino⁴

¹Department of Crop Physiology, Tamil Nadu Agricultural University, Coimbatore, 641 003 Tamil Nadu, India

²Department of Plant Biotechnology, Tamil Nadu Agricultural University, Coimbatore, 641 003 Tamil Nadu, India

³Department of Food Process Engineering, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

⁴Department of Fruit Science, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

*Email: jeyakumar@tnau.ac.in

 OPEN ACCESS

ARTICLE HISTORY

Received: 23 December 2024

Accepted: 21 January 2025

Available online

Version 1.0 : 02 March 2025

Version 2.0 : 18 March 2025



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Devi M, Jeyakumar P, Ravichandran V, Arul L, Balakrishnan M, Kavino M. Melatonin application extends banana shelf life by delayed ripening. *Plant Science Today*. 2025; 12(1): 1-12. <https://doi.org/10.14719/pst.6867>

Abstract

Bananas are among the worlds' most economically significant fruit crops, valued for their distinctive flavour and aroma, despite being highly susceptible to rapid postharvest physiological deterioration. Melatonin, a non-toxic bioactive compound, regulates several plant physiological processes and is widely studied in postharvest management. With this background, the study investigated the effect of different concentrations of melatonin on three banana varieties, namely Grand Naine.(AAA), Poovan (AAB) and Ney Poovan (AB). The fruits were stored under ambient (AS) ($27 \pm 2^\circ\text{C}$) and cold storage (CS) (17°C) with 90-95% relative humidity. Melatonin treatment (1000 μM) effectively reduced weight loss, retained firmness and total soluble solids, reduced ethylene production, enhanced chlorophyll retention and delayed starch degradation. Grand Naine showed the most pronounced effects, followed by Ney Poovan and Poovan. Overall, 1000 μM melatonin was the most effective treatment for extending banana shelf life and maintaining quality under ambient and cold storage conditions.

Keywords

banana; melatonin; ripening; shelf life; storage conditions

Introduction

Melatonin (N-acetyl-5-methoxytryptamine) is a naturally occurring indoleamine with diverse roles in plant physiological processes (1). Melatonin synthesis starts from the amino acid tryptophan and is produced naturally in various fruit tissues such as peel, pulp and seeds (2, 3). Melatonin is a non-toxic compound involved in multiple physiological processes, including seed and root development, plant growth and flowering (4, 5). It also influences fruit development, ripening, senescence and responses to biotic and abiotic stresses (6-9). Moreover, melatonin is widely studied in postharvest applications as it has efficacy in prolonging storage life and enhancing the quality of fruits and vegetables (10).

Notably, the exogenous application of melatonin delays ethylene production during the ripening process by inhibiting the expression of 1-aminocyclopropane-1-carboxylic acid synthase (ACS) and 1-aminocyclopropane-1-carboxylic acid oxidase (ACO) enzyme activities (11). Melatonin helps maintain several fruit quality attributes such as weight loss, respiration rate, titrable acidity, fruit firmness, soluble sugars, organic acids, carotenoid content and aromatic compounds (12, 13). Studies have shown that exogenous melatonin

significantly delays fruit ripening in apple, sweet cherry, peach, pear and mango (14-18). It also reduces decay in fruits like kiwifruit(18), litchi(19) and strawberry(20).

Bananas are among tropical and subtropical regions' most widely cultivated and economically essential fruits (21). Bananas are a natural and instant energy source due to their carbohydrate content and important nutrients like vitamin C, B₆, potassium, iron, magnesium and dietary fibre, making them a healthy addition to the diet (22). In India, banana cultivation covers an area of 994.14 thousand hectares ('000 ha) with a production of 366.14 lakh tonnes and a productivity of 36.83 MT ha⁻¹ (23). The climacteric nature of bananas causes them to ripen rapidly after harvest, which shortens their postharvest shelf life. Postharvest handling and marketing chain involves significant challenges under various storage conditions, accelerating maturations physiological and metabolic processes and producing fruit spoilage. Postharvest losses of perishable commodities like fruits and vegetables can reach 20-40%, occurring at various stages such as harvesting, transportation, storage and marketing (24). Notably, delaying the ripening before consumption is a significant challenge for supply chain management in perishable fruits, including bananas (25, 26). During the climacteric phase, bananas change chlorophyll breakdown, cell wall softening and conversion of starch into sugars. These changes affect the fruits colour, firmness, aroma and shelf life (27).

Optimizing postharvest ripening is crucial for improving banana quality and shelf life, especially for market distribution. During postharvest ripening, there is a notable rise in ethylene production and respiration rate, which coincides with the fruit climacteric phase. Ethylene is an essential gaseous phytohormone linked to the ripening process. It orchestrates the timely expression of genes that govern various ripening and senescence events, including increased respiration, autocatalytic ethylene production and alterations in fruit colour, texture, aroma and flavour. Enzymes such as ACS and ACO play a key role in ethylene biosynthesis, which is triggered during postharvest ripening (21, 28, 29, 30). Ethylene production in climacteric fruits regulates ripening and quality-related processes, while ethylene inhibitors markedly postpone the ripening process (11). However, various postharvest management techniques aim to extend the shelf life of perishable fruits that effectively control ethylene production, maintaining the sensory and nutritional quality of the fruits (24).

Postharvest treatments vary according to regional agricultural practices, local regulations and market demands (25). Fresh fruits are highly perishable and deteriorate rapidly during storage, resulting in reduced taste, nutritional quality, poor marketability and acceptance and leading to economic loss (19). Optimizing quality preservation and extending the postharvest shelf life of fruit represent significant challenges in the fruit industry. Researchers are focused on various postharvest treatments to improve the shelf life of fruits and enhance their quality. Several postharvest technologies have been developed to maintain and extend the shelf life of fresh fruit by targeting ethylene oxidation, inhibiting ethylene action and modulating the ripening process (26).

Melatonin is gaining attention as a promising approach for improving fruit production and prolonging shelf life. Several studies reported that exogenous melatonin application delayed postharvest ripening in various horticultural produce such as apple, pear, peach, guava and pomegranate (14-16, 27, 28). Melatonin application has shown the potential to enhance disease resistance and reduce deterioration rates in several fruits during storage, including litchi, strawberry and peach (15, 20, 29). However, the use of melatonin in the postharvest management of bananas is limited. Hence, this study aims to elucidate the effect of postharvest application of melatonin on three banana varieties under different storage conditions (ambient and cold). The study aimed to optimize melatonin concentration to extend the shelf life of bananas by modifying morphological, physio-chemical and quality traits.

Materials and Methods

Source of banana and melatonin

Freshly harvested bunches of three banana varieties, Grand Naine (AAA) (V₁), Poovan (AAB) (V₂) and Ney Poovan (AB) (V₃) were sourced from a farmers' field at Thadagam road, Coimbatore (11° .54'N, 76° .56'E). The varieties above are widely cultivated and well-known for their high yield and uniform fruit quality. Moreover, the ripening behaviour of these varieties differs slightly, making it valuable for assessing the broad applicability of melatonin treatments. Uniformly sized, healthy banana fruits at the mature green stage (approximately 75 % maturity) were selected for the study. Melatonin (99 % purity) was purchased from Sigma-Aldrich Pvt. Ltd and used for the experiment.

Treatment details

Banana hands were washed with running tap water, immersed in 100 ppm of sodium hypochloride (NaClO) for 2 min and air dried at room temperature. Sodium hypochlorite acts as a disinfectant in the postharvest storage of bananas, preventing microbial decay during storage. Banana hands were tagged and dipped for 3 min in sterile water (control) and different melatonin concentrations with a non-ionic surfactant of 0.05% Tween 20 to reduce the surface tension. The treatment, T₁ - Control (untreated), T₂ - Melatonin at 200 µM, T₃ - Melatonin at 400 µM, T₄ - Melatonin at 600 µM, T₅ - Melatonin at 800 µM and T₆ - Melatonin at 1000 µM. The control and treated fruits were kept in plastic crates with a size of 54 × 35 × 28 cm and stored under ambient (27 ± 2°C) and cold (17 °C) storage conditions, one set at each. Morpho-physiological, biochemical and quality characteristics were measured at 2-day intervals during storage. The ripening behaviour of the banana was assessed daily and also take the observations on alternate days till the 15th day after treatment. Each replication involved two hands of banana. One hand of banana was used to measure the hand weight and shelf-life longevity, while the other was used to analyze other morpho-physiological, biochemical and quality parameters.

Fruit morphological characteristics

At first, the banana hand, *i.e.*, a cluster of bananas from the same bunch, was selected for each treatment. The

hands were separated from the bunches, weighed individually using an electronic balance and expressed in kg hand⁻¹. Fruit firmness was assessed using a penetrometer (TA-XT2 Texture Analyzer, Stable Micro System, UK). The test was conducted on whole bananas at three points: proximal, distal and mid-portion. The maximum force recorded at these points measured firmness, with values expressed in Newton (N).

Fruit physio-chemical and quality characteristics

Ethylene production was measured using a portable analyzer (Bioconservacion, Spain). Bananas were sealed in containers and the ethylene concentration in parts per million (ppm) was determined directly. The rate of ethylene production was expressed as $\mu\text{L kg}^{-1} \text{h}^{-1}$.

Total soluble solids (TSS) were determined from the clear juice filtrate using a handheld refractometer, with results expressed in °Brix. Chlorophyll content was quantified using the DMSO (Dimethyl sulfoxide) extraction method described by Hiscox and Israelstam (30). About 0.5 g of banana peel was subjected to DMSO extraction by incubating it in darkness for 12 h. Then, the volume was made up to 10 mL with DMSO. Total chlorophyll content was determined by measuring absorbance at 652 nm using a UV-1900i spectrophotometer (Shimadzu, UK) and expressed as $\mu\text{g g}^{-1}$ fresh tissue and calculated using the following Equation 1.

$$\text{Total chlorophyll} = \frac{\text{O.D. Value at 652}}{34.5} \times \frac{V}{1000} \quad (\text{Eqn. 1})$$

Where, V=Volume of the extract, W=Fresh weight in grams

The starch-iodine staining method was determined to assess starch conversion to sugars during the ripening process. A staining solution containing 1 % potassium iodide and 0.25 % iodine was prepared. Bananas were sliced transversely at the midpoint into 2-3 cm thick sections, including the peel. The cut surfaces of the banana sections were immersed in the starch-iodine solution for 5 seconds. The resulting starch patterns were analyzed by comparing them to a starch-iodine staining chart specific to bananas (31, 32).

Postharvest shelf life of fruit

The shelf life of banana fruit was determined by measuring the interval from the first day of storage to the end of the fruits' edible life, as observed visually after the applied treatment. Maturity indices for each fruit were assessed by comparing the peel colour to a standard colour chart and expressed in days (33).

Statistical analysis

The experimental design was under a Factorial Completely Randomized Design (FCRD). Data are expressed as mean \pm standard error of the mean. Mean comparisons were performed using the least significant difference (LSD) method and analysis of variance (ANOVA) with a substantial level of 0.05 was calculated using R software (version 4.3.1) for data analysis and bar diagrams were visualized using Microsoft Excel 2010.

Results and Discussion

Effect of postharvest melatonin treatment on hand weight

Significant differences ($p \leq 0.05$) in hand weight were observed between control and treated fruits under two different storage conditions. The results showed a gradual reduction in hand weight during storage. Under ambient conditions, the maximum weight loss was observed in the control (T_1), with 23.07 %, 32.71 % and 21.05 % recorded for V_1 , V_2 and V_3 respectively. In contrast, bananas treated with 1000 μM of melatonin (T_6) had the lowest weight loss with 21.07 %, 22.62 % and 15.74 % for V_1 , V_2 and V_3 respectively (Table 1). Under cold storage, fruits treated with melatonin at 1000 μM (T_6) depicted a minimum reduction of hand weight of 8.73, 15.78 and 7.34 % in V_1 , V_2 and V_3 compared to control (T_1) (12.25 V_1 , 22.46 V_2 and 8.52 % V_3) on 15th day of observation (Table 2). Postharvest application of melatonin at 1000 and 1500 μM significantly diminished weight loss across different banana varieties. In contrast, untreated fruits experienced more significant weight loss, likely due to elevated respiration and transpiration rates. Melatonin treatment postpones the climacteric peak, thus attenuating metabolic activity and reducing weight loss during ripening (34). Similar findings have shown that melatonin prevents weight loss in apple, orange, blueberry and nectarine compared to untreated fruits (14, 35-37).

Effect of postharvest melatonin treatment on firmness and total soluble solids

Statistically significant differences ($p \leq 0.05$) in fruit firmness were observed among the treatments, varieties and their interactions. During the ripening phase, fruit firmness progressively decreased under both storage conditions. Grand Naine treated with melatonin at 1000 μM maintained fruit firmness of 13.85 Newton (V_1T_6) than the untreated fruits of 5.65 Newton (V_1T_1) on the 9th day of ambient storage condition (Fig. 1). Under cold storage conditions, fruits treated with 1000 μM melatonin (T_6) showed a minor reduction in firmness compared to the control. By the 15th day, firmness values were 20.85 to 4.30 N for Grand Naine, 20.15 to 1.62 N for Poovan and 19.10 to 1.73 N for Ney Poovan (Fig. 2). Similar result was reported by Hu *et al.* (11), who found that melatonin-treated fruits showed retention of fruit firmness in bananas, it might be due to repression of ethylene production during postharvest banana ripening. Exogenous melatonin application lowers ethylene production and respiration rates, reducing CO_2 levels. This prevents the build-up of acetaldehyde and ethanol, thereby preserving fruit firmness and strengthening the fruit skin, enhancing postharvest quality (38, 39).

Total soluble solids (TSS) are crucial for assessing fruit maturity and nutrient richness, serving as a key indicator of fruit quality and influencing economic value in the fruit trade. Bananas treated with 1000 μM melatonin (T_6) showed lower TSS values (19.3, 24.8 and 24.7 °Brix for Grand Naine, Poovan and Ney Poovan, respectively) compared to the control (T_1 : 24.8, 28.5 and 28.8 °Brix) on the 9th day under ambient storage conditions (Fig. 3). At 15th day of cold storage, Grand Naine treated with 1000 μM of melatonin (V_1T_6) showed lower TSS values (4.2 to 22.0 °Brix) compared

Table 1. Effect of melatonin on hand weight (kg) in different banana varieties under ambient storage condition

Treatment details	Ambient storage (27±2°C)					
	1 st day	3 rd day	5 th day	7 th day	9 th day	Mean
V ₁ T ₁	3.172 ± 0.05 ^a	2.796 ± 0.04 ^a	2.666 ± 0.04 ^a	2.508 ± 0.04 ^a	2.440 ± 0.04 ^a	2.716
V ₁ T ₂	2.155 ± 0.03 ^d	1.903 ± 0.03 ^e	1.818 ± 0.03 ^e	1.719 ± 0.03 ^e	1.667 ± 0.03 ^d	1.834
V ₁ T ₃	2.442 ± 0.03 ^c	2.294 ± 0.03 ^c	2.200 ± 0.02 ^c	2.098 ± 0.02 ^c	2.044 ± 0.02 ^c	2.216
V ₁ T ₄	2.776 ± 0.04 ^b	2.456 ± 0.04 ^b	2.356 ± 0.03 ^b	2.236 ± 0.03 ^b	2.164 ± 0.03 ^b	2.398
V ₁ T ₅	2.416 ± 0.04 ^c	2.288 ± 0.03 ^c	2.192 ± 0.03 ^c	2.090 ± 0.03 ^c	2.006 ± 0.03 ^c	2.198
V ₁ T ₆	1.784 ± 0.03 ^{ef}	1.566 ± 0.03 ^h	1.504 ± 0.03 ^f	1.426 ± 0.03 ^f	1.408 ± 0.03 ^{ef}	1.538
V ₂ T ₁	2.152 ± 0.03 ^d	2.068 ± 0.03 ^d	1.938 ± 0.03 ^d	1.786 ± 0.03 ^d	1.448 ± 0.02 ^e	1.878
V ₂ T ₂	1.735 ± 0.03 ^f	1.610 ± 0.03 ^{gh}	1.260 ± 0.02 ^g	1.170 ± 0.02 ^g	1.052 ± 0.02 ^h	1.352
V ₂ T ₃	1.804 ± 0.02 ^{ef}	1.702 ± 0.02 ^f	1.564 ± 0.02 ^f	1.412 ± 0.02 ^f	1.182 ± 0.01 ^g	1.533
V ₂ T ₄	1.628 ± 0.02 ^g	1.432 ± 0.02 ⁱ	1.326 ± 0.02 ^g	1.210 ± 0.02 ^g	1.174 ± 0.02 ^g	1.354
V ₂ T ₅	1.830 ± 0.03 ^e	1.678 ± 0.03 ^{fg}	1.548 ± 0.02 ^f	1.432 ± 0.02 ^f	1.370 ± 0.02 ^f	1.572
V ₂ T ₆	1.786 ± 0.03 ^{ef}	1.666 ± 0.01 ^{fg}	1.550 ± 0.01 ^f	1.436 ± 0.03 ^f	1.382 ± 0.03 ^f	1.564
V ₃ T ₁	0.912 ± 0.01 ^{hi}	0.840 ± 0.01 ^k	0.798 ± 0.01 ⁱ	0.750 ± 0.01 ⁱ	0.720 ± 0.01 ^j	0.804
V ₃ T ₂	0.850 ± 0.01 ^{ij}	0.798 ± 0.01 ^{kl}	0.768 ± 0.01 ⁱ	0.687 ± 0.01 ^{jk}	0.661 ± 0.01 ^{kl}	0.745
V ₃ T ₃	0.966 ± 0.01 ^h	0.918 ± 0.01 ^j	0.888 ± 0.01 ^h	0.858 ± 0.01 ^h	0.828 ± 0.01 ⁱ	0.892
V ₃ T ₄	0.714 ± 0.01 ^k	0.662 ± 0.01 ⁿ	0.660 ± 0.01 ^j	0.630 ± 0.01 ^{kl}	0.584 ± 0.01 ^m	0.650
V ₃ T ₅	0.804 ± 0.01 ^j	0.738 ± 0.01 ^{lm}	0.734 ± 0.01 ⁱ	0.704 ± 0.01 ^{ij}	0.682 ± 0.01 ^{jk}	0.732
V ₃ T ₆	0.724 ± 0.01 ^k	0.676 ± 0.01 ^{mn}	0.648 ± 0.01 ^j	0.622 ± 0.01 ^l	0.610 ± 0.01 ^{lm}	0.656
Mean	1.703	1.561	1.468	1.376	1.301	
SE.d	0.039	0.035	0.033	0.031	0.030	
CD (0.05)	0.078	0.071	0.067	0.062	0.059	

V₁ - Grand Naine (AAA); V₂ - Poovan (AAB); V₃ - Ney Poovan (AB); T₁ - Control (Untreated); T₂ - Melatonin @ 200 µM; T₃ - Melatonin @ 400 µM; T₄ - Melatonin @ 600 µM; T₅ - Melatonin @ 800 µM and T₆ - Melatonin @ 1000 µM. Data represent the mean ± SE (n=4)




Fig 1. Effect of postharvest melatonin treatment on fruit firmness (Newton) at ambient storage (27±2°C) of Grand Naine (AAA) (V₁); Poovan (AAB) (V₂) and Ney Poovan (AB) (V₃); T₁- Control (Untreated); T₂ - Melatonin @ 200 µM; T₃ - Melatonin @ 400 µM; T₄ - Melatonin @ 600 µM; T₅ - Melatonin @ 800 µM and T₆ - Melatonin @ 1000 µM.

Table 2. Effect of melatonin on hand weight (kg) in different banana varieties under cold storage condition

Treatment details	Cold storage (17 °C)								
	1 st day	3 rd day	5 th day	7 th day	9 th day	11 th day	13 th day	15 th day	Mean
V ₁ T ₁	2.954 ± 0.04 ^a	2.768 ± 0.04 ^b	2.744 ± 0.04 ^b	2.712 ± 0.04 ^b	2.682 ± 0.04 ^b	2.654 ± 0.04 ^b	2.628 ± 0.04 ^b	2.592 ± 0.02 ^b	2.717
V ₁ T ₂	2.668 ± 0.04 ^c	2.490 ± 0.04 ^c	2.470 ± 0.04 ^c	2.444 ± 0.04 ^c	2.414 ± 0.04 ^c	2.396 ± 0.04 ^c	2.374 ± 0.04 ^c	2.342 ± 0.02 ^c	2.450
V ₁ T ₃	3.096 ± 0.03 ^a	3.060 ± 0.03 ^a	3.034 ± 0.03 ^a	3.006 ± 0.03 ^a	2.970 ± 0.03 ^a	2.940 ± 0.03 ^a	2.906 ± 0.03 ^a	2.867 ± 0.02 ^a	2.985
V ₁ T ₄	2.272 ± 0.03 ^e	2.238 ± 0.03 ^{de}	2.218 ± 0.03 ^{de}	2.192 ± 0.03 ^d	2.160 ± 0.03 ^d	2.136 ± 0.03 ^d	2.108 ± 0.03 ^d	2.076 ± 0.02 ^d	2.175
V ₁ T ₅	2.482 ± 0.04 ^d	2.448 ± 0.04 ^c	2.424 ± 0.04 ^c	2.396 ± 0.04 ^c	2.362 ± 0.03 ^c	2.338 ± 0.03 ^c	2.312 ± 0.03 ^c	2.278 ± 0.02 ^c	2.380
V ₁ T ₆	1.968 ± 0.04 ^f	1.940 ± 0.04 ^f	1.920 ± 0.04 ^f	1.898 ± 0.04 ^e	1.868 ± 0.04 ^f	1.848 ± 0.03 ^f	1.822 ± 0.03 ^f	1.796 ± 0.02 ^e	1.883
V ₂ T ₁	2.288 ± 0.03 ^e	2.250 ± 0.03 ^d	2.230 ± 0.03 ^d	2.198 ± 0.03 ^d	2.170 ± 0.03 ^d	2.140 ± 0.03 ^d	2.106 ± 0.03 ^d	1.774 ± 0.02 ^e	2.145
V ₂ T ₂	1.934 ± 0.03 ^f	1.788 ± 0.03 ^g	1.776 ± 0.03 ^g	1.654 ± 0.02 ^h	1.510 ± 0.02 ^h	1.494 ± 0.02 ^h	1.476 ± 0.02 ⁱ	1.458 ± 0.01 ^g	1.636
V ₂ T ₃	1.888 ± 0.02 ^f	1.626 ± 0.02 ^h	1.610 ± 0.02 ^h	1.582 ± 0.02 ^h	1.556 ± 0.02 ^h	1.534 ± 0.02 ^h	1.520 ± 0.02 ⁱ	1.426 ± 0.01 ^{gh}	1.593
V ₂ T ₄	2.292 ± 0.03 ^e	2.168 ± 0.03 ^e	2.150 ± 0.03 ^e	2.122 ± 0.03 ^d	2.076 ± 0.03 ^e	2.054 ± 0.03 ^e	2.030 ± 0.03 ^e	1.842 ± 0.02 ^e	2.092
V ₂ T ₅	1.802 ± 0.03 ^g	1.784 ± 0.03 ^g	1.770 ± 0.03 ^g	1.742 ± 0.03 ^g	1.702 ± 0.03 ^g	1.674 ± 0.02 ^g	1.606 ± 0.02 ^h	1.363 ± 0.01 ^h	1.680
V ₂ T ₆	1.900 ± 0.04 ^f	1.870 ± 0.04 ^f	1.838 ± 0.03 ^g	1.820 ± 0.03 ^f	1.794 ± 0.03 ^f	1.774 ± 0.03 ^f	1.720 ± 0.03 ^g	1.600 ± 0.02 ^f	1.790
V ₃ T ₁	1.002 ± 0.01 ^h	0.990 ± 0.01 ⁱ	0.982 ± 0.01 ⁱ	0.970 ± 0.01 ⁱ	0.952 ± 0.01 ⁱ	0.944 ± 0.01 ⁱ	0.934 ± 0.01 ^j	0.916 ± 0.01 ⁱ	0.961
V ₃ T ₂	0.806 ± 0.01 ^{ij}	0.798 ± 0.01 ^{jk}	0.788 ± 0.01 ^{jk}	0.780 ± 0.01 ^{jk}	0.764 ± 0.01 ^k	0.754 ± 0.01 ^k	0.742 ± 0.01 ^l	0.730 ± 0.01 ^k	0.770
V ₃ T ₃	0.872 ± 0.01 ⁱ	0.866 ± 0.01 ^j	0.862 ± 0.01 ^j	0.856 ± 0.01 ^j	0.844 ± 0.01 ^j	0.836 ± 0.01 ^j	0.824 ± 0.01 ^k	0.808 ± 0.00 ^j	0.846
V ₃ T ₄	0.678 ± 0.01 ^k	0.670 ± 0.01 ^l	0.668 ± 0.01 ^l	0.660 ± 0.01 ^l	0.654 ± 0.01 ^l	0.646 ± 0.01 ^l	0.634 ± 0.01 ^m	0.618 ± 0.00 ^l	0.654
V ₃ T ₅	0.712 ± 0.01 ^k	0.706 ± 0.01 ^l	0.700 ± 0.01 ^l	0.698 ± 0.01 ^l	0.690 ± 0.01 ^{kl}	0.682 ± 0.01 ^{kl}	0.670 ± 0.01 ^{lm}	0.658 ± 0.01 ^l	0.690
V ₃ T ₆	0.736 ± 0.01 ^{jk}	0.728 ± 0.01 ^{kl}	0.724 ± 0.01 ^{kl}	0.716 ± 0.01 ^{kl}	0.708 ± 0.01 ^{kl}	0.704 ± 0.01 ^{kl}	0.696 ± 0.01 ^{lm}	0.682 ± 0.01 ^{kl}	0.712
Mean	1.797	1.733	1.717	1.691	1.660	1.642	1.617	1.546	
SE.d	0.039	0.039	0.039	0.038	0.037	0.038	0.036	0.035	
CD (0.05)	0.077	0.077	0.078	0.076	0.075	0.074	0.073	0.070	

V₁ - Grand Naine (AAA); V₂ - Poovan (AAB); V₃ - Ney Poovan (AB); T₁ - Control (Untreated); T₂ - Melatonin @ 200 µM; T₃ - Melatonin @ 400 µM; T₄ - Melatonin @ 600 µM; T₅ - Melatonin @ 800 µM and T₆ - Melatonin @ 1000 µM. Data represent the mean ± SE (n=4).




Fig 2. Effect of postharvest melatonin treatment on fruit firmness (Newton) at cold storage (17°C) of Grand Naine (AAA) (V₁); Poovan (AAB) (V₂) and Ney Poovan (AB) (V₃); T₁ - Control (Untreated); T₂ - Melatonin @ 200 µM; T₃ - Melatonin @ 400 µM; T₄ - Melatonin @ 600 µM; T₅ - Melatonin @ 800 µM and T₆ - Melatonin @ 1000 µM.




Fig 3. Effect of postharvest melatonin treatment on total soluble solids ($^{\circ}$ Brix) at ambient storage ($27\pm 2^{\circ}\text{C}$) of Grand Naine (AAA) (V_1); Poovan (AAB) (V_2) and Ney Poovan (AB) (V_3); T₁- Control (Untreated); T₂- Melatonin @ 200 μM ; T₃- Melatonin @ 400 μM ; T₄- Melatonin @ 600 μM ; T₅- Melatonin @ 800 μM and T₆- Melatonin @ 1000 μM .

to the control (V_1T_1 : 7.4 to 24.2 $^{\circ}$ Brix), followed by Ney Poovan and Poovan (Fig. 4). Consistent with the current findings, Bal (34) revealed that melatonin treated at 1000 $\mu\text{M L}^{-1}$ concentration decreased the TSS content in plum. As a climacteric fruit, accelerated ripening and increased respiration convert complex carbohydrates into simple sugars, primarily glucose and fructose, which contribute to an increase in peaches total soluble solids (TSS) (40).

Effect of postharvest melatonin treatment on ethylene evolution

The data showed that ethylene production was significantly increased over the storage period across all the treatments. The climacteric peak of ethylene production was observed on day 7 and day 11 of ambient and cold storage conditions, consistent across all varieties. Among the three varieties, Grand Naine (V_1) displayed lower production of ethylene (0.78 $\mu\text{L kg}^{-1} \text{h}^{-1}$) throughout the storage period compared to Poovan (V_2 : 1.04 $\mu\text{L kg}^{-1} \text{h}^{-1}$) and Ney Poovan (V_3 : 0.89 $\mu\text{L kg}^{-1} \text{h}^{-1}$). Melatonin treatment at 1000 μM significantly reduced the ethylene production peak in all three varieties: V_1T_6 (0.59 $\mu\text{L kg}^{-1} \text{h}^{-1}$), V_2T_6 (0.80 $\mu\text{L kg}^{-1} \text{h}^{-1}$) and V_3T_6 (0.62 $\mu\text{L kg}^{-1} \text{h}^{-1}$) compared to the control (Fig. 5). Fruits stored under cold condition exhibited a gradual increase in ethylene production throughout the observation period. Dipping fruits in melatonin at 1000 μM resulted in the lowest ethylene production in Grand Naine (V_1T_6 : 0.47 $\mu\text{L kg}^{-1} \text{h}^{-1}$), Poovan (V_2T_6 : 0.87 $\mu\text{L kg}^{-1} \text{h}^{-1}$) and Ney Poovan (V_3T_6 : 0.68 $\mu\text{L kg}^{-1} \text{h}^{-1}$) compared to the control (V_1T_1 : 1.08; V_2T_1 : 1.17 and V_3T_1 : 1.12 $\mu\text{L kg}^{-1} \text{h}^{-1}$) at the end of cold storage condition (Fig. 6). Melatonin treatment has been shown to delay and suppress climacteric ethylene production and ACC accumulation, which aligns with melatonin mediated

reductions in the activities of ACS and ACO in mango (17). Exogenous melatonin treatment has been reported to reduce ethylene production and inhibit the expression of ACS and ACO in banana and pear ripened under ambient storage conditions (11, 41).

Effect of postharvest melatonin treatment on chlorophyll

Chlorophyll content was decreased progressively during the ripening period under both storage conditions. Ambient-stored fruits showed a more rapid decline in chlorophyll than cold conditions across all three varieties. On the 9th day of observation, melatonin-treated fruits (T_6 -1000 μM) retained higher total chlorophyll content in Grand Naine 18.52 $\mu\text{g g}^{-1}$ followed by Ney Poovan (11.30 $\mu\text{g g}^{-1}$) and Poovan (10.43 $\mu\text{g g}^{-1}$) compared to untreated. Compared to control, melatonin-treated fruits recorded a lower reduction of total chlorophyll content among all three varieties (Fig. 7).

Grand Naine retained chlorophyll content for a more extended period in melatonin-treated fruits (V_1T_6 : 9.41 $\mu\text{g g}^{-1}$) of total chlorophyll followed by Ney Poovan (V_3T_6 : 4.64 $\mu\text{g g}^{-1}$) and Poovan (V_2T_6 : 4.43 $\mu\text{g g}^{-1}$) than control on 15th day of cold condition (Fig. 8). Dipping lettuce in melatonin at concentrations of 100 and 1000 $\mu\text{mol L}^{-1}$ effectively delayed the decrease in total chlorophyll content compared to the control (42). Similarly, Broccoli treated with 100 $\mu\text{mol L}^{-1}$ melatonin maintained 24.15 % higher chlorophyll content than the control on the 6th day of storage (43). Chlorophyll degradation during postharvest storage is primarily caused by reduced chlorophyll-binding proteins and increased chlorophyllase enzyme activity, which melatonin may counteract by reducing reactive oxygen species (ROS) and lipid radicals




Fig. 4. Effect of postharvest melatonin treatment on total soluble solids ($^{\circ}$ Brix) at cold storage (17°C) of Grand Naine (AAA) (V_1); Poovan (AAB) (V_2) and Ney Poovan (AB) (V_3); T₁ - Control (Untreated); T₂ - Melatonin @ 200 μM ; T₃ - Melatonin @ 400 μM ; T₄ - Melatonin @ 600 μM ; T₅ - Melatonin @ 800 μM and T₆ - Melatonin @ 1000 μM .




Fig. 5. Effect of postharvest melatonin treatment on ethylene evolution ($\mu\text{L kg}^{-1} \text{h}^{-1}$) at ambient storage ($27\pm 2^{\circ}\text{C}$) of Grand Naine (AAA) (V_1); Poovan (AAB) (V_2) and Ney Poovan (AB) (V_3); T₁ - Control (Untreated); T₂ - Melatonin @ 200 μM ; T₃ - Melatonin @ 400 μM ; T₄ - Melatonin @ 600 μM ; T₅ - Melatonin @ 800 μM and T₆ - Melatonin @ 1000 μM .




Fig. 6. Effect of postharvest melatonin treatment on ethylene evolution ($\mu\text{L kg}^{-1} \text{h}^{-1}$) at cold storage (17°C) of Grand Naine (AAA) (V_1); Poovan (AAB) (V_2) and Ney Poovan (AB) (V_3); T₁ - Control (Untreated); T₂ - Melatonin @ 200 μM ; T₃ - Melatonin @ 400 μM ; T₄ - Melatonin @ 600 μM ; T₅ - Melatonin @ 800 μM and T₆ - Melatonin @ 1000 μM .




Fig. 7. Effect of postharvest melatonin treatment on total chlorophyll ($\mu\text{g g}^{-1}$) at ambient storage ($27\pm 2^{\circ}\text{C}$) of Grand Naine (AAA) (V_1); Poovan (AAB) (V_2) and Ney Poovan (AB) (V_3); T₁ - Control (Untreated); T₂ - Melatonin @ 200 μM ; T₃ - Melatonin @ 400 μM ; T₄ - Melatonin @ 600 μM ; T₅ - Melatonin @ 800 μM and T₆ - Melatonin @ 1000 μM .




Fig. 8. Effect of postharvest melatonin treatment on total chlorophyll ($\mu\text{g g}^{-1}$) at cold storage (17°C) of Grand Naine (AAA) (V_1); Poovan (AAB) (V_2) and Ney Poovan (AB) (V_3); T₁ - Control (Untreated); T₂ - Melatonin @ 200 μM ; T₃ - Melatonin @ 400 μM ; T₄ - Melatonin @ 600 μM ; T₅ - Melatonin @ 800 μM and T₆ - Melatonin @ 1000 μM .

and thus potentially delaying chlorophyll degradation (44).

Effect of postharvest melatonin treatment on starch degradation

Starch degradation began in the core of the pulp, the placenta and the endocarp, while the mesocarp retained starch for the most extended period during maturation (45). Fig. 9 and 10 illustrates the starch breakdown in cross-sections of three banana varieties taken from the centre position of the fruit during the ripening period. Starch exhibits a blue-black colouration, whereas regions with depleted starch appear white. During ripening, starch depletion progresses from the centre of the banana towards the periphery. Differences were noticed in the changes in starch degradation pattern on banana pulp between the treatments until the 9th and 15th day of ambient and cold storage conditions, respectively. Bananas treated with 1000 μM melatonin exhibited the most prolonged ripening duration and extended shelf life compared to the control under both storage conditions. The starch pattern of each fruit was compared using the starch pattern of each fruit with the starch iodine staining chart for bananas developed by Blankenship and Ellsworth (31). Melatonin-treated fruits showed about 5 % of starch, while the untreated (control) fruits had >65 % on the 9th and 15th day of ambient and cold storage conditions (Table 3-4). Starch-to-sugar conversion is influenced by the levels of O_2 and CO_2 , which can inhibit the enzyme activities involved in the starch hydrolysis process (46, 47). Immersing bananas in melatonin from 0.05 to 0.5 mM suppressed ethylene production and starch breakdown,

delaying the ripening process during postharvest storage.

Effect of postharvest melatonin treatment on shelf life

Fruit shelf life is a critical factor significantly affecting the quality of horticultural commodities, especially bananas, during the postharvest storage period. Melatonin treatments offer a potential solution for extending shelf life. It is determined through continuous assessment of the fruits' physical appearance and quality analysis tests results throughout the storage period (48). Maximum shelf life was recorded in T₆ (treated with melatonin at 1000 μM) of Grand Naine (10.25 days), Poovan (9.00 days) and Ney Poovan (9.45 days) compared with control (T₁) fruits at ambient storage conditions. The shelf life of fruits was prolonged when stored under cold conditions. Among the varieties, Grand Naine had the most extended shelf life of 17.58 days, followed by Poovan (16.98 days) and Ney Poovan (16.28 days) in fruits treated with 1000 μM of melatonin, compared to the control (T₁). The results are presented in Fig. 11. However, fruits dipped in 1000 μM of melatonin were considered the best treatment among the three banana varieties, viz., Grand Naine, Poovan and Ney Poovan, under two different storage conditions to increase the fruit shelf life during the postharvest storage period.




Fig. 9. Effect of postharvest melatonin treatment on starch degradation at 1st and 9th day of ambient storage condition of Grand Naine (AAA) (V₁); Poovan (AAB) (V₂) and Ney Poovan (AB) (V₃); T₁ - Control (Untreated); T₂ - Melatonin @ 200 µM; T₃ - Melatonin @ 400 µM; T₄ - Melatonin @ 600 µM; T₅ - Melatonin @ 800 µM and T₆ - Melatonin @ 1000 µM.



Fig. 10. Effect of postharvest melatonin treatment on starch degradation at 1st and 15th day of cold storage condition of Grand Naine (AAA) (V₁); Poovan (AAB) (V₂) and Ney Poovan (AB) (V₃); T₁ - Control (Untreated); T₂ - Melatonin @ 200 µM; T₃ - Melatonin @ 400 µM; T₄ - Melatonin @ 600 µM; T₅ - Melatonin @ 800 µM and T₆ - Melatonin @ 1000 µM.

Table 3. Effect of postharvest melatonin treatment on starch conversion analysis by starch-iodine test on 1st and 9th day of ambient storage condition

Treatments	Scale (1-10) based on starch content					
	Ambient storage (27 ± 2 °C)					
	Grand Naine (AAA)	Poovan (AAB)	Ney Poovan (AB)	Grand Naine (AAA)	Poovan (AAB)	Ney Poovan (AB)
T ₁	1	1	1	4	10	4
T ₂	1	1	1	2	9	2
T ₃	1	1	1	2	9	1
T ₄	1	1	1	1	8	1
T ₅	1	1	1	1	7	1
T ₆	1	1	1	1	4	1

V₁ - Grand Naine (AAA); V₂ - Poovan (AAB); V₃ - Ney Poovan (AB); T₁ - Control (Untreated); T₂ - Melatonin @ 200 µM; T₃ - Melatonin @ 400 µM; T₄ - Melatonin @ 600 µM; T₅ - Melatonin @ 800 µM and T₆ - Melatonin @ 1000 µM. Treatment groups are scaled from 1 to 10 on different storage conditions. The data shown was one of the four replicates.

Table 4. Effect of postharvest melatonin treatment on starch conversion analysis by starch-iodine test on 1st and end 15th day of cold storage condition

Treatments	Scale (1-10) based on starch content					
	Cold storage (17 °C)					
	1 st day of observation			15 th day of observation		
	Grand Naine (AAA)	Poovan (AAB)	Ney Poovan (AB)	Grand Naine (AAA)	Poovan (AAB)	Ney Poovan (AB)
T ₁	1	1	1	8	10	4
T ₂	1	1	1	7	9	3
T ₃	1	1	1	6	7	2
T ₄	1	1	1	5	8	1
T ₅	1	1	1	4	6	1
T ₆	1	1	1	4	6	1

V₁ - Grand Naine (AAA); V₂ - Poovan (AAB); V₃ - Ney Poovan (AB); T₁ - Control (Untreated); T₂ - Melatonin @ 200 µM; T₃ - Melatonin @ 400 µM; T₄ - Melatonin @ 600 µM; T₅ - Melatonin @ 800 µM and T₆ - Melatonin @ 1000 µM. Treatment groups are scaled from 1 to 10 on different storage conditions. The data shown was one of the four replicates.

Conclusion

Bananas have a short postharvest shelf life due to climacteric ripening, making effective techniques crucial for extending shelf life and maintaining its quality. Recent studies showed that melatonin can influence postharvest physiology and prolong shelf life in various crops. Our findings revealed that melatonin significantly prolonged shelf life and maintained the morpho-physiological, biochemical and fruit quality characteristics of Grand Naine, Poovan and Ney Poovan varieties compared to untreated control. The 1000 µM melatonin concentration produced the most substantial effect across all three banana varieties. These results suggest that postharvest melatonin treatment could be a promising approach for enhancing bananas' quality and storage potential.

Acknowledgements

The authors thank the Department of Crop Physiology, Tamil Nadu Agricultural University, Coimbatore, for providing research facilities and their support.

Authors' contributions

MD, PJ, VR, LA, MB and MK conceived the concept, designed the study and prepared the manuscript. MD carried out the experimental procedures and statistical analyses. All authors reviewed and endorsed the final manuscript.

Compliance with ethical standards

Conflict of interest: No potential conflict of interest was reported by the author(s).

Ethical issues: None

References

- Aghdam MS, Fard JR. Melatonin treatment attenuates postharvest decay and maintains nutritional quality of strawberry fruits (*Fragaria × ananassa* cv. Selva) by enhancing GABA shunt activity. *Food Chem.* 2017;221:1650–57. <https://doi.org/10.1016/j.foodchem.2016.10.123>
- Arnao MB, Hernández-Ruiz J. Protective effect of melatonin against chlorophyll degradation during the senescence of barley leaves. *J Pineal Res.* 2009;46(1):58–63. <https://doi.org/10.1111/j.1600-079X.2008.00625.x>
- Back K, Tan DX, Reiter RJ. Melatonin biosynthesis in plants: multiple pathways catalyze tryptophan to melatonin in the cytoplasm or chloroplasts. *J Pineal Res.* 2016;61(4):426–37. <https://doi.org/10.1016/j.postharvbio.2021.111571>
- Bal E. Effect of melatonin treatments on biochemical quality and postharvest life of nectarines. *J Food Meas Charact.* 2021;15(1):288–95. <https://doi.org/10.1007/s11694-020-00636-5>
- Bal E. Physicochemical changes in s'anta Rosa' plum fruit treated with melatonin during cold storage. *J Food Meas Charact.* 2019;13:1713–20. <https://doi.org/10.1007/s11694-019-00088-6>
- Bantayehu M, Alemayehu M. Efficacy of postharvest technologies on ripening behavior and quality of banana varieties grown in Ethiopia. *Int J Fruit Sci.* 2020;20(1):59–75. <https://doi.org/10.1080/15538362.2019.1583623>
- Blankenship SM, Ellsworth DD, Powell RL. A ripening index for banana fruit based on starch content. *Hort Techn.* 1993;3(3):338–39. <https://doi.org/10.21273/HORTTECH.3.3.338>
- Cao S, Bian K, Shi L, Chung H-H, Chen W, Yang Z. Role of melatonin in cell-wall disassembly and chilling tolerance in cold-stored peach fruit. *J Agric Food Chem.* 2018;66(22):5663–70. <https://doi.org/10.1021/acs.jafc.8b02055>
- Cheng J, Zheng A, Li H, Huan C, Jiang T, Shen S. Effects of melatonin treatment on ethanol fermentation and ERF expression in kiwifruit cv. Bruno during postharvest. *Scientia Horticulturae.* 2022;293:110696. <https://doi.org/10.1016/j.scienta.2021.110696>
- Dwivany FM, Aprilyandi AN, Suendo V, Sukriandi N. Carrageenan edible coating application prolongs Cavendish banana shelf life. *Int J Food Sci.* 2020;2020(1):8861610. <https://doi.org/10.1155/2020/8861610>
- Fan S, Li Q, Feng S, Lei Q, Abbas F, Yao Y. Melatonin maintains fruit quality and reduces anthracnose in postharvest papaya via enhancement of antioxidants and inhibition of pathogen development. *Antioxidants.* 2022;11(5):804. <https://doi.org/10.3390/antiox11050804>
- Fan S, Xiong T, Lei Q, Tan Q, Cai J, Song Z. Melatonin treatment improves postharvest preservation and resistance of guava fruit (*Psidium guajava* L.). *Foods.* 2022;11(3):262. <https://doi.org/10.3390/foods11030262>
- Gao H, Zhang ZK, Chai HK, Cheng N, Yang Y, Wang DN. Melatonin treatment delays postharvest senescence and regulates reactive oxygen species metabolism in peach fruit. *Postharvest Biol Technol.* 2016;118:103–0. <https://doi.org/10.1016/j.postharvbio.2016.03.006>
- Gurjar P, Singh S, Verma A, Mishra M. Post-harvest melatonin application reduced browning in minimally processed lettuce (*Lactuca sativa* L.) during low temperature storage. *J Hortic Sci.* 2023;18(1):195–200. <https://doi.org/10.24154/jhs.v18i1.2163>
- Hiscox J, Israelstam G. A method for the extraction of chlorophyll from leaf tissue without maceration. *Canad J Bot.* 1979;57(12):1332–4. <https://doi.org/10.1139/b79-163>

16. Miao HM, Ying LJ, Ping RJ. Effect of melatonin on ripening and senescence of postharvest kiwifruits. *Food Sci.* 2018;39(19):226–32. <https://doi.org/10.7506/spkx1002-6630-201819035>
17. Hu W, Yang H, Tie W, Yan Y, Ding Z, Liu Y. Natural variation in banana varieties highlights the role of melatonin in postharvest ripening and quality. *J Agric Food Chem.* 2017;65(46):9987–94. <https://pubs.acs.org/doi/abs/10.1021/acs.jafc.7b03354>
18. Inaba A, Liu X, Yokotani N, Yamane M, Lu W-J, Nakano R. Differential feedback regulation of ethylene biosynthesis in pulp and peel tissues of banana fruit. *J Exp Bot.* 2007;58(5):1047–57. <https://doi.org/10.1093/jxb/ert265>
19. Jayarajan S, Sharma R. Impact of nitric oxide on shelf life and quality of nectarine (*Prunus persica* var. *nucipersica*). *Acta Physiol Plant.* 2018;40(12):207. <https://doi.org/10.1007/s11738-018-2779-4>
20. Li S, Xu Y, Bi Y, Zhang B, Shen S, Jiang T. Melatonin treatment inhibits gray mold and induces disease resistance in cherry tomato fruit during postharvest. *Postharvest Biol Technol.* 2019;157:110962. <https://doi.org/10.1016/j.postharvbio.2019.110962>
21. Liu C, Zheng H, Sheng K, Liu W, Zheng L. Effects of melatonin treatment on the postharvest quality of strawberry fruit. *Postharvest Biol Technol.* 2018;139:47–55. <https://doi.org/10.1016/j.postharvbio.2018.01.016>
22. Liu J, Yang J, Zhang H, Cong L, Zhai R, Yang C. Melatonin inhibits ethylene synthesis via nitric oxide regulation to delay postharvest senescence in pears. *J Agric Food Chem.* 2019;67(8):2279–88. <https://doi.org/10.1021/acs.jafc.8b06580>
23. Liu S, Huang H, Huber DJ, Pan Y, Shi X, Zhang Z. Delay of ripening and softening in 'Guifei' mango fruit by postharvest application of melatonin. *Postharvest Biol Technol.* 2020;163:111136. <https://doi.org/10.1016/j.postharvbio.2020.111136>
24. Lorente-Mento JM, Guillén F, Castillo S, Martínez-Romero D, Valverde JM, Valero D. Melatonin treatment to pomegranate trees enhances fruit bioactive compounds and quality traits at harvest and during postharvest storage. *Antioxidants.* 2021;10(6):820. <https://doi.org/10.3390/antiox10060820>
25. Lustriane C, Dwivany FM, Suendo V, Reza M. Effect of chitosan and chitosan-nanoparticles on post harvest quality of banana fruits. *J Plant Biotechnol.* 2018;45(1):36–44. <https://doi.org/10.5010/JPB.2018.45.1.036>
26. Ma Q, Lin X, Wei Q, Yang X, Zhang Yn, Chen J. Melatonin treatment delays postharvest senescence and maintains the organoleptic quality of 'Newhall' navel orange (*Citrus sinensis* (L.) Osbeck) by inhibiting respiration and enhancing antioxidant capacity. *Sci Hortic.* 2021;286:110236. <https://doi.org/10.1016/j.scienta.2021.110236>
27. Mahajan PV, Caleb OJ, Singh Z, Watkins CB, Geyer M. Postharvest treatments of fresh produce. *Philos Transac Royal Soc Math Physic Engin Sci.* 2014;372(2017):20130309. <https://doi.org/10.1098/rsta.2013.0309>
28. Maqbool M, Ali A, Alderson PG, Zahid N, Siddiqui Y. Effect of a novel edible composite coating based on gum arabic and chitosan on biochemical and physiological responses of banana fruits during cold storage. *J Agric Food Chem.* 2011;59(10):5474–82. <https://doi.org/10.1021/jf200623m>
29. Mubarak S, Rahman MI, Kamaluddin N, Solihin E. Impact of 1-methylcyclopropene combined with chitosan on postharvest quality of tropical banana 'Lady Finger'. *Int J Food Prop.* 2022;25(1):1171–85. <https://doi.org/10.1080/10942912.2022.2074028>
30. Onik JC, Wai SC, Li A, Lin Q, Sun Q, Wang Z. Melatonin treatment reduces ethylene production and maintains fruit quality in apple during postharvest storage. *Food Chem.* 2021;337:127753. <https://doi.org/10.1016/j.foodchem.2020.127753>
31. Ringer T, Blanke M. Non-invasive, real time in-situ techniques to determine the ripening stage of banana. *J Food Meas Charact.* 2021;15(5):4426–37. <https://doi.org/10.1007/s11694-021-01009-2>
32. Sánchez M, Villalta R, Guzmán M. Combate de enfermedades poscosecha en banano (*Musa AAA*) con extractos metanólicos de hojas de *Musa* spp. con diferente grado de resistencia a sigatoka negra. *CORBANA.* 2021;66:15–28.
33. Sati H, Bhardwaj R, Fawole OA, Pareek S. Postharvest melatonin application preserves quality and imparts chilling tolerance in peaches. *J Food Biochem.* 2023;2023(1):8126640. <https://doi.org/10.1155/2023/8126640>
34. Shang F, Liu R, Wu W, Han Y, Fang X, Chen H. Effects of melatonin on the components, quality and antioxidant activities of blueberry fruits. *Lebensmittel-Wissenschaft and Technologie (LWT).* 2021;147:111582. <https://doi.org/10.1016/j.lwt.2021.111582>
35. Sun Q, Zhang N, Wang J, Zhang H, Li D, Shi J. Melatonin promotes ripening and improves quality of tomato fruit during postharvest life. *J Exp Bot.* 2015;66(3):657–68. <https://doi.org/10.1093/jxb/eru332>
36. Tijero V, Munoz P, Munne-Bosch S. Melatonin as an inhibitor of sweet cherries ripening in orchard trees. *Plant Physiol Biochem.* 2019;140:88–95. <https://doi.org/10.1016/j.plaphy.2019.05.007>
37. Wang C, Yin LY, Shi XY, Xiao H, Kang K, Liu XY. Effect of cultivar, temperature and environmental conditions on the dynamic change of melatonin in mulberry fruit development and wine fermentation. *J Food Sci.* 2016;81(4):M958–M67. <https://doi.org/10.1111/1750-3841.13263>
38. Wang Z, Pu H, Shan S, Zhang P, Li J, Song H. Melatonin enhanced chilling tolerance and alleviated peel browning of banana fruit under low temperature storage. *Postharvest Biol Technol.* 2021;179:111571. <https://doi.org/10.1016/j.postharvbio.2021.111571>
39. Wei W, Yang Y-y, Wu C-j, Kuang J-f, Chen J-y, Lu W-j. MaMADS1-MaNAC083 transcriptional regulatory cascade regulates ethylene biosynthesis during banana fruit ripening. *Hortic Res.* 2023;10(10):uhad177. <https://doi.org/10.1093/hr/uhad177>
40. Xiao Y-y, Chen J-y, Kuang J-f, Shan W, Xie H, Jiang Y-m. Banana ethylene response factors are involved in fruit ripening through their interactions with ethylene biosynthesis genes. *J Exp Bot.* 2013;64(8):2499–510. <https://doi.org/10.1093/jxb/ert108>
41. Ze Y, Gao H, Li T, Yang B, Jiang Y. Insights into the roles of melatonin in maintaining quality and extending shelf life of postharvest fruits. *Trends Food Sci Technol.* 2021;109:569–78. <https://doi.org/10.1016/j.tifs.2021.01.051>
42. Zhai R, Liu J, Liu F, Zhao Y, Liu L, Fang C. Melatonin limited ethylene production, softening and reduced physiology disorder in pear (*Pyrus communis* L.) fruit during senescence. *Postharvest Biol Technol.* 2018;139:38–46. <https://doi.org/10.1016/j.postharvbio.2018.01.017>
43. Zhang H, Liu X, Chen T, Ji Y, Shi K, Wang L. Melatonin in apples and juice: Inhibition of browning and microorganism growth in apple juice. *Molecules.* 2018;23(3):521. <https://doi.org/10.3390/molecules23030521>
44. Zhang H, Wang L, Shi K, Shan D, Zhu Y, Wang C. Apple tree flowering is mediated by low level of melatonin under the regulation of seasonal light signal. *J Pineal Res.* 2019;66(2):e12551. <https://doi.org/10.1111/jpi.12551>
45. Zhang N, Zhang H-J, Sun Q-Q, Cao Y-Y, Li X, Zhao B. Proteomic analysis reveals a role of melatonin in promoting cucumber seed germination under high salinity by regulating energy production. *Sci Rep.* 2017;7(1):503. <https://doi.org/10.1038/s41598-017-00566-1>
46. Zhang W, Cao J, Fan X, Jiang W. Applications of nitric oxide and melatonin in improving postharvest fruit quality and the separate and crosstalk biochemical mechanisms. *Trends Food Sci Technol.* 2020;99:531–41. <https://doi.org/10.1016/j.tifs.2020.03.024>
47. Zhang Z, Wang T, Liu G, Hu M, Yun Z, Duan X. Inhibition of downy blight and enhancement of resistance in litchi fruit by postharvest application of melatonin. *Food Chem.* 2021;347:129009. <https://doi.org/10.1016/j.foodchem.2021.129009>
48. Zhu L, Hu H, Luo S, Wu Z, Li P. Melatonin delaying senescence of postharvest Broccoli by regulating respiratory metabolism and antioxidant activity. *Trans Chinese Soc Agric Engin.* 2018;34(3):300–8.