



RESEARCH ARTICLE

In-situ decomposition of sugarcane trash using microbial consortium and its impact on ratoon cane

Sugumaran M P^{1*}, Gayathry Gunavijayan^{2*}, Porkodi G³, Kalaiselvi P⁴, Kiruba M⁵, Sassikumar D⁶, Natarajan K⁷ & Bharathi Kumar K⁸

¹Department of Environmental Sciences, Tamil Nadu Agricultural University, Krishi Vigyan Kendra, Vridhachalam 606 001, Tamil Nadu, India

²Department of Agricultural Microbiology, Tamil Nadu Agricultural University, Krishi Vigyan Kendra, Vridhachalam 606 001, Tamil Nadu, India

³Department of Soil Science and Agricultural Chemistry, Agricultural College and Research Institute, Kudumiyamalai 625 104, Pudhukottai, Tamil Nadu, India

⁴Department of Agronomy, Tamil Nadu Agricultural University, Krishi Vigyan Kendra, Vridhachalam 606 001, Tamil Nadu, India

⁵Department of Forestry, Tamil Nadu Agricultural University, Krishi Vigyan Kendra, Sandhiyur 636 203, Tamil Nadu, India

⁶Department of Plant Breeding and Genetics, Sugarcane Research Station, Tamil Nadu Agricultural University, Cuddalore 607 001, Tamil Nadu, India

⁷Department of Seed Science and Technology, Tamil Nadu Agricultural University, Krishi Vigyan Kendra, Vridhachalam 606 001, Tamil Nadu, India

⁸Department of Plant Breeding and Genetics, Tamil Nadu Agricultural University, Krishi Vigyan Kendra, Vridhachalam 606 001, Tamil Nadu, India

*Correspondence email - gayathryg@tnau.ac.in; sugumaran.mp@tnau.ac.in

Received: 30 December 2024; Accepted: 04 March 2025; Available online: Version 1.0: 22 April 2025

Cite this article: Sugumaran MP, Gayathry G, Porkodi G, Kalaiselvi P, Kiruba M, Sassikumar D, Natarajan K, Bharathi KK. *In-situ* decomposition of sugarcane trash using microbial consortium and its impact on ratoon cane. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.6969>

Abstract

Sugarcane trash, consisting of leaves and stalk residues left after harvesting, poses challenges such as slow decomposition, nutrient imbalance and environmental concerns. This study explores an eco-friendly approach to sugarcane trash management using a microbial consortium to accelerate decomposition and improve soil health and ratoon cane yield. These microorganisms facilitate sugarcane trash decomposition by reducing the C: N ratio and breaking down lignocellulosic compounds. Applying a trash decomposer at 30 kg/ha on the 3rd, 15th and 30th days after harvest, along with 100% RDF fertilizers, significantly reduced sugarcane trashes' C: N ratio to 18.25 by the 60th day. This treatment also improved growth and yield parameters in the ratoon crop. FTIR analysis confirmed compost maturity, shifting from aliphatic to aromatic compounds. The nitrate band at 1384 per cm intensified, while peak ratio changes indicated intensified decomposition progress of the trash. A higher proportion of cellulose-degrading consortia in trash D promoted the breakdown of complex molecules and reduced humification during *in-situ* decomposition. The findings suggest that integrating microbial decomposition strategies with lignocellulosic residues could enhance soil fertility, thus enhancing nutrient availability.

Keywords: decomposition; microbial consortium; ratoon cane; sugarcane; trash; *in-situ*

Introduction

Sugarcane (*Saccharum officinarum* L.) is a vital crop in the agro-industry, primarily cultivated for sugar production and bioethanol. However, sugarcane cultivation generates significant residual biomass, known as sugarcane trash, which remains in the field post-harvest. Improper management of this residue poses challenges such as soil degradation, increased pest incidence and environmental pollution (1). Sustainable alternatives, including microbial-assisted decomposition, are gaining attention as eco-friendly solutions for efficient sugarcane trash management (2). After harvesting, sugarcane trash accounts for 10-20% of the total crop biomass (3). If effectively decomposed, it can contribute significant nutrients to the soil, including 30 kg of nitrogen, 9 kg of phosphorus and 24 kg of potassium per hectare (4). Sugarcane's remarkable dry matter output makes it a valuable

crop for biofuel, which is also used to produce sugar. Climate, soil type, crop management strategies, fertilizer management (5), irrigation timing and soil moisture availability during the growth period are some elements that affect sugarcane productivity. However, due to its high lignocellulose and silica content, natural decomposition is slow, necessitating external interventions such as microbial decomposition for rapid nutrient recycling. Sugarcane trash can be beneficial when appropriately managed. Mulching with sugarcane trash improves soil moisture retention, reduces erosion and enhances microbial activity (6). However, its high carbon-to-nitrogen (C:N) ratio (often exceeding 124:1) limits microbial decomposition, making external microbial inoculation necessary to accelerate breakdown and improve nutrient availability (5, 7).

Microbial inoculation has been widely studied to accelerate organic matter degradation (8, 9). Specific microbial consortia, including fungi and actinobacteria, have been shown to enhance cellulose degradation and improve substrate utilization (10). However, sugarcane trash presents decomposition challenges due to its high silica content and large C:N ratio (124:1), requiring microbial amendments for efficient breakdown. This study evaluates the potential of a microbial consortium-based trash decomposer to enhance *in-situ* decomposition and improve ratoon cane yield (11, 12). Wider usage of specific trash-decomposing fungal genera such as *Chaetomium globosum* and silica/potash solubilizing *Bacillus mucilaginosus* along with yeast molasses, rock phosphate and gypsum is highly limiting in most of the previous research and the present study aims to fill the research gap so to develop a microbial consortium for *in-situ* trash decomposition, especially in ratoon cane.

Materials and Methods

The experiment was conducted from July 2022 to August 2023 at the Sugarcane Research Station, Cuddalore. The site has a clayey loam soil type, with an annual mean rainfall of 1210 mm and an altitude of 4.6 m above sea level (Fig. 1).

Experimental design

A microbial consortium, Trash Decomposer (Trash D), was developed to accelerate sugarcane trash decomposition.

TNAU Trash D formulation

Trash D formulation consisted of the following components per tonne of trash (Fig. 3):

- *Bacillus mucilaginosus* (10^9 cfu/mL at 40 mL)
- *Chaetomium globosum* TNAU cg 6 and *Trichoderma viridi* TNAU TV1 each (10^6 cfu/mL at 30 mL each).
- Yeast molasses (20 L)

- Rock phosphate (20 kg)

- Gypsum (20 kg)

Treatments

T ₁	:	Trash D at 20 kg/ha* + 125 % RDF
T ₂	:	Trash D at 30 kg/ha* + 125 % RDF
T ₃	:	Trash D at 20 kg/ha* + 100 % RDF
T ₄	:	Trash D at 30 kg/ha* + 100 % RDF
T ₅	:	Trash D at 20 kg/ha* + 75 % RDF
T ₆	:	Trash D at 30 kg/ha* + 75 % RDF
T ₇	:	100% RDF alone + untreated trashes (Control)

* Trash D was applied in three stages: (i) on the 3rd day after harvest, (ii) on the 15th day and (iii) on the 30th day. Irrigation was provided uniformly to enhance microbial activity. Standard ratoon management practices were followed across all treatments, including stubble shaving, root pruning, thinning and earthing up.

Compost analysis

Trash decomposition was monitored by collecting samples on the 10th, 30th and 60th days after harvest. Organic carbon was estimated per Walkley and Blacks' procedures and total nitrogen was calculated using the Macro-Kjeldhal method (13, 14). The samples were analyzed using standard procedures for pH, organic carbon, total nitrogen, phosphorus, potassium and C: N ratio.

Effect of Trash D on crop yield

At harvest, tiller population, cane height, number of nodes, cane girth, individual cane weight, Commercial Cane Sugar (CCS%) and total cane yield (t/ha) were recorded.

Statistical Analysis

The experimental data were statistically scrutinized using the standard method (15). The critical difference was achieved at a 5 % (0.05) probability.




Fig. 1. Study area.

Result and Discussion

Characterization of soil

The experimental soil had a pH of 6.9 (slightly acidic to neutral) and an EC of 0.25 dSm^{-1} , indicating low to moderate salinity. Organic carbon content was 0.32%, while available nitrogen, phosphorus and potassium levels were 157.5, 35.0 and 145.0 kg/ha , respectively, sufficient for sugarcane growth.

Overall, the soil characteristics in this Table suggest that the soil has a pH close to neutral and moderate organic carbon content. Favourable levels of essential nutrients such as nitrogen, phosphorus and potassium (Fig. 2). If soil organic matter is consistently cultivated for agricultural

production, it decomposes quickly due to changes in the soils' physical qualities, particularly in tropical and semiarid countries (16). This results in a decrease in soil productivity and soil carbon depletion (17). Applying appropriate input management techniques can improve carbon sequestration and decrease atmospheric CO_2 enrichment (18). Furthermore, the addition of plant remains may have an impact on the soil microclimate (19).

FTIR analysis

FTIR analysis confirmed compost maturity, shifting from aliphatic to aromatic compounds. The nitrate band at 1384 cm^{-1} intensified, while peak ratio changes ($1384/2925$ and $1034/1384$) indicated decomposition progress (Fig. 4). The presence of amides, alkanes, phenols and aromatics further validated microbial activity in degrading sugarcane trash. Vibrations of the methylene group in the spectral range from 1375 to 1150 per cm can be observed in the FTIR spectra (Table 1). Similar results were observed in straw materials. Cellulose and hemicellulose are the most easily broken down by bacteria, whereas lignin is the most difficult (20).

In the closed pile, the quantity of this band was gradually increased and then decreased in composting. This band intensity of inoculated compost by microorganisms increased slightly after the composting process (21) and the highest intensity was observed in closed piles in 16 days (22). The FTIR spectra of the dissolved organic matter showed a decline in aliphatic compounds, protein class and carbohydrates in the infected pile compared to the closed pile (Fig. 7). FTIR absorption peaks confirmed and supplemented the previous results related to the rapid disappearance of compost readily biodegradable components and increased concentration of aromatic compounds due to the compost of microorganisms (23).

Scanning electron microscopy (SEM) analysis

The morphology of sugarcane fresh leaf and composted material was analyzed using SEM, providing insights into their shape, growth mechanisms and size. SEM analysis revealed the structural breakdown of sugarcane trash after decomposition. Composted material showed reduced fibre integrity and increased microbial colonization compared to fresh leaves, confirming microbial action in lignocellulose degradation (Fig. 5).




Fig. 3. Trash decomposer (Trash D formulation).




Fig. 2. Chemical properties of the experimental soil.






Fig. 4. FTIR analysis of trash D treated sugarcane trash.

Table 1. Effect of Trash D on sugarcane trash *in-situ* decomposition by FTIR analysis

Absorbed Peak value (cm ⁻¹)	Bonds	Functional groups
3918.30	C-H stretch	Alkanes (Aliphatic)
3776.30		
2062.80	N-H stretch	1°, 2° amines, amides, alkalides
2374.86		
1654.92	N-C=O	Amide carbonyl stretching vibration in carbon bond
1960.41	C=C	Secondary aromatic
2508.33		
1384.89	O-H	Phenolic; ligneous



A) Fresh sugarcane trash



B) Composted sugarcane trash

Fig. 5 A. SEM analysis of fresh; **B.** Composted sugarcane trash.

Effect of TNAU Trash D on C: N ratio

The graphical data on different treatments (T_1 to T_7) and their impact on trash properties, including total carbon percentage, total nitrogen percentage and the carbon to nitrogen (C: N) ratio at various time points such as the initial, 10th day, 30th day and 60th day is presented in Fig. 6. The total carbon percentage represents the proportion of carbon in the sample. A reduced C: N ratio typically indicates higher decomposition and nutrient mineralization. The C: N ratio varies among treatments and periods, reflecting differences in nutrient dynamics and decomposition rates (Table 2). Sugarcane trash mulch conserves soil organic matter, essential for maintaining soil fertility and better crop growth over a more extended period

(24). These parameters are crucial for understanding nutrient cycling, decomposition processes and soil health.

Trash D significantly reduced the C: N ratio, indicating enhanced decomposition and nutrient mineralization. The initial C:N ratio was significantly varied and was recorded as 40.75:0.59 at and after the different treatment effects; the lowest C: N ratio (18.25) was observed in trash D at 30 kg/ha + 100% RDF. At the same time, the control had the slowest breakdown. The treatment details are provided in Table 3. The data helps identify treatments that may promote more favourable carbon and nitrogen dynamics in the soil, influencing plant growth and crop yield. Treatments with a reduced C: N ratio (trash D @ 30 kg/ha+ 100 % RDF) indicate




Fig. 6. C: N ratio of sugarcane trash decomposed *in-situ* by trash D




Fig. 7. Field application of trash D over the trashes and its effect on growth.

more efficient decomposition and potential for higher nutrient mineralization. Conversely, treatments with a higher C: N ratio (control) suggest slower organic matter breakdown and nutrient release. Overall, in all treatments, compost maturity was obtained after 60 days of decomposition except for control (25). The C: N ratio is considered one of the simple indices used to evaluate any organics for its suitability for soil application and it is the index traditionally used to establish the maturity degree of the compost (26). The integrated system of composting with bioinoculants followed by vermicomposting in recent years has gained worldwide attention to overcome the problem of the degradation of lignocellulosic waste of various agricultural residues and waste industrial by-products (27). In another study, sugarcane trash treated with EM-F exhibited C: N ratio value (14.44:1) on 60 days of composting (28). Effective microorganisms extract (EM), yeast (*Saccharomyces*), actinomycetes (*Streptomyces*), fungi (*Aspergillus*); EM-F rich in lactic acid bacteria (*Lactobacillus*) were used in the above study (29).

Table 2. Effect of Trash D on sugarcane trash *in-situ* decomposition and its impact on C: N ratio

Treatments	Total carbon (%)		Total nitrogen (%)	
	30 th day	60 th day	30 th day	60 th day
T ₁	28.54	21.97	0.84	1.11
T ₂	26.2	21.24	0.92	1.20
T ₃	28.97	22.91	0.88	1.16
T ₄	25.37	21.56	0.9	1.22
T ₅	29.74	22.51	0.89	1.08
T ₆	28.94	21.78	0.85	1.13
T ₇	33.67	25.89	0.73	0.92
SE±	0.69	0.59	0.02	0.02
CD (0.05)	1.49	1.31	0.04	0.04

Yield and quality parameters

Yield was highest in Trash D at 30 kg/ha + 125% RDF (137.1 t/ha), followed by Trash D at 30 kg/ha + 100% RDF (135.2 t/ha). Control recorded the lowest yield (115 t/ha). Trash D treatments also improved cane height, girth and CCS %, aligning with previous studies (30, 31). The research data highlight that trash mulching had additional benefits, including moisture conservation, weed control, enhanced soil biological activity and increased number of earthworms, which finally leads to higher yield. It also found that garbage mulching enhances sugarcane productivity and soil health (32).

Table 3. Effect of TNAU Trash D on *in-situ* sugarcane trash decomposition and yield attributes of sugarcane

Treatments	Tiller population ('000/ha)	Yield parameters					CCS (%)
		Cane height (cm)	No. nodes / cane	Cane girth (cm)	Individual cane weight (kg)	Cane yield (t/ha)	
T ₁	181.5	210	20	2.30	1.88	126.9	11.4
T ₂	186.1	208	22	2.88	1.99	137.1	12.2
T ₃	181.5	202	20	2.36	1.82	126.5	11.6
T ₄	184.4	204	23	2.56	1.87	135.2	12.0
T ₅	182.3	200	20	2.48	1.76	120.2	11.6
T ₆	183.5	203	22	2.49	1.80	129.4	11.4
T ₇	166.7	189	18	2.17	1.72	115.0	10.2
SEd	3.15	3.86	0.31	0.05	0.04	1.57	NS
CD (0.05)	6.86	8.40	0.67	0.10	0.09	3.43	-

Conclusion

This study demonstrates that microbial decomposition using trash decomposer (Trash D) at 30 kg/ha, combined with balanced fertilization, significantly enhances soil health and sugarcane yield. Trash D treatments at 30 kg/ha + 100% RDF and at 30 kg/ha + 125% RDF improved decomposition efficiency, reduced the C: N ratio and increased crop performance, confirming the benefits of integrating microbial technology in sustainable agriculture. Compost maturity was achieved within 60 days in all Trash D-treated plots, enhancing soil nutrient dynamics. The significant improvements in yield and soil fertility observed in T₂ (30 kg Trash D + 125% RDF) and T₄ (30 kg Trash D + 100% RDF) highlight the potential of microbial decomposition as an eco-friendly alternative to traditional trash management methods. Future studies should optimize microbial formulations for different soil types and climates. Large-scale field trials and economic analyses will further validate the feasibility of microbial trash decomposers in sustainable sugarcane production systems.

Acknowledgements

The authors kindly acknowledge with special thanks the approval of the University research project by Tamil Nadu Agricultural University and the successful research project at Sugarcane Research Station, Cuddalore, Tamil Nadu, India.

Authors' contributions

MP, GG worked on conceptualization. MP, GG and SD on formal analysis. Funding acquisition was done by MP, GG, KP and GG. Original draft was written by GG, MPS, KP, KN. PG, MPS, GG, SD, KN contributed to writing, reviewing and editing of the manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None

Reference

- Bernabe GA, Almeida S, Ribeiro CA, Crespi MS. Evaluation of organic molecules originated during composting process. *J Thermal Anal Calorimetry*. 2011;106(3):773–78. <https://doi.org/10.1007/s10973-011-1420-1>
- Liu H, Li D, Huang Y, Lin Q, Huang L, Cheng S, et al. Addition of bacterial consortium produced high-quality sugarcane bagasse compost as an environmental-friendly fertilizer: Optimizing arecanut (*Areca catechu* L.) production, soil fertility and microbial community structure. *Appl Soil Ecol*. 2023;188:104920. <https://doi.org/10.1016/j.apsoil.2023.104920>
- Powar RV, Mehetre SA, Powar TR, Patil SB. End-use applications of sugarcane trash: A comprehensive review. *Sugar Tech*. 2022;24(3):699–714. <https://doi.org/10.1007/s12355-022-01107-5>
- Ikram K, Niaz Y, Atif M, Mansha MZ, Nadeem M, Ghani MU, et al. Optimizing mechanical and crop physical factors for sugarcane trash recovery at farm. *Pak J Agri Sci*. 2021;58(4): 1387–1393. <https://doi.org/10.21162/PAKJAS/21.1541>
- Kumar N, Rana L, Singh AK, Pramanick B, Gaber A, Alsulhaibani AM, et al. Precise macronutrient application can improve cane yield and nutrient uptake in widely spaced plant-ratoon cycles in the Indo-Gangetic plains of India. *Front Sustain Food Syst*. 2023;7:1223881. <https://doi.org/10.3389/fsufs.2023.1223881>
- Yang X, Soothar RK, Rahu AA, Wang Y, Li B, Mirjat MU, et al. Integrated effects of water stress and plastic film mulch on yield and water use efficiency of grain maize crop under conventional and alternate furrow irrigation method. *Water*. 2023;15(5):924. <https://doi.org/10.3390/w15050924>
- de Aquino GS, de Conti Medina C, da Costa DC, Shahab M, Santiago AD. Sugarcane straw management and its impact on production and development of ratoons. *Ind Crops Prod*. 2017;102:58–64. <https://doi.org/10.1016/j.indcrop.2017.03.018>
- Saravanan P, Palanisamy K, Kulandaivelu S. Spectroscopic assessment of sugarcane bagasse mediated vermicompost for qualitative enrichment of animal wastes *Elephas maximus* and *Bos taurus*. *Waste Biomass Valor*. 2023;14(7):2133–49. <https://doi.org/10.1007/s12649-022-02011-5>
- Aranganathan L, Rajasree SR, Suman TY, Remya RR, Gayathri S, Jayaseelan C, et al. Comparison of molecular characteristics of Type A humic acids derived from fish waste and sugarcane bagasse co-compost influenced by various alkaline extraction protocols. *Microchem J*. 2019;149:104038. <https://doi.org/10.1016/j.microc.2019.104038>
- Bassey MS, Kolo MG, Daniya E, Odofin AJ. Trash mulch and weed management practice impact on some soil properties, weed dynamics and sugarcane (*Saccharum officinarum* L.) genotypes plant crop productivity. *Sugar Tech*. 2021;23(2):395–406. <https://doi.org/10.1007/s12355-020-00899-8>
- Dlamini NE, Zhou M. Soils and seasons effect on sugarcane ratoon yield. *Field Crops Res*. 2022;284:108588. <https://doi.org/10.1016/j.fcr.2022.108588>
- Devi GK, Vignesh K, Chozhavendhan S. Effective utilization of sugarcane trash for energy production. In: Kumar RP, Gnansounou E, Raman JK, Baskar G, editor. Refining biomass residues for sustainable energy and bioproducts. New York: Academic Press; 2020. p. 259–73 <https://doi.org/10.1016/B978-0-12-818996-2.00012-0>

13. Walkley A, Black IA. An examination of the Degjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci.* 1934;37:29–38. <https://doi.org/10.1097/00010694-193401000-00003>
14. Piper CS. *Soil and plant analysis*. Hans Publishers, Bombay; 1966
15. Panse VG, Sukhatme PV. *Statistical methods for agricultural workers*. By. New Delhi: Indian Council of Agricultural Research; 1961
16. Ashagrie Y, Zech W, Guggenberger G, Mamo T. Soil aggregation and total and particulate organic matter following conversion of native forests to continuous cultivation in Ethiopia. *Soil Tillage Res.* 2007;94(1):101–08. <https://doi.org/10.1016/j.still.2006.07.005>
17. Ranjan S, Kumar S, Dutta SK, Sow S, Kumar S, Sushant. Long-term organic amendment application improves soil fertility status, nutrient accumulation and crop productivity under rice-wheat cropping system. *Commun Soil Sci Plant Anal.* 2023;54(18):2579–89. <https://doi.org/10.1080/00103624.2023.2227240>
18. Prosdocimi M, Tarolli P, Cerda A. Mulching practices for reducing soil water erosion: A review. *Earth-Sci Rev.* 2016;161:191–203. <https://doi.org/10.1016/j.earscirev.2016.08.006>
19. Laik R, Singh SK, Pramanick B, Kumari V, Nath D, Dessoky ES, et al. Improved method of boron fertilization in rice (*Oryza sativa* L.)-mustard (*Brassica juncea* L.) cropping system in upland calcareous soils. *Sustain.* 2021;13(9):5037. <https://doi.org/10.3390/su13095037>
20. Dashtban M. Fungal bioconversion of lignocellulosic residues; Opportunities and perspectives. *Int J Biol Sci.* 2009;5:578–95. <https://doi.org/10.7150/ijbs.5.578>
21. Shan G, Lu H, Li Q. The properties and dynamic changes of DOM subfractions during food waste and sugarcane leaves co-composting. *Environ Sci Pollut Res.* 2018;25:7433–42. <https://doi.org/10.1007/s11356-017-1083-y>
22. Xu J, Lu Y, Shan G, He XS, Huang J, Li Q. Inoculation with compost-born thermophilic complex microbial consortium induced organic matters degradation while reduced nitrogen loss during co-composting of dairy manure and sugarcane leaves. *Waste Biomass Valor.* 2019;10:2467–77. <https://doi.org/10.1007/s12649-018-0293-y>
23. Bhattacharyya P, Bhaduri D, Adak T, Munda S, Satapathy BS, Dash PK, et al. Characterization of rice straw from major cultivars for best alternative industrial uses to cutoff the menace of straw burning. *Ind Crops Prod.* 2020;143:111919. <https://doi.org/10.1016/j.indcrop.2019.111919>
24. Tegene S, Terefe H, Tena E, Dejene M, Tegegn G, Ayalew A. Sugarcane Smut (*Sporisorium scitamineum*), a major threat to sugarcane production: designing sustainable management option for sugar and cane productivity in Ethiopia. *Biocontrol Sci Technol.* 2024;34(10):885–914. <https://doi.org/10.1080/09583157.2024.2391322>
25. Shan G, Xu J, Jiang Z, Li M, Li Q. The transformation of different dissolved organic matter subfractions and distribution of heavy metals during food waste and sugarcane leaves co-composting. *Waste Manag.* 2019;87:636–44. <https://doi.org/10.1016/j.wasman.2019.03.005>
26. Towles TB, Catchot AL, Gore J, Cook DR, Caprio MA, Daves C, et al. Determining yield effects of simulated stand loss in field corn (*Zea mays*). *Midsouth Entomol.* 2022;15:1–9.
27. Janssen VC, van Rijn F. Towards a sustainable sugarcane industry in India: improving livelihoods and increasing water use efficiency in sugarcane growing in India through adoption of improved practices and technologies: evaluating progress between 2016 and 2021. *Wageningen Economic Res;* 2022. <https://doi.org/10.18174/576255>
28. Phukongchai W, Kaewpradit W, Rasche F. Inoculation of cellulolytic and ligninolytic microorganisms accelerates decomposition of high C/N and cellulose rich sugarcane straw in tropical sandy soils. *Appl Soil Ecol.* 2022;172:104355. <https://doi.org/10.1016/j.apsoil.2021.104355>
29. Shukla SK, Jaiswal VP, Sharma L, Tiwari R, Pathak AD, Gaur A, et al. Trash management and *Trichoderma harzianum* influencing photosynthesis, soil carbon sequestration and growth and yield of sugarcane ratoon in subtropical India. *Eur J Agron.* 2022;141:126631. <https://doi.org/10.1016/j.eja.2022.126631>
30. Kumar N, Sow S, Rana L, Ranjan S, Singh AK. Trash amended with *Trichoderma* effects on cane yield, soil carbon dynamics and enzymatic activities under plant-ratoon system of sugarcane in calcareous soil. *Sugar Tech.* 2024;1–4. <https://doi.org/10.1007/s12355-024-01467-0>
31. Matoso ES, Reis VM, Giacomini SJ, Silva MT, Avancini AR, Silva SD. Diazotrophic bacteria and substrates in the growth and nitrogen accumulation of sugarcane seedlings. *Sci Agric.* 2020;78(1):e20190035. <https://doi.org/10.1590/1678-992X-2019-0035>
32. Bairwa R, Jha CK, Thakur SK, Mamta, Roy DK, Brajendra. Carbon pools and indices under activated trash treatments in sugarcane plant-ratoon system grown in calcareous soil of subtropics. *Sugar Tech.* 2023;25(6):1433–44. <https://doi.org/10.1007/s12355-023-01280-1>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc
See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.