



RESEARCH ARTICLE

# Molecular and morphological identification of some oligochaete species as indicator for ecological diversity of the Hilla River in Babylon Province-Iraq

Abd Al-Rezzaq AJ & AL- Yasari WAK\*

Department of Biology, College of Science, University of Babylon, Babylon 00964, Iraq

\*Email: [sci.wameedh.adil@uobabylon.edu.iq](mailto:sci.wameedh.adil@uobabylon.edu.iq)



## ARTICLE HISTORY

Received: 03 February 2025

Accepted: 04 March 2025

Available online

Version 1.0 : 04 April 2025

Version 2.0 : 15 April 2025



Check for updates

## Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonepublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonepublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See [https://horizonepublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

## CITE THIS ARTICLE

Abd Al - Rezzaq AJ & AL - Yasari WAK. Molecular and morphological identification of some oligochaete species as indicator for ecological diversity of the Hilla River in Babylon Province-Iraq. Plant Science Today. 2025; 12(2): 1-8. <https://doi.org/10.14719/pst.7578>

## Abstract

The present study was carried out in Hilla river for the identification of Oligochaetes using morphological and molecular methods. The morphological study confirms the identification of five species (*Aulodrilus pigueti*, *Branchiura sowerbyi*, *Eiseniella tetraedra*, *Limnodrilus hoffmeisteri*, *Stylaria lacustris*) using scanning electron microscope (SEM) technique. Out of all species, *Eiseniella tetraedra* received the greatest percentage of density (55.19 %). Their diversity was assessed using several indices. Value of Shannon-Wiener index ranged between 0.23-0.93 and that of, Simpson index ranged between 0.32-0.91. The results of these and other biodiversity indicate the presence of environmental stress. Mitochondrial cytochrome c oxidase subunit I (COI) region was amplified and sequenced to find the similarity and variations among species and confirm the diagnosis of species (*Paranais litoralis*, *Paranais frici*, *Stylaria lacustris*, *Monopylephorus rubroniveus*, *Tubifex tubifex*) and all sequence and strains matched to the complete genetic content of the samples at the NCBI to ensure the results. This study is considered the first study for the diagnosis of oligochaetes in the middle Euphrates region by using the molecular technique to confirm the identification of numerous unrecorded species that share morphological characteristics with other species, making it challenging to categorize them morphologically.

## Keywords

biodiversity indices; DNA barcoding; Hilla river, oligochaetes

## Introduction

Biodiversity is studied at three levels, genetic diversity, species diversity and ecosystem diversity (1). Mostly, morphological species richness is used as the mine scale for measuring biodiversity (2). Despite morphological importance, many studies and research have clarified that the species concepts dependent on the shape and morphology have miscalculated how crucial biodiversity is to ecological systems (3). Other studies found that genetic diversity, especially freshwater organisms, has given new concepts to diversity and recording species through their genetic content and species delimitation (4). This raises many concerns about the true number of species on earth and the true diversity of living organisms (5), as well as the influence of geology and geography on the spread and diversity of species (6). Few researchers have reported cryptic species, which are similar morphologically but different genetically using molecular techniques (7).

Recent studies of environmental pollutants have confirmed that measures of biodiversity and its indices are indicators for pollution monitoring

and environmental changes (8). Some diversity indices, such as Shannon diversity index, Dominance index (Simpson index), Relative abundance and Uniformity index, were used in this study (36). Since the beginning of interest in measuring biodiversity, the number of species and its changes have been considered biodiversity (9), which consists of two components, species richness and evenness (10) and can be regarded as factors of sustainability and healthiness for most ecosystems (11). Diversity is affected by many factors, including human activity, sampling size, farming, rivers, underground waters and the aquatic environment (12). Most freshwater invertebrates are characterized by the small size and complexity of their life cycle and contain several life stages, so it is often difficult to rely on morphological identification in biomonitoring, classification and biodiversity (13). Furthermore, when classifying to the species level, the adult stage is required and it is impossible to rely on the larval or immature stages to identification (14). Modern techniques such as gene sequencing are very successful in the genetic diagnosis of many species of aquatic invertebrates (15). In addition, through genetic biodiversity, it is possible to evaluate freshwater quality through biomonitoring of freshwater invertebrates (16).

Oligochaetes are the most diverse and abundant group in freshwater habitats (17). Aquatic oligochaetes are very small, ranging from less than 0.5 mm up to 28 centimeters in length, although most are smaller than 3 cm (18). More than 1700 species of aquatic oligochaetes have been identified and about 1100 species inhabit freshwater. The Tubificidae is the most diverse group, with more than 1000 species described; most of these species inhabit freshwater and about 60 species of megadriles are also considered aquatic habitats (19).

Limited studies in Iraq concerning the annelid fauna, especially oligochaeta, have been conducted. Three new tubificid species (*Limnodrilus profundicola*, *Embolacephalus velutinus* and *Aulodrilus pigueti*) were identified in the marsh in southern of Iraq (20). Tubificid worms such as *Tubifex tubifex*, *Limnodrilus claparedianus*, *Limnodrilus profundicola*, *Branchiura sowerbyi*, *Limnodrilus maumeensis* and *Limnodrilus hoffmeisteri* have also been recorded in various aquatic habitats (21-26). Most studies on the oligochaetes have been focused on Naididae fauna, 37 naidid species have been previously reported (27-30). The aquatic oligochaetes community in AL-Abbasyia River (Al-Najaf province) identified three families Naididae, Lumbriculidae and Enchytraeidae (31). Also, the studies of genetic variation between and within species at a molecular level are very few; The genetic techniques for molecular identification and classification of species (Annelida; Oligochaeta) by using DNA barcoding in Kurdistan-Iraq. By using morphological and molecular methods, *Limnodrilus claparedianus* was identified (32).

Finally, the morphological identification of oligochaetes to the species level depends on the arrangement of the chaetae, located dorsally on the prepatellar segments, but identification of some species needs to be done at mature stage, as their classification depends on the reproductive system. Therefore, our study focused on molecular analyses and Scanning Electron Microscope (SEM) as modern techniques for identification of the oligochaete species.

## Materials and Methods

### Collection of samples

The study samples were collected seasonally (Spring 2021 to Winter 2022) from Hilla river. Five sites were selected, including three stations from Hilla river and two streams branching from the river. The first site (St1) represents the Hilla river; It is situated at the river's entrance to the village of Anana (32°33'6.61"N; 44°25'14.22"E). The second site (St2) represents the Hilla river before entering the city center (Betta bridge) (32°31'1.35"N; 44°25'37.40"E). The third site (St3) depicts the Hilla river immediately following its departure from the governorate's center in Al-Farsi region (32°28'7.31"N; 44°26'23.79"E). The fourth site (St4) is a stream that arises from the Hilla river and flows through the village of Al-Ghalis (32°26'15.59"N; 44°26'53.56"E). The fifth site (St5) is a stream that splits off from the Hilla river near Nazim Dora and travels through the region between Al-Dolab and Al-Dabla (32°25'4.08"N; 44°29'14.67"E) (Fig. 1). An Ekman Birge grab sampler was used to collect sediment samples during the study period. Three replicates for each sample were brought to the laboratory and formalin (4 %) and ethanol (75 %) were used to fix and preserve the worms (34).

### Analysis of diversity indices

Analysis of diversity includes the calculation of Relative abundance (35), Shannon index, Uniformity index and Simpson index (36).

### Morphological identification

Scanning Electron Microscope (SEM) type (INSPECT S50) was used for identification of the specimens; the sample was coated with gold. Species identification was conducted using appropriate keys (37-42).

### Molecular identification

**DNA extraction:** Cetyl trimethyl ammonium bromide "CTAB" Kit was used to extract genomic DNA from samples following manufacturers protocol provided by Macrogen company. Since the study was conducted to obtain genetic diversity among oligochaetes, 18s rRNA were selected for this study (32).

**Preparation of primers:** Primers were obtained from <https://www.ncbi.nlm.nih.gov/> and designed using use free online tool <https://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi/>. The lyophilized primers were sourced from Macrogen company, List of primers used in this study are shown in Table 1.

**Polymerase chain reaction (PCR):** Conventional PCR was used to amplify the target DNA using specific primer pairs. It includes three consecutive steps that are repeated for specific number of cycles to get PCR product which could be finally visualized after agarose gel electrophoresis. The thermal cycling conditions are mentioned in Table 2. PCR products were sent to Macrogen company in Korea to be matched to the complete genetic content of the samples at NCBI to ensure the sequencing results.



**Fig. 1.** Maps of the Hilla river show the studied sites.

**Table 1.** Primer used for characterization of oligochaetes species in the current study

Primer	Sequence 5' to 3'	Product Size (base pairs)	Annealing temperature
Primer- F	GGGCGTAATGAAAGTGAAGG	500	56 °C
Primer- R	TAACCGGACGTTTGGTTCAT		

## Results and Discussion

A total of 683 oligochaeta individuals belonging to five species were reported. The five identified species include *Aulodrilus pigueti* (5), *Branchiura sowerbyi* (178), *Limnodrilus hoffmeisteri* (9), *Stylaria lacustris* (114) and *Eiseniella tetraedra* (377). These represented 0.73 %, 26.06 %, 1.3 %, 16.78 % and 55.19 % of the total sample respectively. These results represent a new record for Hilla river (Fig. 2, Table 3). Regarding their distribution throughout the study sites and periods, the highest number (88) of individuals recorded in St1 during spring, while the lowest number (5) recorded in St5 during summer. As can be seen from (Table 3), it turned out that the decline in the numbers of benthic invertebrates coincided with the rise in temperatures during the hottest season of the year (43 & 44). *Eiseniella tetraedra* was the most abundant species in majority of the sites across seasons (45).

Previously, reliance was placed on shape and phenotypic characteristics to identify and classify species. However, now the combined use of molecular techniques and morphology has been preferred for the identification of the species. Also, species identification using DNA technology has helped diagnose the young individuals of these species at different ages, which was previously possible only at adult stages. This is the first DNA molecular study in Babylon province to identify and classify the oligochaetes species by DNA barcoding using the COX1 gene

and find the phylogenetic relationship among species. The Basic Local Alignment Search Tool (BLAST) in NCBI is used for nucleotide sequence comparison. In addition, the information obtained from GenBank was used to compare sequences of different species and draw a phylogenetic tree (Fig. 3). According to the phylogenetic tree, the results of sample 1 were two species belonging to the Naididae family and sample 2 had five species belonging to the Naididae family too. BLAST similarity analysis showed that the obtained sequences have 99 - 100 % bootstrapping with the sequences in GenBank. The current study focused on the molecular identification of oligochaetes for many reasons, frequently has intricate life cycles, dispersal abilities and quick local adaptations, all of which may promote intraspecific divergence and interspecific gene flow (46).


The values of the diversity indexes were calculated for the diagnosed species isolated from the stations. According to the Shannon-Wiener index, the present study indicated that the (St4) recorded the highest value (0.93) in the summer and the (St5) recorded the lowest value (0.17) in the autumn (Table 4). This index's values dipped in the autumn, perhaps as a result of increased turbidity and suspended debris that affects the variety of food (47).

The Shannon-Wiener index estimates the diversity of species within a community, its high values are evidence of high diversity (48) and a value higher than 3 refers to high diversity of a healthy community living in a good

**Table 2.** PCR conditions for the current study

Primer	Sequence 5' to 3'	Product Size (base pairs)	Conditions steps (temperature, time)
Primer- F	GGGCGTAATGAAAGTGAAGG	500	1: 95 C°, 2 min. 2: 95 C°, 30 sec. 3: 56 C°, 30 sec.
Primer- R	TAACCGGACGTTTGGTTCAT		4: 72 C°, 20.0 sec. 6: 72 C°, 5 min. 7: 4 C°, forever






**Fig. 2.** SEM pictures of the identified species : A-B, *Aulodrilus pigueti* ; C-D, *Branchiura sowerbyi*; E-F, *Limnodrilus hoffmeisteri*; G-H, *Stylaria lacustris*; I-J, *Eiseniella tetraedra* .

**Table 3.** Seasonal changes in the numbers and percentage of the recorded species during the study period

Species	St 1 (village of Anana)				St 2(Betta Bridge)				St 3 (Al-Farsi region)				St 4 (village of Al-Ghalis)				St 5 (Nazim Dora)				Total No.	Percentage %
	Spring	Summer	Autumn	Winter	Spring	Summer	Autumn	Winter	Spring	Summer	Autumn	Winter	Spring	Summer	Autumn	Winter	Spring	Summer	Autumn	Winter		
<i>Aulodrilus pigueti</i>	0	0	0	0	0	0	0	0	3	0	0	0	0	2	0	0	0	0	0	0	5	0.73
<i>Branchiura sowerbyi</i>	28	5	8	0	15	4	0	10	26	10	9	4	19	6	0	4	12	5	0	13	178	26.06
<i>Limnodrilus hoffmeisteri</i>	0	0	0	3	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	5	9	1.3
<i>Stylaria lacustris</i>	17	8	9	3	0	7	9	5	0	11	12	0	0	0	8	6	5	4	2	8	114	16.78
<i>Eiseniella tetraedra</i>	43	15	17	20	46	24	19	13	43	23	12	3	35	10	6	7	39	0	22	0	377	55.19
Total number of Individuals	88	28	34	26	61	35	28	28	72	44	33	8	54	18	14	17	56	5	24	26	683	
number of Species	3	3	3	3	2	3	2	3	3	3	3	3	2	3	2	3	3	2	2	3		

**Fig. 3.** Phylogenetic tree (according to BLAST) of the studied species.**Table 4.** Seasonal changes in biodiversity indices for the study stations during the study period

stations	Seasons	Simpson Index	Shannon Wiener index	Relative abundance (%)	Uniformity index
St1	Spring	0.5	0.73	12.8	0.66
	Summer	0.54	0.72	4.09	0.66
	Autumn	0.5	0.76	4.97	0.69
	Winter	0.82	0.36	3.8	0.33
St2	Spring	0.87	0.23	8.93	0.33
	Summer	0.69	0.54	5.12	0.49
	Autumn	0.8	0.32	4.09	0.46
	Winter	0.44	0.82	4.09	0.75
St3	Spring	0.48	0.8	10.54	0.73
	Summer	0.53	0.72	6.44	0.66
	Autumn	0.44	0.83	4.83	0.76
	Winter	0.32	0.69	1.17	0.63
St4	Spring	0.9	0.19	7.9	0.27
	Summer	0.39	0.93	2.63	0.85
	Autumn	0.71	0.41	2.04	0.59
	Winter	0.4	0.88	2.48	0.8
St5	Spring	0.6	0.63	8.19	0.57
	Summer	0.66	0.45	0.73	0.65
	Autumn	0.91	0.17	3.51	0.24
	Winter	0.74	0.48	3.8	0.44

environment, while when the Shannon-Wiener value lower than 1 meaning the presence of environmental change which leads to the disappearance of sensitive species and their migration (44), so, worms cannot grow in the current unstable environment due to stresses that may be brought on by pollution. The seasonal changes lead to changes in diverse values, which in turn affects the nature of the life cycle of each species and our results agree with previous reports (49).

The dominance (Simpson index) was measured in the current study and shows a high value (0.91) in (St5) during autumn and a low value (0.32) in (St3) during winter (Table 4). These findings showed that the communities are stable and mature because many different species share dominance. It is acknowledged that low diversity, which typically displays values near zero, indicates that the communities are under stress (50). The current study's findings demonstrated that the values of the Uniformity index fluctuated between a low value (0.24) in (St5) in autumn and a high value (0.85) in (St4) during summer, as shown in Table 4. The decrease in the values of the Uniformity index indicates the predominance of a few species with high densities, may be due to the presence of environmental pressure or some species being more competitive (51, 53).

The relative abundance index's value varied from a low value (0.73) in (St5) during summer to a high value (12.8) in St1 (Table 4). The results indicate that the current species were rare in all stations and seasons because their relative abundance was less than 10 % (52).

## Conclusion

The biodiversity of oligochaetes fluctuated throughout the Hilla river among all seasons and we noticed there were little species richness and evenness of oligochaetes according to biodiversity indices values. Regarding the recorded species, there were four species (*Aulodrilus pigueti*, *Branchiura sowerbyi*, *Limnodrilus hoffmeisteri*, *Stylaria lacustris*) with low density and individual numbers while *Eiseniella tetraedra* had a good density during the study period may be more tolerant to environmental conditions. Ultimately, there was a decrease in diversity, which suggests that the river water may be under environmental stress due to changes in its chemical and physical characteristics which lead to the presence of pollution. We recommend reliance on molecular methods and SEM to prove the identification of the oligochaetes.

## Acknowledgements

The authors express their gratitude for the support from Department of Biology, College of Science, University of Babylon/ Iraq.

## Authors' contributions

AJAAR and WAKAY designed and performed the experiment. AJAAR analyzed data and wrote the first draft of manuscript; WAKAY performed statistical analysis and revised the manuscript. Both authors read and approved the final manuscript.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interest to declare.

**Ethical issues:** None

## References

- Spangenberg JH. Biodiversity pressure and the driving forces behind. *Ecological Economics*. 2007;61(1):146-58. <https://doi.org/10.1016/j.ecolecon.2006.02.021>
- De Queiroz K. Species concepts and species delimitation. *Systematic Biology*. 2007;56(6):879-86. <https://doi.org/10.1080/10635150701701083>
- Monro AK, Mayo SJ (eds). *Cryptic species: Morphological stasis, circumscription and hidden diversity*. Cambridge University Press. Cambridge. Chapter 4. 2022. <https://doi.org/10.1017/9781009070553>
- White BP, Pilgrim EM, Boykin LM, Stein ED, Mazor RD. Comparing four species-delimitation methods applied to a DNA barcode data set of insect larvae for use in routine bioassessment. *Freshwater Science*. 2014;33(1):338-48. <https://doi.org/10.1086/674982>
- Zachos FE, Apollonio M, Bärmann EV, Festa-Bianchet M, Göhlich U, Habel JC, et al. Species inflation and taxonomic artefacts -A critical comment on recent trends in mammalian classification. *Mammalian Biology*. 2013;78(1):1-6. <https://doi.org/10.1016/j.mambio.2012.07.083>
- Spitzer B. Genetic structure in populations of *Baetis tricaudatus* in the San Bernardino mountains. *Western North American Naturalist*. 2014;74(4):434-45. <https://doi.org/10.3398/064.074.0409>
- Bickford D, Lohman DJ, Sodhi NS, Ng PKL, Meier R, Winker K, et al. Cryptic species as a window on diversity and conservation. *Trends in Ecology and Evolution*. 2007;22(3):148-55. <https://doi.org/10.1016/j.tree.2006.11.004>
- Zaghloul A, Saber M, Gadow S, Awad F. Biological indicators for pollution detection in terrestrial and aquatic ecosystems. *Bulletin of the National Research Centre*. 2020;44(1):127. <https://doi.org/10.1186/s42269-020-00385-x>
- Williams J. Metrics for assessing the biodiversity values of agricultural landscapes. *Pacific Conservation Biology*. 2004;10(3):145-63. <https://doi.org/10.1071/PC040145>
- Magurran AE. *Measuring biological diversity*. Wiley-Blackwell Publishing, Oxford. Edition 1. 2004:256.
- Simba DL, Beest M, Hawkins HJ, Larson KW, Palmer AR, Sandström C, et al. Wilder rangelands as a natural climate opportunity: Linking climate action to biodiversity conservation and social transformation. *Ambio*. 2024;53(5):678-96. <https://doi.org/10.1007/s13280-023-01976-4>
- Afshani SA, Norani M, Mahdavi M, Norian A. Applicable authority SPSS17. *Bishe Application*. 2009:437.
- Strachan SR, Chester ET, Robson BJ. Freshwater invertebrate life history strategies for surviving desiccation. *Springer Sci Rev*. 2015;3(1):57-75. <https://doi.org/10.1007/s40362-015-0031-9>
- Stein ED, White BP, Mazor RD, Jackson JK, Battle JM, Miller PE, et al. Does DNA barcoding improve performance of traditional stream



- bioassessment metrics? *Freshwater Biology*. 2014;33(1):302–11. <https://doi.org/10.1086/674782>
15. Webb JM, Jacobus LM, Funk DH, Zhou X, Kondratieff B, Geraci CJ, et al. A DNA barcode library for North American Ephemeroptera: progress and prospects. *PLoS One*. 2012;7(5):e38063. <https://doi.org/10.1371/journal.pone.0038063>
  16. Carter JL, Resh VH. Analytical approaches used in stream benthic macroinvertebrate biomonitoring programs of state agencies in the United States. 2013. <https://doi.org/10.3133/ofr20131129>
  17. Atanacković A, Popović N, Marinković N, Duknić J, Stanković J, Paunović M. Effects of environmental factors on the distribution and diversity of aquatic oligochaetes, *Water* (Basel). 2023;15 (22):3873. <https://doi.org/10.3390/w15223873>
  18. Krieger KA, Stearns AM. Atlas of the aquatic oligochaete worms (Phylum Annelida: Class Clitellata: Superorder Microdrili) recorded at the old woman Creek National Estuarine Research Reserve and State Nature Preserve, Ohio National Center for Water Quality Research Heidelberg University Tiffin, Ohio, USA.2010. [www.Heidelberg.edu/wql/educationoutreach/reports](http://www.Heidelberg.edu/wql/educationoutreach/reports)
  19. Martin P, Martinez-Ansemil E, Pinder A, Timm T, Wetzel MJ. Global diversity of oligochaetous clitellates (Oligochaeta: Clitellata) in freshwater animal diversity assessment. *Hydrobiology*. 2008;595 (1):117-27. <https://doi.org/10.1007/s10750-007-9009-1>
  20. Jaweir HJ. New record of three tubificid species (Annelida: ligochaeta) from Al-Hawiezah Marsh, Iraq. *Mesopot. J. Mar Sci*. 2011; 26(2):114-21. <https://doi.org/10.58629/mjms.v26i2.177>
  21. Jaweir HJ, Al-Lami A, Hassan LA. Study of population structure of *Limnodrilus hoffmeisteri* (Annelida: Oligochaeta) in one of the muddy pond of Al-Zaafarana fish farm. *Journal of College of Education for Women*. 2002;13(4):772-78.
  22. Al-Khafaji NF. A study of population structure and some biological aspects of *Branchiura sowerbyi*, Beddard, Iraq. 2002.
  23. Al-Rubaiy HH. The toxic effects of the herbicide's glyphosate and propanil on *Limnodrilus claparedeanus*, Ratzl, 1868. (Annelida: Oligochaeta) (thesis). Iraq: University of Baghdad. 2002.
  24. Farman KS. Effects of salinity and sediment on toxicity of heavy metals on survival of *Limnodrilus claparedeanus* Ratzl, 1868 (thesis). Iraq: University of Baghdad. 2005.
  25. Ali LA. A study of macroinvertebrates community in the middle sector of greater Zab River (thesis). Iraq: University of Baghdad. 2007.
  26. Sebtie HA. An ecological study of the benthic macroinvertebrates community in the southern marshes of Iraq (thesis). Iraq: University of Baghdad. 2009.
  27. Nashaat MR. Impact of AL-Durah power plant effluents on physical, chemical and invertebrates biodiversity in Tigris River Southern Baghdad (thesis). Iraq: University of Baghdad. 2010.
  28. Al-Abbad MYM. New records of twelve species of oligochaeta (Naididae and Aeolosomatidae) from the Southern Iraqi Marshes, Iraq. *JJBS*. 2012;5(2):105-11.
  29. Jaweir HJ, Al-Janabi ESO. Biodiversity and abundance of aquatic oligochaetes family Naididae in the middle sector of Euphrates river, Al-Mussayab City/ Iraq. *The International Journal of the Environment and Water*. 2012;1(1):122-30.
  30. Radhi MM. Sorting and identification of Naididae species (Annelida-Oligochaeta) from different aquatic sites within Baghdad city (thesis). Iraq: University of Baghdad. 2012.
  31. Al-Ameen NI, Jawair HJ. A study of aquatic oligochaetes community in the upper part of Al- Abbasyia river/ middle of Iraq. *Iraqi Journal of Agricultural Sciences*. 2020;51(3):930-37. <https://doi.org/10.36103/ijas.v51i3.1048>
  32. Othman H.H, Ahmad S.T. Molecular identification of some earthworm species (Annelida; Oligochaeta) in Kurdistan-Iraq. *Zanco J Pure Appl Sci*. 2020;32(6):46-53. <https://doi.org/10.21271/ZJPAS.32.6.5>
  33. Zaar EO, Jaweir HJ. Morphological and molecular identification of *Limnodrilus claparedeanus*, 1862 Species (clitellata: naididae) in Tigris River, Baghdad/Iraq. *Baghdad Science Journal*. 2021;18 (2):0231. <https://doi.org/10.21123/bsj.2021.18.2.0231>
  34. Anderson JT, Zilli FL, Montalto L, Marchese MR, McKinney M, Park YL. Sampling and processing aquatic and terrestrial invertebrates in wetlands. *Wetlands Techniques*. Dordrecht: Springer. 2013;5:143–95. [https://doi.org/10.1007/978-94-007-6931-1\\_5](https://doi.org/10.1007/978-94-007-6931-1_5)
  35. Salahi T, Pourbabaie H, Salahi M, Karamzadeh S. An investigation on plant species composition and diversity in the coniferous and broadleaved plantations: Case study of Bibi Yanlu Forest Park, Astara, Iran. *Biodiversitas*. 2017;18(3):958-63. <https://doi.org/10.13057/biodiv/d180313>
  36. Magurran AE. Ecological diversity and its measurement. *Springer*, Netherlands. 2013.
  37. Brinkhurst RO, Jameison BG. Aquatic oligochaetes of the world, University of Toronto press. Toronto, Canada. 1971:859.
  38. Brinkhurst RO. A guide for the identification of British aquatic oligochaeta. *Freshwater Biological Association, Scientific Publication No. 22*. Kendal, UK, Wilson and Sons. 1971.
  39. Stimpson KS, Klem DJ, Hiltunen JK. A Guide to the freshwater tubificidae (Annelida: Clitellata: Oligochaeta) of North America. EPA. 1982;600/3.82-033.
  40. Brinkhurst RO. Guide to the freshwater aquatic microdrile oligochaetes of North America. *Can Spec Publ Fish Aquat Sci*. 1986;84:259.
  41. Timm T. Guide to freshwater oligochaeta and polychaeta of northern and central Europe. *Lauterbornia*. 2009;66:1-23.
  42. Pinder A. Tools for identifying Australian aquatic oligochaetes of the families Phreodrilidae, Lumbriculidae and Capilloventridae (Clitellata: Annelida). *Museum Victoria Science Reports*. 2013;18:1–20. <https://doi.org/10.24199/j.mvrs.2013.18>
  43. Jaweir HJ, Radhi MM. Naididae (Clitellata: Oligochaeta) and Aeolosomatidae (Polychaeta: Aphanoneura) species associated with aquatic plants in Tigris River, Baghdad-Iraq. *Baghdad Science Journal*. 2013;10(1):116–25. <https://doi.org/10.21123/bsj.2013.10.1.116-125>
  44. Jaweir HJ, Al-Seria MHM. A Study of the benthic invertebrates community in Dalmage Lake - Middle of Iraq. *Iraqi Journal of Science*. 2015;56(3): 2513-22.
  45. Javidkar M, Abdoli A, Ahmadzadeh F, Nahavandi Z, Yari M. Molecular evidence reveals introduced populations of *Eiseniella tetraedra* (Savigny, 1826) (Annelida, Lumbricidae) with European origins from protected freshwater ecosystems of the southern Alborz mountains. *Mari Freshwa Res*. 2020;72(1):44. <https://doi.org/10.1071/MF20004>
  46. Cristescu ME. From barcoding single individuals to metabarcoding biological communities: towards an integrative approach to the study of global biodiversity. *Trends in Ecology & Evolution*. 2014;29 (10):566-71. <https://doi.org/10.1016/j.tree.2014.08.001>
  47. Abdulwahab S, Rabee AM. Ecological factors affecting the distribution of the zooplankton community in the Tigris River at Baghdad region, Iraq. *Egyptian Journal of Aquatic Research*. 2015;41(2):187-96. <https://doi.org/10.1016/j.ejar.2015.03.003>
  48. Burton TM, Uzarski DG, Genet JA. Development preliminary invertebrate index of biotic integrity for Lake Huron coastal wetlands. *Wetlands J*. 1999;19(4):869-82. <https://doi.org/10.1007/BF03161789>
  49. Ibrahim SS. Biological diversity of invertebrates in Al-Dagara and Al-Diwaniya Rivers/Iraq (thesis). Iraq: Al-Qadisiyah University; 2005:1-177(in Arabic).

50. Dash. Fundamentals of Ecology Third Edition, New Delhi: Mc Graw Hill India. 2009.
51. Proto-Neto VF. Zooplankton as bioindicator of environmental quality in the tamandanereff system (Pernambuco-Brazil): Anthropogenic influences and interaction with mangroves (thesis) Brazil: University Bremen. 2003.
52. Omori M, Ikeda T. Methods in marine zooplankton ecology. Wiley and Sons, New York. 1984.
53. Nico E, Stefan S. Earthworms as drivers of the competition between grasses and legumes. Soil Biology and Biochemistry. 2008;40 (10):2650-59. <https://doi.org/10.1016/j.soilbio.2008.07.010>