



REVIEW ARTICLE

Turning trash into treasure: Valorization of fruit waste for sustainable bioeconomy

Divya Peekaa^{1*}, P S Kavitha², K S Vijai Selvaraj³, I Muthuvel⁴, V Veeranan Arun Giridhari⁵ & R Gurusamy⁶

¹Department of Fruit Science, Horticultural College & Research Institute, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

²Directorate of Planning and Monitoring, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

³Vegetable Research Station, Tamil Nadu Agricultural University, Palur 697 102, Tamil Nadu, India

⁴Department of Fruit Science, Horticultural College & Research Institute (W), Tamil Nadu Agricultural University, Trichy 620 009, Tamil Nadu, India

⁵Department of Food Nutrition, Horticultural College & Research Institute, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

⁶Department of Food Process Engineering, Agricultural Engineering College & Research Institute, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

*Correspondence email - kavitha.ps@tnau.ac.in

Received: 16 February 2025; Accepted: 30 June 2025; Available online: Version 1.0: 31 July 2025; Version 2.0: 15 August 2025

Cite this article: Divya P, Kavitha PS, Vijai SKS, Muthuvel I, Veeranan VAG, Gurusamy R. Turning trash into treasure: Valorization of fruit waste for sustainable bioeconomy. Plant Science Today. 2025; 12(sp1): 1-6. <https://doi.org/10.14719/pst.7760>

Abstract

The effective utilization of fruit waste offers a significant opportunity to advance sustainable resource management and reduce environmental burdens. This comprehensive overview brings together current research on the diverse applications of fruit by-products, showcasing their potential across various industries. Fruit waste, comprising peels, seeds and pulp, is highly valuable to the food, pharmaceutical and cosmetic industries due to its rich content of bioactive compounds such as vitamins, fibre and antioxidants. In addition to industrial applications, the integration of fruit waste into agricultural practices, such as compost or biofertilizer, improves soil health and enhances crop productivity. These organic amendments reduce dependence on synthetic inputs and promote environmentally responsible farming practices. Beyond agriculture and industry, fruit waste is increasingly explored for its role in sustainable material and energy development. Notably, its high carbohydrate and cellulose content make it suitable for biofuel production, while its film-forming properties support the creation of biodegradable packaging as a replacement for plastics. Furthermore, processed fruit residues are proving to be effective animal feed components, offering nutritional benefits and lowering feed costs in livestock production. This review stresses the economic and environmental advantages of fruit waste utilization and urges the development of scalable, cost-effective processing technologies to maximize its potential. It underscores the importance of ongoing research and cross-sector collaboration among academia, industry and policymakers to unlock the full value of fruit by-products. Such efforts are essential to advancing circular economy principles and reducing the environmental impact of food production systems.

Keywords: agricultural benefits; economic and environmental impact; innovative products; nutrient-rich by-products; sustainable use of fruit waste

Introduction

A significant proportion, ranging from 25-50 % of fruits and vegetables produced globally is lost along the supply chain from farm to table, a phenomenon widely recognized as post-harvest loss (1). Transport and handling account for approximately 25-30 % of these losses. The fruit juice industry produces 5.5 million tonnes (Mt) of waste, with 20-30 % treated (2). These losses vary by crop, with waste percentages differing for various fruits as detailed in Table 1.

Fruit by-products like peels, seeds and pomace contribute significantly to municipal solid waste. Peels and seeds are often discarded during processing despite their nutritional value, while pomace remains a major by-product of the juice industry (8). Fruit waste, rich in protein, sugars and

Table 1. Different types of fruit wastage and their percentage losses

Fruits	Type of waste	Percentage of losses (%)	References
Mango	Seed, peel	25.51	(3)
Banana	Peel	26.5	(4)
Pineapple	Peel	32.12	(5)
Grapes	Stem, seeds, skin	53.00	(6)
Orange	Peel, seeds	29.00	(7)

minerals, can address feed scarcity in developing countries. It may be fed directly or processed through drying and ensiling (9). Additionally, its phytochemicals, fibers and vitamins make it a nutritious alternative to conventional feed ingredients. Using fruit waste in dairy cow diets enhances milk's fatty acid content without reducing production.

Different wastages in fruit crops

Mango

Mango processing generates significant waste, mainly peels and seeds (10). Its peel makes up 15-20 % of the fruit's weight and seeds 20-60 %, depending on the variety (11-13). Mango peel is rich in bioactive compounds like dietary fiber, vitamins C and E, carotenoids and flavonoids, making it a potential functional food (14). It is also a valuable source of aromatic compounds for flavoring and volatile compound extraction (15).

Gallate and penta-O-galloylglucoside from mango exhibit antioxidant, anti-tumor and antimicrobial properties, highlighting their pharmaceutical potential (16). Mango seed kernels, though low in protein, contain essential amino acids like leucine and lysine (17). Mango peel pectin has superior gelling properties, while its flour is used in various food products, including bread, cookies and jams (16). Additionally, *Pichia pinus* yeast is employed to produce lactic acid and single-cell proteins from mango peel.

Banana

Banana processing generates significant waste, with peels accounting for 40 % of the fruit's weight (18). Banana cultivation also yields large amounts of biomass, with pseudo-stems and peduncles being valuable fiber sources. These fibers are used in industries for paper, cardboard, tea bags, currency notes and polymer composites (19). Rich in cellulose and fiber, banana peels are increasingly studied for their potential use as an alternative flour source, adding value to this by-product (20). Plantain roots and stems are traditionally used to bind leaves that serve as baskets, mats and natural packaging materials (21). Old leaves serve as bunch covers, large leaves as umbrellas and dried leaves as fuel for oyster mushroom cultivation.

Pineapple

Pineapple processing generates by-products like peel and core, rich in fibers, pectins, sugars, proteins, vitamins, minerals and phenolic compounds such as ferulic acid and catechin (22, 23). Pineapple peel contains 6.5-35 % hemicellulose, 12-50 % cellulose and 5-30 % lignin, while the core has 41.1-43.5 % cellulose, 5.78 % lignin and 28.5 % hemicellulose (24). The crown has 67.1-85 % cellulose, 13.8 % lignin and 21-21.8 % hemicellulose. These by-products are being explored for eco-friendly biodiesel production, offering economic and environmental benefits.

Citrus

Citrus processing produces various wastes, including solid waste (peels, seeds, rags) and liquid waste (wash water, can cooler overflows and effluents) (25). Citrus peel, a byproduct of juice production, is rich in phenols and flavonoids with antioxidant and health benefits, but lacks commercial value. This study focuses on citrus extracts, particularly Eco-enzyme, known for its phenolic compounds and biological effects (26). Orange peel essential oils mainly consist of limonene (94 %), with smaller amounts of myrcene, linalool and other compounds. Limonene is known for its insecticidal properties (27). Citrus seed processing yields oil (20-40 % of seed weight) and seed meal, with the oil being rich in unsaturated fatty acids, especially linoleic acid, along with oleic, stearic, palmitic

and α -linolenic acids (28).

Jackfruit

Jackfruit waste includes unused peels and seeds, which are rich in pectin, fiber and starch (29, 30). While edible, jackfruit seeds have a short shelf life, but drying them deactivates antinutritional compounds, making them suitable for flour production.

Apple

Apple pomace, a byproduct of apple processing, is rich in carbohydrates, fiber, vitamins and antioxidants, with polyphenols concentrated in the peels (9). It contains 10-15 % pectin and significant amounts of insoluble and soluble fiber (31). Additionally, waste from apple orchards, including leaves and branches, can be processed into biofuels like hydrogen, ethanol and methane (32).

Degradable packaging materials

The increased use of plastic has harmed the environment and animal welfare (33). Developing biodegradable packaging from renewable sources like starch, cellulose and pectin derived from fruit peels and seeds is essential for replacing plastic materials (34).

Biodegradable packaging research can be categorized into four types: 1) Anti-corrosive for durable metals, 2) Pharmaceutical for packaging medical items, 3) Electronic packaging for protecting components and 4) Food packaging to prevent contamination and extend shelf life (35).

Fruit processing byproducts contain carbohydrates such as cellulose and starch, which can replace traditional starches in films and coatings (36).

Mango

Mangoes are rich in pectin, a natural gelling agent (37). Plasticizers like glycerol, sorbitol and polyethylene glycol enhance flexibility and permeability by disrupting polymer hydrogen bonds (38). Sorbitol improves film strength and elasticity (39). Nayak and Rayaguru's method for extracting starch from mango seed kernels involved washing, slicing, peeling and sun-drying the kernels, followed by oven drying at 105 °C. The dried kernels were ground and soaked in sulfur dioxide for four days at varying mass-to-liquid ratios (1:5, 1:10, 1:15, 1:20), with starch yield depending on the ratio and soaking time (40). Mango Kernel Flour (MKF)/glycerol films were prepared by mixing distilled water with 5 % w/w MKF, heating to 90 °C and stirring for 30 min. Glycerol (50 % w/w of MKF) was added, followed by further stirring. The mixture was sonicated to remove bubbles and poured into Petri plates to dry for 48 hrs (41).

Banana

Banana pseudo-stems are rich in cellulose fibers (42), yet remain underutilized. One innovative application involves the use of the ionic liquid 1-allyl-3-methylimidazolium chloride ([AMIm][Cl]) along with delignified fibers to create regenerated cellulose films for sustainable mango packaging (43). Additionally, banana peel fibers, often from food industry waste, offer a sustainable alternative to polyethylene or polystyrene (44). Banana peels are soaked in 0.2 M sodium metabisulfite solution for 45 min to act as a preservative and

antioxidant (45). After boiling for 30 min, the peels are drained, dried and blended into a paste. To prepare filmogenic suspensions, banana peel flour, cornstarch and water were heated to 90 °C, stirred and homogenized. Glycerol was added and the mixture was stirred further before filtering and pouring into Petri dishes to form starch banana peel film (46).

Citrus

Orange peels, rich in starch and pectin, hold potential for bioplastic production due to their strength-enhancing properties (47). Limonene in the peels can be oxidized with CO₂ to create bioplastics, but industrial use remains limited (48). Sweet orange peel was washed, sun-dried for 20 hrs, oven-dried at 60 °C for 18 hrs and ground into fine powder (100 µm) before storage at room temperature (49). PVA dissolved in hot water was heated to 90 °C and stirred for 2 hrs. OPP (5-20 %) was added, stirred further and the mixture was poured onto a glass plate to set before removing the film. (50-52).

Jackfruit

Starch, a cheap and renewable resource, is widely used for biodegradable films (48). Using jackfruit seed starch for bioplastics can help reduce waste from the jackfruit seed (53). Jackfruit seeds were cleaned, cubed and blended with water. The slurry was filtered and the starch was separated after settling for 24 hrs. The wet starch was oven-dried at 65 °C for 12 hrs, ground and sifted into a fine powder (54). Jackfruit starch solution was stirred for 24 hrs, gelatinized at 95 °C and mixed with glycerol and anthocyanin. The solution was poured onto a Mylar plate, spread evenly and dried at 30 °C for 24 hrs (55).

Apple

Red apple pomace, comprising 25-30 % of raw apples, is rich in phenolic compounds. PVA films with apple pomace exhibit antioxidant properties, but whole pomace reduces tensile strength and elongation (56). Apple peels were washed, sun-dried, ground and sieved (60-mesh) to obtain a fine powder, which was then stored in an airtight container. Apple peel-based edible films were made using high-pressure homogenization. Increasing glycerol concentration (23-44 %) reduced moisture content and improved barrier properties, making them suitable for food wrapping.

Animal feed in agriculture

Mango

Mango peels, rich in sugars (13.2 %), are energy-rich feed but unsuitable for ruminants due to high moisture and acidity. Adding nitrogen or protein improves energy utilization and a mixture of rice straw and beans promotes fermentation for palatable silage (57). Incorporating *Leucaena* leaves into mango peel silage with rice straw improved digestibility to 60 % (58). Substituting maize bran with up to 37.5 % processed mango fruit boosted sheep growth without affecting welfare (59).

Banana

Goats, dairy cattle, beef cattle and rabbits readily consume banana waste due to its digestibility (60–63). Calves fed banana waste showed improved growth (62), Dairy cows fed banana peels increased milk production. Dried, ripe plantain

peels replace corn for goats, while dried banana peels can be fed to pigs up to 20 % without affecting growth (64).

Pineapple

Pineapple juice extraction produces waste (peels, pomace, crowns), accounting for 33 % of processing by-products (64). Raw pineapple waste is high in dry matter and fiber but low in minerals, containing soluble sugars, crude protein and pectin (65). Silage with pineapple waste and chicken litter reduces feed costs and pineapple waste can replace up to 50 % of roughage in dairy calf feed without affecting milk production (66). Anaerobic digestion slurry can be used as animal feed (67).

Citrus

Pigs can consume up to 5 % dried citrus pulp and fresh pulp should be mixed with hay or straw for better ensiling. Cattle enjoy ensiled orange pulp, which improves silage quality (64). Citrus pulp, mixed with straw, creates high-quality silage and helps buffer rumen pH due to its high fiber. It is a safer feed alternative to grain for dairy cows on high-concentrate diets (68).

Apple

Apple pomace is a valuable feed for calves and provides 1.86 Mcal of metabolizable energy per kg of dry matter for lactating dairy cows (69). Holstein cows fed 15 % ensiled apple pomace showed improved milk yield and composition (70).

Biofuel production: Fruit wastes, rich in sugars and carbohydrates, can be used for bioethanol production through enzymatic hydrolysis with xylanase and cellulase enzymes. Pineapples and bananas are effective sources and cleaning fruit wastes with water and steam, followed by fermentation with *Saccharomyces cerevisiae*, produces bioethanol (71). Fermenting apple pomace and overripe bananas with baker's yeast can yield alcohol concentrations of 48 % after distillation (72).

Bioethanol can be produced from pineapple and jackfruit waste through acid hydrolysis and fermentation with *Saccharomyces cerevisiae*, yielding 0.090 % ethanol from pineapple and 0.045 % from jackfruit (73). Orange and mango peels can also be fermented with bacterial strains like *Enterobacter cloacae*, *Pseudomonas aeruginosa* and *Bacillus cereus* to produce ethanol (74).

Banana peels, with high cellulose and low lignin, are ideal for bioethanol production. Ethanol yield depends on pH, typically around 4.5 to 5.5 for maximum enzyme activity during hydrolysis (75). Fruit wastes like bananas, apples and pomegranates can also serve as substrates, reducing bioethanol production costs. Converting daily household waste into ethanol offers a promising solution.

Conclusion

Fruit waste holds immense potential for contributing to sustainable practices across multiple sectors. Rich in bioactive compounds, fibers and nutrients, these by-products can be efficiently transformed into high value outputs such as biofertilizers, biodegradable packaging, biofuels, these by-products can be efficiently transformed into high value outputs

and animal feed. This review emphasizes the need for innovative processing techniques and collaborative efforts to maximize the value of fruit by-products. Adopting these strategies supports a circular economy, minimizes environmental impact and promotes sustainable development.

Authors' contributions

DP and PSK prepared the original draft. KSVS, IM, VWAG and RG corrected and edited the manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

Declaration of generative AI and AI-assisted technologies in the writing process: Used Grammarly AI software to improve grammar and language.

References

- Bancal V, Ray RC. Overview of food loss and waste in fruits and vegetables: From issue to resources. In: Fruits and vegetable wastes: Valorization to bioproducts and platform chemicals. Springer. 2022;3–29. https://doi.org/10.1007/978-981-16-9527-8_1
- Ali A, Riaz S, Sameen A, Naumovski N, Iqbal MW, Rehman A, et al. The disposition of bioactive compounds from fruit waste, their extraction and analysis using novel technologies: A review. Processes. 2022;10(10):2014. <https://doi.org/10.3390/pr10102014>
- Ali SY, Hossain MM, Zakaria M, Hoque MA, Ahiduzzaman M. Postharvest losses of mangoes at different stages from harvesting to consumption. Int J Bus Soc Sci Res. 2019;7(4): 21–6.
- Mebratie MA, Haji J, Woldetsadik K, Ayalew A, Ambo E. Determinants of postharvest banana loss in the marketing chain of central Ethiopia. Food Sci Qual Manag. 2015;37:52–63.
- Ningombam S, Noel A, Singh J. Post-harvest losses of pineapple at various stages of handling from the farm level up to the consumer in Manipur. Int J Agric Sci. 2019;11:9235–7.
- Rajabi S, Lashgarara F, Omid M, Hosseini SJF. Quantifying the grapes losses and waste in various stages of supply chain. In: Biological Forum. Research Trend. 2015:225.
- Thapa S, Sapkota S, Adhikari D. Effect of different postharvest treatments on prolonging shelf life and maintaining quality of sweet orange (*Citrus sinensis* Osbeck.). Sustainability in Food and Agriculture. 2020;1(2):69–75. https://doi.org/10.1007/978-981-16-9527-8_1
- Lau KQ, Sabran MR, Shafie SR. Utilization of vegetable and fruit by-products as functional ingredient and food. Front Nutr. 2021;8:661693. <https://doi.org/10.3389/fnut.2021.661693>
- Soni D, Saxena G. Hidden potential of fruit waste and its utilization. Sustainability, Agri, Food and Environmental Research. 2021;9(3):469–79. <https://doi.org/10.7770/safer-V0N0-art2213>
- Cheok CY, Mohd Adzahan N, Abdul Rahman R, Zainal Abedin NH, Hussain N, Sulaiman R, et al. Current trends of tropical fruit waste utilization. Crit Rev Food Sci Nutr. 2016;58(3):1–27. <https://doi.org/10.1080/10408398.2016.1176009>
- Sagar NA, Pareek S, Sharma S, Yahia EM, Lobo MG. Fruit and vegetable waste: Bioactive compounds, their extraction and possible utilization. Compr Rev Food Sci Food Saf. 2018;17(3):512–31. <https://doi.org/10.1111/1541-4337.12330>
- Mwaurah PW, Kumar S, Kumar N, Panghal A, Attkan AK, Singh VK, et al. Physicochemical characteristics, bioactive compounds and industrial applications of mango kernel and its products: A review. Compr Rev Food Sci Food Saf. 2020;19(5):2421–46. <https://doi.org/10.1111/1541-4337.12598>
- Sharma L, Saini CS, Sharma HK, Sandhu KS. Biocomposite edible coatings based on cross linked-sesame protein and mango puree for the shelf life stability of fresh-cut mango fruit. J Food Process Eng. 2019;42(1):e12938. <https://doi.org/10.1111/jfpe.12938>
- Serna-Cock L, García-Gonzales E, Torres-León C. Agro-industrial potential of the mango peel based on its nutritional and functional properties. Food Reviews International. 2016;32(4):364–76. <https://doi.org/10.1080/87559129.2015.1094815>
- Bonneau A, Boulanger R, Lebrun M, Maraval I, Valette J, Guichard É, et al. Impact of fruit texture on the release and perception of aroma compounds during in vivo consumption using fresh and processed mango fruits. Food Chem. 2018;239:806–15. <https://doi.org/10.1016/j.foodchem.2017.07.017>
- Wadhwa M, Bakshi MPS, Makkar HPS. Wastes to worth: value added products from fruit and vegetable wastes. CABI Reviews. 2016;(2015):1–25. <https://doi.org/10.1079/PAVSNNR201510043>
- Ravani A, Joshi D. Mango and its by product utilization-a review. Energy (kcal). 2013;74(44):2013.
- Sharma R, Oberoi HS, Dhillon GS. Fruit and vegetable processing waste. In: Agro-industrial wastes as feedstock for enzyme production. Elsevier. 2016:23–59. <https://doi.org/10.1016/B978-0-12-802392-1.00002-2>
- Matani AG, Bhatkar PP, Doifode SK. Currency note paper and banana fiber pulp waste utilization and its effect of density, moisture and temperature on wood products. Group (New York).9:0–33.
- Gomes AAB, Ferreira ME, Pimentel TC. Bread with flour obtained from green banana with its peel as partial substitute for wheat flour: Physical, chemical and microbiological characteristics and acceptance. Int Food Res J. 2016;23(5).
- Mohapatra D, Mishra S, Sutar N. Banana and its by-product utilisation: an overview. 2010.
- Li T, Shen P, Liu W, Liu C, Liang R, Yan N, et al. Major polyphenolics in pineapple peels and their antioxidant interactions. Int J Food Prop. 2014;17(8):1805–17. <https://doi.org/10.1080/10942912.2012.732168>
- Sarangi PK, Singh AK, Srivastava RK, Gupta VK. Recent progress and future perspectives for zero agriculture waste technologies: Pineapple waste as a case study. Sustainability. 2023;15(4):3575. <https://doi.org/10.3390/su15043575>
- Polania AM, Londoño L, Ramírez C, Bolívar G, Aguilar CN. Valorization of pineapple waste as novel source of nutraceuticals and biofunctional compounds. Biomass Convers Biorefin. 2023;13(5):3593–618. <https://doi.org/10.1007/s13399-022-02811-8>
- Maqbool Z, Khalid W, Atiq HT, Koraqi H, Javaid Z, Alhag SK, et al. Citrus waste as source of bioactive compounds: extraction and utilization in health and food industry. Molecules. 2023;28(4):1636. <https://doi.org/10.3390/molecules28041636>
- Benny N, Shams R, Dash KK, Pandey VK, Bashir O. Recent trends in utilization of citrus fruits in production of eco-enzyme. J Agric Food Res. 2023;13:100657. <https://doi.org/10.1016/j.jafr.2023.100657>
- Ramadhani AN, Malik AF, Fitriana WR. Utilization of wasted cooking oil and essential oil of sweet orange peel (*Citrus sinensis*) as aromatherapy candles. Equilibrium Journal of Chemical Engineering. 2023;7(2):191. <https://doi.org/10.20961/equilibrium.v7i2.80308>
- Kaur S, Panesar PS, Chopra HK. Citrus processing by-products: an overlooked repository of bioactive compounds. Crit Rev Food Sci Nutr. 2023;63(1):67–86.

29. Pragalyaashree MM, Monicka A, Blessie F, Deivanayagame T. Jackfruit processing and utilization of its waste: A review. *Asia-Pacific J Sci Technol*. 2023;28(4).
30. Sarangi PK, Srivastava RK, Singh AK, Sahoo UK, Prus P, Dziekański P, et al. The utilization of Jackfruit (*Artocarpus heterophyllus* L.) waste towards sustainable energy and biochemicals: The attainment of zero-waste technologies. *Sustainability*. 2023;15(16):12520. <https://doi.org/10.3390/su151612520>
31. Wang ChaoJin WC, Lai QingHua LQ. *Advances in comprehensive utilization of fruit peel in China*. 2016.
32. Duan Y, Mehariya S, Kumar A, Singh E, Yang J, Kumar S, et al. Apple orchard waste recycling and valorization of valuable product-A review. *Bioengineered*. 2021;12(1):476–95. <https://doi.org/10.1080/21655979.2021.1872905>
33. Drzal LT. *Natural fibers, biopolymers and biocomposites*. CRC Press;2005.
34. Birania S, Kumar S, Kumar N, Attkan AK, Panghal A, Rohilla P, et al. Advances in development of biodegradable food packaging material from agricultural and agro-industry waste. *Journal of Food Process Engineering*. 2022;45.
35. Gupta P, Toksha B, Rahaman M. A review on biodegradable packaging films from vegetative and food waste. *The Chemical Record*. 2022;22(7):e202100326. <https://doi.org/10.1002/tcr.202100326>
36. Yadav A, Kumar N, Upadhyay A, Pratibha, Anurag RK. Edible packaging from fruit processing waste: A comprehensive review. *Food Reviews International*. 2023;39(4):2075–106. <https://doi.org/10.1080/87559129.2021.1940198>
37. Nguyen HDH, Nguyen HVH, Savage GP. Properties of pectin extracted from vietnamese mango peels. *Foods*. 2019;8(12):629. <https://doi.org/10.3390/foods8120629>
38. Bourtoom T, Chinnan MS. Preparation and properties of rice starch–chitosan blend biodegradable film. *LWT – Food Science and Technology*. 2008;41(9):1633–41. <https://doi.org/10.1016/j.lwt.2007.10.014>
39. Susmitha A, Sasikumar K, Rajan D, Padmakumar MA, Nampoothiri KM. Development and characterization of corn starch-gelatin based edible films incorporated with mango and pineapple for active packaging. *Food Biosci*. 2021;41:100977. <https://doi.org/10.1016/j.fbio.2021.100977>
40. Admase AT, Sendekie ZB, Alene AN. Biodegradable film from mango seed kernel starch using pottery clay as filler. *J Polym Environ*. 2022;30(8):3431–46. <https://doi.org/10.1007/s10924-022-02449-7>
41. Gomez-Caturla J, Ivorra-Martinez J, Quiles-Carrillo L, Balart R, Garci-Garcia D, Dominici F, et al. Improvement of the barrier and mechanical properties of environmentally friendly mango kernel flour/glycerol films by varying the particle size of mango kernel flour. *Ind Crops Prod*. 2022;188:115668. <https://doi.org/10.1016/j.indcrop.2022.115668>
42. Guimarães JL, Frollini E, da Silva CG, Wypych F, Satyanarayana KG. Characterization of banana, sugarcane bagasse and sponge gourd fibers of Brazil. *Ind Crops Prod*. 2009;30(3):407–15. <https://doi.org/10.1016/j.indcrop.2009.07.013>
43. Ai B, Zheng L, Li W, Zheng X, Yang Y, Xiao D, et al. Biodegradable cellulose film prepared from banana pseudo-stem using an ionic liquid for mango preservation. *Front Plant Sci*. 2021;12:625878. <https://doi.org/10.3389/fpls.2021.625878>
44. Yusof NN, Yusoff M. Investigation of chemical analysis and physical properties of bio-polymer waste banana peel fibre composite. *IOP Conf Ser Earth Environ Sci*. 2020;596(1):012042. <https://doi.org/10.1088/1755-1315/596/1/012042>
45. Arjun J, Manju R, Rajeswaran SR, Chandru M. Banana peel starch to biodegradable alternative products for commercial plastics. *GSC Biol Pharm Sci*. 2023;22(2):234–44. <https://doi.org/10.30574/gscbps.2023.22.2.0066>
46. Medeiros Silva VD, Coutinho Macedo MC, Rodrigues CG, Neris dos Santos A, de Freitas e Loyola AC, Fante CA. Biodegradable edible films of ripe banana peel and starch enriched with extract of *Eriobotrya japonica* leaves. *Food Biosci*. 2020;38:100750. <https://doi.org/10.1016/j.fbio.2020.100750>
47. Yaradoddi JS, Banapurmath NR, Ganachari SV, Soudagar MEM, Sajjan AM, Kamat S, et al. Bio-based material from fruit waste of orange peel for industrial applications. *Journal of Materials Research and Technology*. 2022;17:3186–97. <https://doi.org/10.1016/j.jmrt.2021.09.016>
48. Abdul Rahman W, Syafiqah Ismail A, Aishatun Majid N. Preparation and characterization of biocomposite film derived from microcrystalline cellulose (MCC) of jackfruit rind waste. *Mater Today Proc*. 2022;66:4055–60. <https://doi.org/10.1016/j.matpr.2022.06.071>
49. Sambudi NS, Lin WY, Harun NY, Mutiari D. Modification of poly (lactic acid) with orange peel powder as biodegradable composite. *Polymers (Basel)*. 2022;14(19):4126. <https://doi.org/10.3390/polym14194126>
50. Ahmed I, Zia MA, Hussain MA, Akram Z, Naveed MT, Nowrouzi A. Bioprocessing of citrus waste peel for induced pectinase production by *Aspergillus niger*; its purification and characterization. *J Radiat Res Appl Sci*. 2016;9(2):148–54. <https://doi.org/10.1016/j.jrras.2015.11.003>
51. Balaji A, Karthikeyan B, Swaminathan J, Sundar Raj C. Thermal behavior of cardanol resin reinforced 20mm long untreated bagasse fiber composites. *Int J Polymer Anal Characterization*. 2018;23(1):70–7. <https://doi.org/10.1016/j.jrras.2015.11.003>
52. Perumal AB, Sellamuthu PS, Nambiar RB, Sadiku ER. Development of polyvinyl alcohol/chitosan bio-nanocomposite films reinforced with cellulose nanocrystals isolated from rice straw. *Appl Surf Sci*. 2018;449:591–602. <https://doi.org/10.1016/j.apsusc.2018.01.022>
53. Vargas-Rechia CG, Reicher F, Sierakowski MR, Heyraud A, Driguez H, Szymanska-Chargot M, et al. A profile of the South African table grape market value chain. *Bioresour Technol*. 2015;6(1):1–14.
54. Wahidin M, Srimarlita A, Sulaiman I, Indarti E. Transparency and thickness of jackfruit and durian seed starch edible film. *IOP Conf Ser Earth Environ Sci*. 2021;667(1):012030. <https://doi.org/10.1088/1755-1315/667/1/012030>
55. Costa LA da, Diógenes ICN, Oliveira M de A, Ribeiro SF, Furtado RF, Bastos M do SR, et al. Smart film of jackfruit seed starch as a potential indicator of fish freshness. *Food Science and Technology*. 2021;41(2):489–96. <https://doi.org/10.1590/fst.06420>
56. Lan W, Wang S, Zhang Z, Liang X, Liu X, Zhang J. Development of red apple pomace extract/chitosan-based films reinforced by TiO₂ nanoparticles as a multifunctional packaging material. *Int J Biol Macromol*. 2021;168:105–15. <https://doi.org/10.1016/j.ijbiomac.2020.12.051>
57. Lalramhlimi B, Mukherjee D, Chakraborty I, Ghosh N, Chattopadhyay A, Dey RC. Fruit and vegetable wastes as livestock feeds. In: *Fruits and vegetable wastes: Valorization to bioproducts and platform chemicals*. Springer;2022:139–68. https://doi.org/10.1007/978-981-16-9527-8_6
58. Sruamsiri S, Silman P. Nutritive value and nutrient digestibility of ensiled mango by-products. 2009.
59. Ibrahim TA, Jamilu M, Agboola LO, Abdu SB, Hasaan MR, Kailani AI. Growth performance of Yankasa sheep fed increasing levels of processed mango (*Mangifera indica*) fruits replacing maize bran. *Livestock Research for Rural Development*. 2020;32(9):144.
60. Álvarez S, Méndez P, Martínez-Fernández A. Fermentative and nutritive quality of banana by-product silage for goats. *J Appl Anim Res*. 2015;43(4):396–401.

61. Katongole CB, Bareeba FB, Sabiiti EN, Ledin I. Nutritional characterization of some tropical urban market crop wastes. *Anim Feed Sci Technol.* 2008;142(3-4):275–91. <https://doi.org/10.1016/j.anifeedsci.2007.09.002>
62. Xue Z, Mu L, Cai M, Zhang Y, Wanapat M, Huang B. Effect of using banana by-products and other agricultural residues for beef cattle in southern China. *Trop Anim Health Prod.* 2020;52:489–96. <https://doi.org/10.1007/s11250-019-02031-9>
63. Ekwe OO, Osakwe II, Nweze BO. The effect of replacing maize with cassava “sievate” using banana leaves as basal forage in the diet of weaned rabbit. *Ozean J Appl Sci.* 2011;4(1):52–8.
64. Wadhwa M, Saini JK, Bakshi MPS. Potential of bottle gourd pomace as feed for goat bucks. *Small Rumin Res.* 2013.
65. Pereira ES, Regadas Filho JGL, Freitas ER, Neiva JNM, Candido MJD. Energetic value from by-product of the Brazil agroindustria; 2009.
66. Sruamsiri S. Agricultural wastes as dairy feed in Chiang Mai. *Animal Sci J.* 2007;78(4):335–41. <https://doi.org/10.1111/j.1740-0929.2007.00445.x>
67. Rani DS, Nand K. Ensilage of pineapple processing waste for methane generation. *Waste Management.* 2004;24(5):523–8. <https://doi.org/10.1016/j.wasman.2003.10.010>
68. Crawshaw R. Co-product feeds: animal feeds from the food and drinks industries R Crawshaw Nottingham University Press, Nottingham, 2001. *J Sci Food Agric.* 2003;83(4):362–2. <https://doi.org/10.1002/jsfa.1326>
69. Council NR, Nutrition C on A, Nutrition S on DC. Nutrient requirements of dairy cattle: 2001. National Academies Press; 2001.
70. Ghoreishi SF, Pirmohammadi R, Yansari AT. Effects of ensiled apple pomace on milk yield, milk composition and DM intake of Holstein dairy cows; 2007.
71. Jahid M, Gupta A, Sharma DK. Production of bioethanol from fruit wastes (banana, papaya, pineapple and mango peels) under milder conditions. *J Bioprocess Biotech.* 2018;8(3):1–11. <https://doi.org/10.4172/2155-9821.1000327>
72. Borah D, Mishra V. Production of biofuel from fruit waste. *Int J Adv Biotechnol Res.* 2011;1(1):71–4.
73. Gosavi P, Chaudhary Y, Durve-Gupta A. Production of biofuel from fruits and vegetable wastes. *Yeast.* 2004;17.
74. Vinotha T, Umamaheswari N, Pandiyan J, Al-Ghanim KA, Nicoletti M, Govindarajan M. Biofuel Production from mango and orange peel and tapioca shells by fermentation using consortium of bacteria: Agricultural and food waste valorization. *Fermentation.* 2023;9(7):678. <https://doi.org/10.3390/fermentation9070678>
75. Hamzah MA, Alias AB, Ahmad NE. Production of biofuel (bio-ethanol) from fruitwaste: Banana peels. *Int J Eng Adv Technol.* 2019;9(1):5897–901. <https://doi.org/10.35940/ijeat.A3024.109119>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc
See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.