



RESEARCH ARTICLE

Nutrient-rich or anti-nutritional challenge? Evaluating *Senna spectabilis* (DC.) H.S. Irwin & Barneby for sustainable animal feed production

Bargavi S^{1*}, Baranidharan K^{1*}, Tilak M², Revathi R³, Ganesan K N⁴, Ragunath K P⁵, Vijayabhama M⁶, Hemalatha P¹, Ravi R¹, Kabinesh V¹ & Suwethaasri D¹

¹Department of Forest Products and Wildlife, Forest College and Research Institute, TNAU, Mettupalayam 641 301, Tamil Nadu, India

²Department of Agroforestry, Forest College and Research Institute, TNAU, Mettupalayam 641 301, Tamil Nadu, India

³Department of Forest Biology and Tree Improvement, Forest College and Research Institute, Mettupalayam, Tamil Nadu 641 301, India

⁴Department. of Rice, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

⁵Centre for Water and Geospatial studies, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

⁶Department of Physical Sciences and Information Technology, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

*Email: baranidharan.k@tnau.ac.in, bargavi.phdfor2022@tnau.ac.in



ARTICLE HISTORY

Received: 27 February 2025

Accepted: 28 March 2025

Available online

Version 1.0 : 20 April 2025

Version 2.0 : 23 April 2025



Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonepublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Bargavi S, Baranidharan K, Tilak M, Revathi R, Ganesan KN, Ragunath KP, Vijayabhama M, Hemalatha P, Ravi R, Kabinesh V, Suwethaasri D. Nutrient-rich or anti-nutritional challenge? Evaluating *Senna spectabilis* (DC.) H.S. Irwin & Barneby for sustainable animal feed production. Plant Science Today. 2025; 12(2): 1-8. <https://doi.org/10.14719/pst.7953>

Abstract

This study investigated the nutritional composition and anti-nutritional factors of *Senna spectabilis* as a potential fodder for animal feed. Key parameters such as crude protein, crude fiber, dry matter, digestibility and anti-nutritional factors like phenols, tannins, saponins and nitrates were analyzed from the five formulations named from SS₁ to SS₅. The results showed significant variations across the species in terms of digestibility and energy content, with formulations SS₁ and SS₄ demonstrating superior digestibility and higher Total Digestible Nutrients (TDN). On the other hand, formulation SS₅ exhibited higher crude fiber content and lower digestibility, making it less efficient as a feed source. Anti-nutritional factors such as saponins and nitrates were found to be highest in formulation SS₁, which could affect feed intake and livestock health if present in excessive amounts. The study also noted moderate levels of tannins and phenols in several species, which could impact nutrient absorption for animals. Additionally, concentrate feed mixtures were considered in comparison to fodder-based feeds for their influence on livestock growth and performance. These findings suggest that careful selection of feed ingredients, considering both nutritional and anti-nutritional parameters, is essential for optimizing livestock nutrition. Formulations with lower levels of anti-nutritional compounds like SS₂, offer more sustainable feed options, while further research is required to understand the effects of these compounds on livestock health and performance.

Keywords

animal feed pellets; anti-nutritional factors; nutritional composition; *Senna spectabilis*

Introduction

Invasive alien species cause significant damage to native ecosystems. These plant species are introduced by humans, either intentionally or accidentally, into regions outside their natural habitats. Without natural predators or competitors, these species can spread quickly, severely impacting native biodiversity, even in protected areas (1). Invasive plants often share traits that help them dominate; they tend to grow rapidly, have short life cycles, high

reproductive capacity, strong competitive abilities and sometimes even release chemicals that harm other plants (allelopathy) (2). These adaptations enable them to outcompete native species, disturbing entire ecosystems. The effects of plant invasions are widespread and well-documented, making them one of the most serious threats to the integrity of natural and semi-natural ecosystems globally (3-7). Invasive Alien Plants (IAPs) often compete with native species for essential resources such as light, water and nutrients, leading to declines in the abundance and diversity of native plants (8). These invasions disrupt ecosystem balance, altering plant and animal community structures. Such changes can affect food webs, nutrient cycling and overall ecosystem health. Furthermore, IAPs can modify the physical and chemical environment, creating conditions that disadvantage native species even further (9).

Senna spectabilis is a fast-growing, medium-sized tree native to Central and South America. Its name is derived from its historical use as a laxative, with Arabic roots referring to its purgative effects. Part of the *Senna* genus, which includes over 200 species (10), it was first noticed in India in the Sathyamangalam Forest and Wayanad Wildlife Sanctuary of Kerala (11). This tree can grow over 10 meters tall, with young branches often covered in fine hair. Its leaves are large, with multiple pairs of leaflets averaging 20 per leaf. Each leaflet is lance-shaped, measuring 3-8 cm in length and 2 cm in width, with a hairy underside and a smoother, shiny top. Its yellow flowers appear in clusters at the ends of branches, with five unevenly sized petals, the largest reaching up to 2.5 cm. The plant also has three types of stamens, some of which are rudimentary. Its fruit is a long, flat pod that opens along one side (12) (Fig. 1).

Originally introduced to Indian gardens for its ornamental value, *S. spectabilis* has unfortunately spread beyond cultivation and invaded forested areas in regions like Mysore and Sikkim. This aggressive invader thrives in Wayanad and Tamil Nadu, producing numerous seedlings that threaten native species. As a result, it has been classified as an 'Invasive Alien Species', causing significant harm to biodiversity. What was once an ornamental tree has now become a serious ecological threat, overtaking large areas of natural forest and displacing native species (13). Even

herbivores, which might typically help control its spread, inadvertently aid its expansion by consuming and dispersing its seeds (14). The real challenge, however, lies in the tree's incredible resilience. *S. spectabilis* is extremely difficult to remove, as its seeds remain viable for long periods and even damaged branches, cut stems, or uprooted trees can regenerate new growth. This persistence makes it a major obstacle in efforts to protect native biodiversity (15, 16).

Despite being considered a dangerous invader, *S. spectabilis* may offer an unexpected benefit to livestock farmers. The tree can form a symbiotic relationship with nitrogen-fixing bacteria called rhizobia. These bacteria help convert atmospheric nitrogen into usable forms, enriching the plant's leaves with nutrients like protein and fiber. As a result, these nutrient-dense leaves could serve as a supplementary feed source for livestock, including goats, sheep and cattle, particularly in times of scarcity, such as during droughts or periods of overgrazing (17, 18).

Additionally, the nutrient composition of *S. spectabilis* leaves suggests potential use as a component in livestock feed concentrates. Concentrates are feed blends that supply essential nutrients such as proteins, sugars and fats at elevated levels while containing less than 18 % crude fiber and over 60 % total digestible nutrients (TDN). These are categorized into energy-rich and protein-rich concentrates, depending on their crude protein content. Energy-rich concentrates contain less than 18 % crude protein, while protein-rich concentrates contain over 18 % (19). Given its nitrogen-fixing ability and high protein content, *S. spectabilis* leaves could be explored as an alternative protein-rich concentrate, reducing reliance on conventional feed sources. However, further research is necessary to assess its digestibility, anti-nutritional factors and long-term effects on livestock health. Based on these findings, *S. spectabilis* leaves have been selected for this study as a potential component in animal feed pellets. Their nitrogen-fixing ability, high protein content and resilience make them a promising alternative protein-rich feed source. Further analysis will focus on their nutritional composition, digestibility and anti-nutritional factors to determine their suitability for livestock feed formulations.



a. *S. spectabilis* coppiced shoots



b. *S. spectabilis* tree

Fig. 1. *S. spectabilis* (DC.) H.S. Irwin & Barneby.

Materials and Methods

Its ability to thrive in diverse ecosystems makes it a significant species in both natural and cultivated landscapes. To explore its viability in livestock nutrition, different feed pellet formulations were developed using *S. spectabilis* leaf powder as the primary ingredient. The formulations were supplemented with various oilcakes, rice bran, maize grain, mineral mixtures, molasses and salt to enhance nutritional value and palatability.

The tender leaves of *S. spectabilis* species were collected, shade dried, pulverized and formed into pellets using a pelleting machine in five formulations (SS₁ - SS₅), to explore their potential as a fodder resource. The table below presents the different formulations used in the study (Table 1.) (Fig. 2 and 3).

The nutritional profiling involved in assessing Dry Matter (DM) by oven-drying until a constant weight was achieved, while ash content was measured by incinerating the samples at 550 °C (20). Crude Protein (CP) was analyzed using the Micro-Kjeldahl method and Crude Fat (CF) was determined through Soxhlet extraction with ethanol (20). Crude Fiber was quantified through acid and alkali treatments, whereas Acid Detergent Fiber (ADF) and Neutral Detergent Fiber (NDF) were assessed using detergent solutions (20). The Nitrogen-Free Extract (NFE) was calculated as $100 - [CP + CF + \text{crude fiber} + \text{ash}]$ (21). Digestible Dry Matter (DDM) was estimated using the formula $DDM = 88.9 - (ADF \times 0.779)$ (22) and Relative Feed Value (RFV) was derived from $(DMI \times DDM) \div 1.29$. Anti-nutritional factors were evaluated using standard procedures. Total phenolic content and Folin tannin-reactive substances were

Table 1. Formulations of *S. spectabilis*-based animal feed pellets

Formulation No.	Formulations
SS ₁	<i>S. spectabilis</i> leaf powder (Control) (100 %)
SS ₂	<i>S. spectabilis</i> leaf powder (50 %) + Groundnut oilcake (15 %) + Rice bran (10 %) + Maize grain (20 %) + Mineral mixture (2 %) + Molasses (2 %) + Salt (1 %)
SS ₃	<i>S. spectabilis</i> leaf powder (50 %) + Gingelly oilcake (15 %) + Rice bran (10 %) + Maize grain (20 %) + Mineral mixture (2 %) + Molasses (2 %) + Salt (1 %)
SS ₄	<i>S. spectabilis</i> leaf powder (50 %) + Coconut oilcake (15 %) + Rice bran (10 %) + Maize grain (20 %) + Mineral mixture (2 %) + Molasses (2 %) + Salt (1 %)
SS ₅	<i>S. spectabilis</i> leaf powder (50 %) + Sunflower oilcake (15 %) + Rice bran (10 %) + Maize grain (20 %) + Mineral mixture (2 %) + Molasses (2 %) + Salt (1 %)



Fig. 2. Formulation of *S. spectabilis*-based animal feed pellets.



Fig. 3. *S. spectabilis* based animal feed pellets.

determined through the Folin-Ciocalteu assay, followed by spectrophotometric measurements at 760 nm and 700 nm, respectively (23).

Total Nitrates were measured following a standard protocol (24), with potassium nitrate as the standard. Saponins were determined using a standard methodology (25) involving vanillin and sulfuric acid. These findings were employed to optimize nutritional value while minimizing anti-nutritional factors.

Statistical analysis

The design of the experiment was a Completely Randomized Design (CRD). Data focused initially on the Invasive Alien Species, *Senna spectabilis* with a total of five formulations. All statistical analyses were performed using the AGRES and TNAUSTAT software.

Results and Discussion

Table 2 presents the proximate composition of different formulations, highlighting variations in Dry Matter (DM), Moisture Content (MC), Ash Content (AC), Crude Protein (CP), Crude Fat (CFa) and Crude Fiber (CFi). These parameters play a crucial role in assessing the nutritional quality of the species, which can influence their suitability for animal feed. The observed differences in composition indicate species-specific variations in nutrient availability and forage quality.

Table 2. Proximate composition of *S. spectabilis* based animal feed pellets

S. No	Formulations	DM (%)	MC (%)	AC (%)	CP (%)	CFa (%)	CFi (%)
1	SS ₁	88.91*	11.09	11.55*	16.18	2.27	4.50*
2	SS ₂	86.70	13.30*	11.07	15.75	2.43*	5.50*
3	SS ₃	84.01	15.99*	10.17	16.62*	2.09	3.0
4	SS ₄	88.95*	11.05	9.89	16.31	2.03	3.0
5	SS ₅	86.44	13.56*	13.25*	16.18	2.67*	5.50*
	Mean	87.00	12.99	11.18	16.20	2.29	4.30
	SEd	0.831	0.189	0.134	0.194	0.033	0.077
	CD (0.05)	1.787	0.407	0.288	0.417	0.070	0.166

DM - Dry Matter, MC - Moisture Content, AC - Ash Content, CP - Crude Protein, CFa - Crude Fat, CFi - Crude Fibre

The proximate composition analysis of five different formulations (SS₁-SS₅) revealed significant variations in nutritional parameters, influencing their suitability as animal feed. The Dry Matter (DM) content ranged from 84.01 % in SS₃ to 88.95 % in SS₄, with a mean of 87.00 %. Dry matter content plays a crucial role in determining nutrient concentration and energy availability, affecting feed efficiency and animal performance (26), who discussed how dry matter varies based on species, plant parts and growing conditions. Moisture content (MC) varied between 11.05 % (SS₄) and 15.99 % (SS₃), staying within the acceptable limit of 12 % as per the SNI standard (SNI 8509:2018). Excess moisture content can negatively impact feed storage and stability, the water content in feed is influenced by the material type and environmental conditions (27). Ash Content (AC), representing the mineral composition of feed, was highest in SS₅ (13.25 %) and lowest in SS₄ (9.89 %), with a mean of 11.18 %, aligning with the previous findings (28), which noted that ash content in cattle feed is critical for maintaining mineral balance.

Crude Protein (CP), essential for livestock growth and nutrient absorption, ranged from 15.75 % (SS₂) to 16.62 % (SS₃), meeting the minimum requirement of 16 % recommended for cattle feed. This aligns with previous research (29), which recorded a minimum crude protein content of 10.80 % in cassava-based pellets. Crude fat (CFa), an important energy source in feed, varied between 2.03 % (SS₄) and 2.67 % (SS₅), aligning with the SNI standard (>2.0 %), consistent with earlier reports (30), which found 5.98 % crude fat in mulberry leaf-based pellets. Crude fiber (CFi), which plays a crucial role in digestion and gut health, was highest in

SS₂ and SS₅ (5.50 %) and lowest in SS₃ and SS₄ (3.0 %), with a mean of 4.30 %. These findings were consistent with those of previous research which observed a similar crude fiber content in mulberry leaf-based pellets and a higher crude fiber content in pineapple leaf-based pellets (30, 31). The overall variations in nutritional composition indicate species-specific differences in feed quality, suggesting that selecting appropriate species can significantly enhance livestock nutrition and feed efficiency.

Table 3 summarizes the key nutritional parameters such as Acid Detergent Fiber (ADF), Neutral Detergent Fiber (NDF), Nitrogen-Free Extract (NFE), Digestible Dry Matter (DDM), Dry Matter Intake (DMI), Relative Feed Value (RFV) and Total Digestible Nutrients (TDN) for the various formulations (SS₁-SS₅), providing insights into their digestibility and feed quality. The nutritional analysis (SS₁-SS₅) revealed significant variations across multiple parameters, influencing their suitability as animal feed. The Acid Detergent Fiber (ADF) content ranged from 36.09 % (SS₁) to 40.94 % (SS₅), with a mean of 38.06 %, indicating moderate fiber levels. The Neutral Detergent Fiber (NDF) content varied between 60.74 % (SS₂) and 66.81 % (SS₅), with an average of 63.92 %. This suggests that species with higher NDF content, such as SS₅, may have lower digestibility. The Nitrogen-Free Extract (NFE) content ranged from 62.4 % (SS₅) to 68.77 % (SS₄), with a mean of 66.00 %. This trend of increasing NFE is linked to a decrease in crude fiber content where reduction in crude fiber content resulted in an increase in NFE (32). Similar findings were observed in a study which reported that a pelleted feed mixture with lower fiber content improved the Total Digestible Nutrients (TDN) value (33).

Digestible Dry Matter (DDM) ranged from 57.00 % (SS₅) to 60.78 % (SS₁), with a mean of 59.24 %, reflecting significant differences in the digestibility of the feed. Dry Matter Intake (DMI) values ranged from 1.79 % (SS₅) to 1.97 % (SS₂), with a mean of 1.87 %, suggesting that species like SS₂ may have a higher intake potential. The Relative Feed Value (RFV), which indicates overall feed quality, ranged from 79.09 % (SS₅) to 89.88 % (SS₂), with a mean of 86.07 %. This implies that species with lower fiber content, such as SS₂, generally offer higher

Table 3. Nutritional composition and digestibility of different formulations of *S. spectabilis* based animal feed pellets (SS₁-SS₅)

S. No	Formulations	ADF (%)	NDF (%)	NFE (%)	DDM (%)	DMI (%)	RFV (%)	TDN (%)
1	SS ₁	36.09	64.70	65.5	60.78	1.85	87.16	91.28
2	SS ₂	38.56	60.74	65.25	58.86	1.97*	89.88*	91.96
3	SS ₃	37.96	64.08	68.12	59.32	1.87	85.99	92.44
4	SS ₄	36.78	63.29	68.77*	60.24	1.89	88.25	92.64
5	SS ₅	40.94*	66.81*	62.4	57.00	1.79	79.09	90.08
	Mean	38.06	63.92	66.00	59.24	1.87	86.07	91.68
	SEd	0.406	0.406	1.028	1.087	0.026	1.081	1.446
	CD (0.05)	0.873	0.873	2.211	2.339	0.056	2.325	N/A

feed value. Finally, Total Digestible Nutrients (TDN) values ranged from 90.08 % (SS₅) to 92.64 % (SS₄), with a mean of 91.68 %, indicating high energy content and digestibility across all species. These results highlight the species-specific differences in feed quality, suggesting that selecting for lower fiber content and higher digestibility can significantly enhance livestock nutrition and feed efficiency.

Table 4 presents the concentrations of phenols, tannin, saponin and nitrates in different species (SS₁-SS₅). Significant differences in saponin and nitrate content were observed across the species, indicating variability in anti-nutritional factors that could influence the nutritional quality and suitability of these species for animal feed. The total phenol content ranged from 0.007 % (SS₁) to 0.011 % (SS₅), with a mean of 0.009 %. Phenols in feed provide crucial information regarding rumen fermentation, biohydrogenation, milk yield, cattle weight and potential anti-inflammatory properties (34). This finding aligns with previous research, where a total phenol content of 0.78 % in *Albizia lebbek* + *Prosopis* pod-based feed pellets, highlighting the impact of phenol content

Table 4. Anti-nutritional factors in different formulations of *S. spectabilis* based animal feed pellets (SS₁-SS₅)

S. No	Formulations	Phenols	Tannin	Saponin	Nitrates
1	SS ₁	0.007	0.101	1.376*	0.317*
2	SS ₂	0.009	0.120*	1.069	0.119
3	SS ₃	0.010*	0.098	1.271*	0.281*
4	SS ₄	0.008	0.100	1.011	0.198
5	SS ₅	0.011*	0.112*	1.220*	0.297*
Mean		0.009	0.106	1.189	0.242
SEd		0.000	0.002	0.017	0.004
CD (0.05)		0.000	0.003	0.036	0.009

ADF - Acid Detergent Fibre, NDF - Neutral Detergent Fibre, NFE - Nitrogen Free Extract, DDM - Digestible Dry Matter, DMI - Dry Matter Intake, RFV - Relative Feed Value, TDN - Total Digestible Nutrients

on feed quality and livestock health (35).

Tannin in a potential animal feed can affect nutrient digestibility and animal performance. The study found that the tannin content ranged from 0.1143 % (SS₄) to 0.1290 % (SS₂), with an average of 0.106 %. Excessive tannin consumption reduces feed intake and hinders growth (36). Similar studies reported a tannin content of 0.28 % in legume-based feed pellets (*Lablab purpureus*, *Mucuna pruriens* and *Calopogonium mucunoides*), indicating a consistent trend of tannin levels affecting feed quality (37). The nitrate content ranged from 1.0173 % (SS₃) to 1.2234 % (SS₅), with a mean of 1.189 %. High nitrate levels in feed can pose significant health risks, such as nitrate poisoning, if present in excessive amounts (38). This was reflected in a previous study where the nitrate value of 0.98 % was noted in *Swietenia mahagoni* + *Prosopis* pod-based feed pellets (35).

Saponins are glycosides with soap-like properties. The saponin content ranged from 0.8169 % (SS₄) to 0.9449 % (SS₅), with a mean of 0.9113 %. While saponins reduce feed intake and nutrient absorption, they also provide health benefits such as antimicrobial and anti-inflammatory effects. A similar study found that unpelleted legume feed had a saponin content of 6.17 %, while pelleted legume feed showed lower values (5.5 %), with *Lablab purpureus*, *Calopogonium mucunoides* and

Mucuna pruriens having saponin values of 5.25 %, 5.75 % and 6.5 %, respectively (37). These studies highlight the species-specific variations in anti-nutritional factors and their potential impact on livestock nutrition and health.

Conclusion

The nutritional and anti-nutritional factor analysis of the five different formulations (SS₁-SS₅) reveals significant variations across all measured parameters. This study assessed essential nutritional factors, such as crude protein, crude fiber, dry matter and digestibility, alongside anti-nutritional factors including phenols, tannins, saponins and nitrates. The results highlight the importance of considering both the nutrient composition and anti-nutritional factors when evaluating feed quality for livestock.

The species under study exhibited substantial differences in their nutritional composition, with the formulations SS₁ and SS₄ showing better overall digestibility and energy content, as reflected by their higher Total Digestible Nutrients (TDN) and Digestible Dry Matter (DDM) values. This suggests these species may be more suitable for promoting livestock growth and improving feed efficiency. On the other hand, SS₅ demonstrated higher levels of crude fiber and lower digestibility, making it less optimal for efficient feed conversion and weight gain.

In terms of anti-nutritional factors, SS₁ exhibited the highest levels of saponins and nitrates, both of which can negatively impact animal health if consumed in excess. The presence of these compounds in feed ingredients can potentially lead to reduced feed intake and digestive issues. The study also found variations in phenol and tannin content, with formulations such as SS₂ and SS₅ showing moderate levels, impacts digestibility and overall feed efficiency. These findings support the need for careful selection and processing of feed materials to minimize the negative impacts of these compounds on livestock performance.

Overall, the study emphasizes the importance of balancing nutritional composition with anti-nutritional factors to optimize livestock feed. The feed formulations with lower levels of anti-nutritional compounds, like SS₂, offer more sustainable and cost-effective solutions for livestock nutrition. Moreover, future research should focus on understanding the precise mechanisms by which these anti-nutritional factors interact with livestock physiology, as well as exploring methods to mitigate their impact through feed processing or supplementation.

Acknowledgements

We express our heartfelt gratitude to the esteemed contributors from Tamil Nadu Agricultural University and Tamil Nadu Forest Department for their invaluable support in the submission of the research article titled "Nutrient-rich or anti-nutritional challenge? Evaluating *Senna spectabilis* (DC.) H.S. Irwin & Barneby for sustainable animal feed production"

We extend our deepest appreciation to:

- Dr. V. Geethalakshmi, Vice Chancellor, Tamil Nadu Agricultural University, for her visionary leadership and

unwavering support.

- Dr. K. Baranidharan, Professor (Forestry) and Head, Department of Forest Products and Wildlife and Thiru. Nihar Ranjan, IFS, Dean (Forestry) - Forest College and Research Institute, Mettupalayam, for his profound insights and expertise.

Their contributions have been instrumental in enriching this work and advancing the field of Forestry and Agriculture. Profound thanks to Tamil Nadu Forest Department for funding the Research work

Authors' contributions

BS worked in the collection of the literatures and writing the manuscript. BK guided in writing the manuscript and bringing it to a sequence. TM assisted in the correction of the manuscript. RR assisted in additional inputs that was needed in the manuscript. GKN, RKP and VM provided an outline and interpretation of the manuscript. RR and HP assisted in correcting the manuscript. KV and SD helped in reviewing and editing the manuscript. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: The authors declare that there is no conflict of interest.

Ethical issues: None

References

1. IUCN. *Senna spectabilis*: Groom, A: The IUCN red list of threatened species 2012: E.T19892105A20141165. IUCN Red List of Threatened Species. IUCN; 2010.
2. Kumar A, Prasad S. Threats of invasive alien plant species. *Int Res J Manag Sci Technol*. 2014; 4:605-24.
3. Hobbs HA. *Invasive species in a changing world*. Island press; 2000.
4. Raizada P, Raghubanshi AS, Singh JS. Impact of invasive alien plant species on soil processes: A review. *Proceedings of the National Academy of Sciences India Section B-Biological Sciences*. 2008;78(PART 4):288-98.
5. Vitousek PM, D'antonio CM, Loope LL, Rejmanek M, Westbrooks R. Introduced species: a significant component of human-caused global change. *New Zealand Journal of Ecology*. 1997:1-6.
6. Rejmánek M, Richardson DM. Trees and shrubs as invasive alien species-2013 update of the global database. *Diversity and Distributions*. 2013;19(8):1093-4. <https://doi.org/10.1111/ddi.12075>
7. Rejmánek M, Richardson DM, Pyšek P. Plant invasions and invasibility of plant communities. *Vegetation Ecology*. 2013;387-424. <https://doi.org/10.1002/9781118452592.ch13>
8. Gaertner M, Den Breëyen A, Hui C, Richardson DM. Impacts of alien plant invasions on species richness in Mediterranean-type ecosystems: a meta-analysis. *Progress in Physical Geography*. 2009;33(3):319-38. <https://doi.org/10.1177/030913330934160>
9. Hejda M, Pyšek P, Jarošík V. Impact of invasive plants on the species richness, diversity and composition of invaded communities. *Journal of Ecology*. 2009;97(3):393-403. <https://doi.org/10.1111/j.1365-2745.2009.01480.x>
10. Irwin HS, Barneby RC. The American Cassiinae: a synoptical revision of Leguminosae tribe Cassieae subtribe Cassiinae in the new world. 1982.
11. Satyanarayana P, Gnanasekaran G. An exotic tree Species *S. spectabilis* (DC.) Irwin & Barneby (Caesalpiniaceae) naturalized in Tamil Nadu and Kerala. *Indian Journal of Forestry*. 2013;36(2):243-6. <https://doi.org/10.54207/bsmps1000-2013-fbbsv3>
12. Garden MB. Tropicos database. St. Louis, Missouri, USA: Missouri Botanical Garden [Internet]. 2016.
13. Harilal K. Impact of invasive alien Plants on understorey vegetation in Tholpetty range of Wayanad Wildlife Sanctuary (Doctoral dissertation, Department of Natural Resource Management, College of Forestry, Vellanikkara). <http://hdl.handle.net/123456789/7993>
14. Anoop NR, Sen S, Vinayan PA, Ganesh T. Native mammals disperse the highly invasive *Senna spectabilis* in the Western Ghats, India. *Biotropica*. 2022;54(6):1310-4. <https://doi.org/10.1111/btp.12996>
15. Lukosi N. <News> A Brief Note on Possible Control of *S. spectabilis*, an Invasive Exotic Tree at Mahale. *Pan Africa News*. 1997;4(2):18.
16. Vinayan PA, Anjankumar BN, Vishnu NM, Vaishnav K, Unais P, Ajayan PA, et al. Mapping the distribution and abundance of the exotic invasive species, *Senna spectabilis* in the Wayanad Wildlife Sanctuary, Kerala. *Ferns-A society for nature conservation & Kerala Forests and Wildlife Department*. 2020.
17. Garba Y, Muhammad IR, Adnan aa. Common fodder fed by small ruminants of the agro-pastoral production system in semi-arid, Nigeria. *Proceedings of the 1st International conference on Drylands*. 2014.
18. Duarte-Vargas JH, Melo O, Mora-Delgado J, Castañeda-Serrano R, Váquiro H. Pod production and dasometric variables, of the tree *S. spectabilis* (Fabaceae) in a tropical dry forest. *Revista de Biología Tropical*. 2021;69(1):218-30. <https://doi.org/10.15517/rbt.v69i1.42792>
19. Kumar S, Singh P, Devi U, Yathish KR, Saujanya PL, Kumar R, et al. An overview of the current fodder scenario and the potential for improving fodder productivity through genetic interventions in India. *Animal Nutrition and Feed Technology*. 2023;23(3):631-44. <https://doi.org/10.5958/0974-181X.2023.00054.9>
20. Helrich K. *Official methods of analysis*; 1990.
21. Schroeder JW. *Interpreting forage analysis*; 1994.
22. Redfearn DD, Zhang H, Caddel JL. *Forage quality interpretations*. Division of Agricultural Sciences and Natural Resources, Oklahoma State University; 2004.
23. CI KC, Indira G. Quantitative estimation of total phenolic, flavonoids, tannin and chlorophyll content of leaves of *Strobilanthes kunthiana* (Neelakurinji). *J Med Plants*. 2016;4:282-6.
24. Cataldo DA, Maroon M, Schrader LE, Youngs VL. Rapid colorimetric determination of nitrate in plant tissue by nitration of salicylic acid. *Communications in Soil Science and Plant Analysis*. 1975;6(1):71-80. <https://doi.org/10.1080/00103627509366547>
25. Hiai S, Oura H, Nakajima T. Color reaction of some sapogenins and saponins with vanillin and sulfuric acid. *Planta Medica*. 1976;29(02):116-22. <https://doi.org/10.1055/s-0028-1097639>
26. Cheng ZJ, Hardy RW. Effects of extrusion processing of feed ingredients on apparent digestibility coefficients of nutrients for rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Nutrition*. 2003;9(2):77-83. <https://doi.org/10.1046/j.1365-2095.2003.00226.x>
27. Syarif R, Halid H. *Food storage technology*. Inter-University Food and Nutrition, Bogor Agricultural University, Bogor, Indonesia. 1993.
28. Afolabi SS, Oyeyode JO, Shafik W, Sunusi ZA, Adeyemi AA.

- Proximate Analysis of Poultry-Mix Formed Feed Using Maize Bran as a Base. International Journal of Analytical Chemistry. 2021;2021(1):8894567. <https://doi.org/10.1155/2021/8894567>
29. Ukachukwu SN. Studies on the nutritive value of composite cassava pellets for poultry: chemical composition and metabolizable energy. Livestock Research for Rural Development. 2005;17(11).
 30. Julaikha Qistina I. Characterisation of Mulberry Leaf Pellet as Rabbit Feed (Doctoral dissertation, Universiti Malaysia Kelantan);2020.
 31. Buliah N, Jamek S, Ajit A, Abu R. Production of dairy cow pellets from pineapple leaf waste. InAIP Conference Proceedings 2019(Vol. 2124, No. 1). AIP Publishing. <https://doi.org/10.1063/1.5117108>
 32. Tillman BA, Pan WL, Ullrich SE. Nitrogen Use by Northern-Adapted Barley Genotypes Under No-Till. Agronomy Journal. 1991;83(1):194-201. <https://doi.org/10.2134/agronj1991.00021962008300010046x>
 33. Raut RG, Rekhate DH, Dhok AP. Nutrient utilization in goats fed arhar (*Cajanus cajan*) straw based complete feed pellets. Indian Journal of Animal Nutrition. 2002;19(2):135-9.
 34. Formato M, Scharenberg F, Pacifico S, Zidorn C. Seasonal variations in phenolic natural products in *Fagus sylvatica* (European beech) leaves. Phytochemistry. 2022;203:113385. <https://doi.org/10.1016/j.phytochem.2022.113385>
 35. Rajanish, Varshini. Validation of multi nutrient animal feed pellets using farm grown fodder trees with *Prosopis* pods. UG Project, Forest College and Research Institute, Tamil Nadu Agricultural University. 2022.
 36. Naumann HD, Tedeschi LO, Zeller WE, Huntley NF. The role of condensed tannins in ruminant animal production: advances, limitations and future directions. Revista Brasileira de Zootecnia. 2017;46:929-49. <https://doi.org/10.1590/s1806-92902017001200009>
 37. Oyaniran DK, Ojo VO, Aderinboye RY, Bakare BA, Olanite JA. Effect of pelleting on nutritive quality of forage legumes. Livestock Research for Rural Development. 2018;30(4).
 38. Lenz M. Risk of nitrate toxicity when grazing annual forages. 2018.