



REVIEW ARTICLE

Understanding the microbiome of ornamental plants: A comprehensive note

Nomita Laishram^{1*}, Arvinder Singh¹, Sourav Sharma¹, Shivam Kanungo¹, Ashutosh Sharma¹, Raj Kumar², Ritika Manhas¹, Shriya Gupta¹ & Kedar Mahadev Gheware¹

¹Division of Floriculture and Landscaping, Sher-e-Kashmir University of Agricultural Sciences and Technology, Chatha 180 009, Jammu & Kashmir, India

²Indian Agriculture Research Institute, Katrain, Kullu 175 129, Himachal Pradesh, India

*Correspondence email - nomitalaishram@gmail.com

Received: 18 March 2025; Accepted: 25 September 2025; Available online: Version 1.0: 26 November 2025; Version 2.0: 11 December 2025

Cite this article: Nomita L, Arvinder S, Sourav S, Shivam K, Ashutosh S, Raj K, Ritika M, Shriya G, Kedar MG. Understanding the microbiome of ornamental plants: A comprehensive note. Plant Science Today. 2025; 12(4): 1-13. <https://doi.org/10.14719/pst.8356>

Abstract

Ornamental plants are linked with a number of microbes present in rhizosphere, phyllosphere and endophytic region. Different plants host distinct microbiome communities. Many of these microorganisms are beneficial but some of them are pathogenic also. Beneficial microorganisms are engaged in putrefaction of organic matter, nutrient mineralization, nitrogen fixation, phosphate solubilization and development of plant-growth hormones. They also help the plants in restricting the development of pathogenic organisms and to withstand against various biotic and abiotic stresses. This review presents an extensive literature search on microbes of ornamental plants. The knowledge of microbiome and its beneficial effect on ornamental plants will be helpful in development of microbial formulations for sustainable flower production which will further lessen the utilization of agrochemicals and maintain ecological balance. Studies on microbiome-based sustainable practices in ornamentals, though promising, are poorly understood.

Keywords: ecosystem; microbiome; ornamental plants; plant nutrition; stress mitigation

Introduction

Plant microbiome is defined as an assembly of microorganisms which are interrelated with rhizosphere, phyllosphere and endosphere of higher plants. Interaction of these microbes with the plants constitutes the microbial ecosystem and play a vital role in plant nutrition as well as increased resilience to plant stresses, stress tolerance and productivity (1). The plant microbiome comprises not only fungi and bacteria, but also some microorganisms like viruses, algae, archaea and protozoa also exist. Microbial colonization in the plant rhizosphere is highly influenced and controlled by the composition of the host-specific exudates which further modify the physico-chemical properties of the soil and thus the microbiome. A few bacterial and fungal species have been isolated from ornamental plant tissues. Some of the microbes termed as “obligate symbionts” cannot survive without a host plant and includes fungi, viruses and some bacteria. The host plant plays its part by providing oxygen, space, carbohydrates and proteins to the microbes and the microbe in turn helps in improving the plant health by developing plant hormones, thus enhancing uptake of nutrients and solubilization of phosphate. Also, they help the plants in suppressing the growth of plant pathogens by producing antibiotics, degrading fungal cell wall enzymes and suppressing siderophores activity (2, 3).

Diversity in microbiome of ornamental plants

The microbiome of ornamental plant is vast and their habitats include the rhizosphere, phyllosphere as well as the endosphere

where microbes engage in an associated, endophytic or epiphytic manner (Fig. 1). When microbes adhere to the plant tissues they are termed as epiphytes whereas when the microorganisms are positioned inside plant tissues then they are termed as endophytes. The rhizospheric microflora comprises of fungi, nematodes, algae, bacteria, protozoa and microarthrops (4). Although only about 2 % of soil microbes can be cultured under laboratory conditions, the remaining 98 % cannot be cultured. It is because identifying, characterizing and describing the role of these uncultured microbes is difficult. Recently, evaluation of molecules from soil samples by using nucleic acid based techniques which included analysis of DNA and rRNA have uncovered enormous heterogeneity in the rhizosphere dwelling microbial flora (5).

A total of twenty-three different species of arbuscular mycorrhizal fungi from rhizospheric soil of various ornamental plants pertaining to five genera i.e. *Acaulospora*, *Entrophospora*, *Gigaspora*, *Glomus* and *Sclerocystis* (6). Among all the genera, most dominant was *Glomus* followed by *Acaulospora*. Among different species of *Glomus*, the most profuse and frequent species reported in various ornamental flowering plants was *G. mosseae* and among the *Acaulospora* species, *Acaulospora laevis* was the most abundant.

Rhizospheric microbiome

The plant root system provides unprecedented ecological niches for soil microflora which inhabit the rhizosphere, roots and to some extent above ground parts. A broad range of bacterial




Fig. 1. Microbial habitats and interactions in ornamental plants.

endophytes colonize the root tissues of plants. Rhizobium bacteria, mycorrhizae, biocontrol microbes and plant growth promoting rhizobacteria (PGPR) are the most often studied rhizosphere organisms that are beneficial to the plants. Most profuse members of the rhizosphere population are mycorrhizal fungi, as they have been found in association with over 200000 species of plants i.e., over 80 % of all the plant species (7). Various rhizosphere bacterial taxa especially relating to *Pseudoxanthomonas* having significant divergent abundances among 30 plant species in angiosperms (8). Several authors described that the roots of *Rosa hybrida* were highly colonized by *Glomus intradices* (9-11).

High levels of phosphorus fertilization rates can lead to a decline in colonization rates of arbuscular mycorrhiza in rose plants (9). However, it was observed that mycorrhizal symbiosis in *Tagetes* and *Zinnia* plants is independent of changes in nutrient levels (12). Plants inoculated with *Glomus fasciculatum* showed higher percentage of colonization in chrysanthemum cultivars “Shyamal”, “Local Yellow” and “Bangalore White” than other species of mycorrhiza (13). When referring to the growth parameters such as number of laterals and plant height in all the cultivars, effect of VAM fungi viz., *Acaulospora laevis*, *Glomus fasciculatum* and *Glomus mosseae* were found to be stimulating in comparison to the control. Colonization by *Glomus fasciculatum* was also associated with the highest flower yield in chrysanthemum (14).

Phyllosphere microbiome

The phyllosphere of the ornamental plants provide exceptional habitat for both endophyte as well as epiphyte diversities. Endophytes usually move systemically through xylem to different parts of the plant like fruits, leaves, flowers and stem (15). These above-ground plant microbiome usually originates from soil, air and seed and are infamously known for their role in promoting disease resistance; alleviate stress tolerance and promoting plant growth. Population of fungal communities are significantly

variable in the phyllosphere of the temperate regions and are usually diverse in tropical regions (16).

The exudations from pollen provide an attractive niche for microbial colonizers. These colonizers include both fungi (*Penicillium* and *Botrytis* molds) and bacteria (*Bacillus* sp.) (17). The primary site of infection for the pathogenic bacterium *Erwinia amylovora* is thought to be stigma; because of the numerous economic losses interrelated with floral infection caused by this pathogen (18).

Recent studies have recognized novel microbial taxa isolated from flowers, highlighting various ascomycetous yeasts belonging to *Wickerhamomyces* and *Candida* which signify that flowers might be an untapped reservoir of microbial diversity (19). Furthermore, the population densities of two epiphytic bacterial communities of petals and leaves of two ornamental plants, *Lotus corniculatus* L. and *Saponaria officinalis* L. were compared and reported that petals and leaves of both the plant species had different community formations (20). Communities from the petals were predominated by family Enterobacteriaceae which had lesser diversity as compared to the population of leaf communities. However, petal communities of two plants were reported to be similar and common taxa comprised the genus *Serratia*. In another study, culturable yeasts were examined from different parts of flowers and reported maximum number of yeast species found in the inner as well as outer corolla and the most abundant species on the surface of plant were reported to be basidiomycetous yeast species (21).

Endosphere microbiome

The endophytes invade and occupy the internal tissues of plants thus creating the endospheric microbiome. The arbuscular mycorrhiza as well as other endophytic fungi are the predominant dwellers of the endosphere. Bacteria are also important endophytes. The diversity and identity of the endophytic microbiome of both the above as well as below-ground tissues varies within the plant.

Factors influencing ornamental plant microbiome composition

Host factors

The host plant's identity plays a crucial role in shaping its unique microbial community. Plant species that are distantly related phylogenetically show greater variation in microbiome composition (22). When different plant species grow adjoining to one another they can harbour different microbiomes. All three factors i.e. host, environment and microbe are inter-linked and they together build the community of microbiome in plants.

Plant species that grow in closely resembling soil environment harbour distinct microbial communities in root compartments as well as rhizosphere. Different genotypes of plants behave differently in respect to interaction with microbiota that attract different microbiome members which confer resistance to biotic as well as abiotic stresses and may help for plant growth and nutrition (23).

Microbiome composition also varies with plant's age, health and surrounding soil environment. Sometimes different species of plants can harbour the same microbiome composition with variation in level of colonization. For instance, as mentioned in the earlier studies 71 % mycorrhizal root colonization was observed in snapdragon (24), 50 % in *Zinnia* and *Tagetes* (12) and 40 % in chrysanthemum (25).

In cases of relationship between pathogens and plants, age-related resistance (onto genic resistance) is highly expressed and correlated with developmental stage of the plant (26). Symbiosis research also indicated that the age and developmental stage of the plant are important factors that affect the microbial communities (27).

The arbuscular mycorrhizal fungal status of fifteen different ornamental flowering plants was investigated in 2012 (6). They observed that percent root colonization by the fungi varies from 14.28 % to 100.00 %. Maximum root colonization and spore count was observed in *Gladiolus grandiflorus* (172.00 ± 4.35) and *Senecio cineraria* (100.00 ± 0.0), respectively.

Some microorganisms produce antibiotics such as penicillin and polymyxin which increase the leakage, exudation of organic materials and altered cell permeability of roots (28) resulting in variable assembly of microbiome.

It is also documented that inoculation with arbuscular mycorrhizal fungi expressed varied colonized status in roots and production of biomass in *Hycinthus orientalis* L. Anna Marie. Only *Funneliformis mosseae* showed the greatest root colonization among other AM fungi species which confirmed that *Funneliformis mosseae* had the highest compatibility with the hyacinth plant species for greater efficiency (29).

Environmental factors

The formation of plant microbiome is highly affected by a range of abiotic and biotic factors. The rhizospheric microbiome is largely influenced by salinity, pH, structure, type, organic matter and moisture content of soil (30) whereas environmental factors like

climate, presence of pathogen, geography and farming practices influence both rhizospheric and phyllospheric microbiota (31).

Land use activities and cultivation practices significantly contribute to the decline in microbial biodiversity (32). Change in the vegetation highly affects the structure and diversity of soil microbiomes. Continuous cultivation leads to the changes in soil properties mainly nutritional, which in turn affects the communities of soil microbiome directly. In prolonged field experiments, it was summarized in 2009 (33) that differently managed soils harboured distinct microbiomes.

In natural conditions, plants and their microbiota when exposed to different environmental conditions, such as changes in humidity, temperature, UV rays and pH, directly or indirectly affect the microbiome composition (34). A study in different ecosystems in Britain demonstrated that temperature and rainfall gradients are major climatic factors that shape the composition of bacterial community in plants (35). Drought is another factor that affects the microbiome composition due to altered oxygen availability and nutrient mobility in soil (36).

Role of microbiomes in ornamental plants

The microbiome of ornamental plant consists of harmful as well as beneficial microorganisms. Beneficial microorganisms of the rhizosphere are involved in disintegration of organic matter, primary productivity and nutrient mineralization (37). They are also involved in fixation of nitrogen (N_2), solubilization of phosphate and manufacturing of plant-growth hormones (38). Some of them help the plants to fight pathogen attack and withstand abiotic and biotic stresses. Exudates released in rhizosphere from plants or from microbes help the plants to communicate with microorganisms for assistance in environmental acclimation to mitigate stresses. The plant exudates are further utilised by microorganisms as a resource for carbon and other nutrients. The microbiome also modifies root architecture of plant (39), by releasing plant growth hormones. Harmful microbiota on the other hand acts as pathogens causing diseases on important crop plants leading to economical loss (Fig. 2).

Biofertilization and bio-stimulation

Biological modifications in soil leading to improving the efficacy of soil microbial community is known as "biofertilization". Biofertilizers are microbial inoculants or biologically active products comprising of one or more beneficial bacteria or fungal strains which conserve, add and mobilize crop nutrients in the soil (Table 1).

Excessive use of chemicals fertilizers in the cultivation system reduced soil microbial population which declines soil physico-chemical properties. Further, these applied fertilizers are lost through volatilization, denitrification, leaching and phosphorus fixation. Consequently, favouring the use of bio-fertilizers for sustainable farming.

Efforts in early 1970's were made to specify the microflora of English daisy, marigold and balsam (40, 41). The beneficial role played by biofertilizers in ornamental plants like jasmine, rose, marigold, chrysanthemum, aster, dahlia, gladiolus, tuberose and lilies (42).

Table 1. The most popular types of bio-fertilizers used in ornamental plants

Sl. No.	Biofertilizers	Examples
1.	Nitrogen (N_2) fixers	<i>Azotobacter</i> and <i>Azospirillum</i>
2.	Phosphorous solubilizers	<i>Bacillus</i> and <i>Pseudomonas</i>
3.	Phosphorous Mobilizers	<i>Gigaspora</i> sp. and <i>Acaulospora</i> sp.
4.	Zinc and silicate solubilizers	<i>Bacillus</i> sp.



Fig. 2. Unveiling the dual nature of microbiomes in plants.

Plant growth promoting rhizobacteria persistently colonize the roots of plants that augment plant growth through mineral solubilization. They also support establishment of other beneficial soil microbes, such as mycorrhizal fungi and help in suppressing the growth of plant pathogens (43).

When phosphorus-solubilizing bacteria is used in combination with Arbuscular mycorrhiza it leads to synergistic effect because mycorrhizal exudations directly affect bacterial communities thus increasing water absorption by host plants (44). The synergistic effect of *Glomus fasciculatum* and *Bacillus subtilis* increased inflorescence numbers by 14-24 % and significantly higher fresh weight in *Tagetes erecta* cv. "Alcosa" compared to untreated plants (45). Similar reports on the combination of *Acaulospora laevis* and *Pseudomonas fluorescens* in enhancing root biomass, phosphatase activity and phosphorus uptake of *Chrysanthemum indicum* L. were documented in the earlier studies (46) and in *Gazania rigens* L. (47). A consortium of *Pseudomonas putida* and *P. fluorescens* strains improves growth and flowering in *Pelargonium peltatum*, *Chrysanthemum* sp. and *Dahlia variabilis* (48).

Some of the researches have highlighted the beneficial effect of *Azotobacter* in crops like *Tagetes erecta*, *Rosa hybrida* (49, 50). Yet others are of the opinion that the interaction of *Azotobacter* and phosphorus solubilizing bacteria are more beneficial in crops like *Gladiolus grandiflorus*, *Polyanthes tuberosa* and *Matthiola incana* than application of *azotobacter* alone (51-53).

Azospirillum is another important microbe used in ornamental crops which is reported to increase growth, early flowering and nutrient uptake in rose (54). A combination of

Azospirillum and phosphate solubilizing bacteria in petunia and carnation improved flower quality and yield (55, 56).

Arbuscular mycorrhizal fungi live in symbiotic association with plants. The root colonization, spore count, dominant species and species diversity varies with the soil nutrient conditions and region (57). This symbiosis can enhance plant growth, flower production, reduce days to flower emergence (58, 59), increase size and weight of flowers (25) and flower longevity (24, 60) in different flower crops. It was observed in rose that there was marked reduction in the transpiration rate of leaves detached from plants that were inoculated with *Glomus* thus suggesting increased longevity for harvested flower stems (10).

The beneficial effect of arbuscular mycorrhizal fungi in improving growth and flowering of annual flowers such as *Zinnia elegans*, *Salvia splendens*, *Tagetes patula*, *Callistephus chinensis*, *Impatiens balsamina*, *Petunia hybrida* and *Antirrhinum majus* was reported by several workers (61-65). Low supply of NPK augmented with mycorrhizal fungi is particularly helpful in increasing phosphorous content in tissues of *Pelargonium hortorum* (66).

AMF inocula applied directly to cuttings of *Chrysanthemum morifolium* shows enhanced rooting rate to 99 % while it was 77 % in non-mycorrhizal inoculated plants (67) and resulted in improved fresh weight and size of flower (25, 68).

It was recommended that for better functioning and establishment of symbiosis, rhizobacteria plays important role in assisting arbuscular mycorrhizal fungi (69). A combination of *Aspergillus niger* and *Glomus fasciculatum* was proved to be economical for flower production of chrysanthemum (70). Application of phosphate-solubilizing bacteria, vesicular arbuscular

mycorrhizas and free-living nitrogen-fixing bacteria in *Eustoma grandiflorum* is also reported to produce more flowers per plant and advanced the flowering by 14 days than application of these microbes individually (71).

Zephyranthes bulbs inoculated with AMF during the storage period showed enhanced growth and flowering in the subsequent growing cycle (72). Likewise, corms produced from AMF-inoculated *Brodiaea* plants contained a greater proportion of non-reducing sugars compared to non-inoculated plants (60). Some microbes also release enzymes like, 1-carboxylic acid-1-aminocyclopropane (ACC) deaminase, which leads to the reduced level of ethylene production in the plants (15).

Some studies have documented earlier flowering (73) with the application of AM fungi whereas others have reported delayed flowering (66, 74), or non-significant effect on flowering (75) due to AM fungi. Prolonged flowering duration (76) as well as reduced flowering duration (77) with AMF application are also reported. Under greenhouse rose production system, inoculation of *Glomus mosseae* resulted in reduction of time needed for flowering for 80 % plants by one month and exhibited increased number of cut flowers (73). Application of *Glomus fasciculatum* in *Gerbera jamesonii* cv. "Bolus" leads to 50 days earlier flower initiation (78).

Pathogenic interactions and host resistance in plant microbiomes

The intricate relationship between plants and their associated microbiomes involves a dynamic interplay with both beneficial and detrimental microorganisms. Plant root exudates, for instance, attract not only advantageous microbes but also potential pathogens (79). While plants exist in close proximity to various bacteria, fungi and viruses, successful infections are frequently averted due to the plant's inherent "non-host resistance" or "horizontal resistance" (80).

However, environmental factors significantly influence fungal disease development, particularly temperature and moisture. For example, leaf blight of marigolds caused by *Alternaria dianthi* is strongly correlated with rainfall, temperature and humidity (81). Fungal mycelia require specific temperature ranges and moist surfaces to survive, though spores can overwinter in harsh conditions, germinating when favourable temperatures and moisture return (82). Upon successful infection, pathogens compromise plant productivity by disrupting physiology and growth. Common fungal diseases include damping-off caused by *Pythium* spp. and *Phytophthora* spp. in crops like carnation and poinsettia; *Ceratobasidium* sp. was newly reported to cause damping-off in marigold in India (83). *Botrytis* spp. is notorious for infecting propagation material and causing substantial post-harvest losses in flowers during transport and storage (84). Specific reports include *Alternaria tenuis* causing black spot-on *Rosa centifolia* (85) and *Septoria* leaf spot on marigold (86). *Fusarium oxysporum* is a widespread pathogen affecting many ornamentals like *Chrysanthemum* spp., *Dianthus* spp., *Gerbera* spp., *Gladiolus* spp. and *Lilium* spp., with specific forma specialis targeting different hosts (87, 88). Its coalition with *Pythium* sp. can cause root rot in gerbera (89) and it also impacts other ornamentals such as *Anoectochilus* spp. and *Ranunculus* sp. (90). Bacterial diseases, driven by phytotoxic proteins and phytohormones, are also devastating, with pathogens like *Erwinia* sp., *Pseudomonas* spp. and *Xanthomonas* spp. maintaining infection cycles via insect vectors, vegetative parts or seeds. Viral diseases lead to significant economic losses, with transmission primarily by insects (leafhoppers, aphids, whiteflies,

thrips) and secondarily by nematodes, fungi and mites, as well as through infected plant parts or mechanical means.

Crucially, the plant microbiome plays an important role in inducing host resistance against these pathogens. Beneficial microbes, often existing in a dormant state, can be "reactivated" by pathogen attack to enhance resistance. This disease resistance, especially against soil-borne diseases, is influenced by the microbiome's composition, structure, functionality and assembly (91). Both rhizospheric and phyllospheric bacteria bolster host resistance through various biocontrol activities, including producing pathogen-inhibiting volatile compounds, siderophores, lytic enzymes and antibiotics (92). Some bacteria also enhance plant immunity by altering hormone levels and inducing systemic resistance (93). The plant microbiome further inhibits pathogen development by competing for resources, microenvironments and through antibiosis or parasitism (94).

Key bacterial genera known for protecting plants include *Pseudomonas*, *Streptomyces*, *Bacillus*, *Paenibacillus*, *Enterobacter*, *Pantoea*, *Burkholderia* and *Paraburkholderia* (94). Specific bacterial taxa like *Firmicutes*, *Actinobacteria* and *Acidobacteria* have been shown to control *Fusarium* wilt (95). For *Fusarium oxysporum*, effective biocontrol agents include *Bacillus*, *Fusarium*, *Glomus* and *Trichoderma* genera (96-98). *Pseudomonas fluorescens* is particularly significant, protecting roots from *Fusarium* sp. and *Pythium* sp. (99). Research indicates that healthy leaves of *Euonymus japonicus* harbour higher fungal and bacterial diversity compared to diseased ones, with powdery mildew altering the abundance of fungal (*Ascomycota*, *Basidiomycota*) and bacterial (*Proteobacteria*, *Firmicutes*) phyla (100).

Successful biocontrol examples abound: powdery mildew in rose can be controlled by *Ampelomyces quisqualis*, *Bacillus subtilis* and *Verticillium lecanii* (101), with *Gliocladium catenulatum* and *Bacillus subtilis* showing comparable efficacy to chemical treatment of mycobutanil (102). *Trichoderma viride* achieved 100% control of rhizoctonia stem rot in chrysanthemum (103) and *Trichoderma viride* and *Trichoderma harzianum* offered 63-73 % control of the same disease in carnation (104). *Trichoderma harzianum* and *Pseudomonas fluorescens* significantly inhibited fusarium wilt in carnation (105, 107, 108), with *Trichoderma asperellum* also reducing disease severity (106). *Pseudomonas fluorescens* and *Trichoderma viride* effectively reduced fusarium wilt in gerbera (109). Additionally, *Glomus mosseae*, *Trichoderma harzianum*, *T. viride* and *Bacillus subtilis* control *Fusarium* wilt in gladiolus (110-112). *Minimedusa polyspora* reduced the incidence of *F. oxysporum* f. sp. *narcissi* by 33% in narcissus (113). *Botrytis cinerea* sporulation was reduced by over 90 % using *Gliocladium roseum* FR136 and *Trichoderma inhamatum* (114) and leaf rot in cyclamen was mitigated by *Ulocladium atrum* and *Gliocladium roseum* (115). Furthermore, *Glomus mosseae*, *G. intraradices*, *Pseudomonas putida*, *Pseudomonas aureofaciens* and *Streptomyces* sp. delayed chrysanthemum yellows phytoplasma infection (116). Finally, *Streptoverticillium* sp., *Trichoderma* sp. and *Gliocladium virens* controlled collar rot in marigold (117) and *Trichoderma asperellum* successfully remediated *Phytophthora ramorum* infested nurseries (118) (Table 2).

Biocontrol of insect-pests

Intricate relationships exist among immune system and microbiome of plants which adds to the capability and versatility of the plant to deal with varying environments. The intrinsic immune system of the

Table 2. Biocontrol of diseases in ornamental plants

Disease in crop plant	Biocontrol agent	Beneficial effect	Reference
Powdery mildew of rose	<i>Ampelomyces quisqualis</i> , <i>Bacillus subtilis</i> , <i>Verticillium lecanii</i> and <i>Trichoderma harzianum</i>	Powdery mildew can be effectively control under the greenhouse conditions	(101)
Powdery mildew of rose	<i>Gliocladium catenulatum</i> and <i>B. subtilis</i>	Significant control of powdery mildew	(102)
Black spot incidence in rose	<i>Chaetomium globosum</i> , <i>T. harzianum</i> and <i>Bacillus subtilis</i>	Reduced disease index	(157)
Rhizoctonia stem rot in chrysanthemum	<i>Trichoderma viride</i>	No incidence of disease	(103)
Rhizoctonia stem rot in carnation	<i>Trichoderma viride</i> and <i>Trichoderma harzianum</i>	Disease control between 63-73 %	(104)
Fusarium wilt in carnation	<i>Trichoderma</i> , <i>Streptomyces</i> , <i>Pseudomonas</i> and <i>Bacillus</i> spp.	<i>Trichoderma harzianum</i> showed highest mycelial growth inhibition (87.78 %) followed by <i>Pseudomonas fluorescens</i> with 86.67 % growth inhibition of mycelia	(105, 106, 158)
Fusarium wilt in carnation	<i>Trichoderma viride</i> and <i>Trichoderma harzianum</i>	The beneficial effect of using <i>Trichoderma viride</i> and <i>T. harzianum</i> for disease control is reported	(107, 108, 159)
Fusarium wilt in gerbera	<i>Pseudomonas fluorescens</i> and <i>Trichoderma viride</i>	Significantly reduced the wilt incidence	(109)
Fusarium wilt in gladiolus	<i>Glomus mosseae</i> and <i>G. etunicatum</i>	Check the infection of <i>Fusarium</i> yellows	(110-112)
Anthrachnose of cyclamen	AM fungi	Decrease the anthrachnose incidence	(127)
Color rot of marigold	<i>Streptovericillium</i> sp., <i>Trichoderma</i> sp. and <i>Gliocladium virens</i>	Effectively control collar rot of marigold caused <i>Rhizoctonia solani</i> , <i>Phytophthora</i> sp. and <i>Sclerotium rolfsii</i>	(117)
Phytophthora infested nnursery plants	<i>Trichoderma asperellum</i>	Infested nurseries were successfully remediated under greenhouse conditions	(118)

plant solitarily is usually not adequate to fight against the antagonists in the phyllosphere. The microbiome of ornamental plants can also contribute to naturally suppressing the pests. The communities of microorganisms in the endosphere, leaf surfaces and rhizosphere protect the plants against the various biotic stresses such as fungal pathogens and insect herbivory (119, 120).

Amelioration of abiotic stress

Plants undergoing various abiotic stresses engage advantageous microorganisms present in the environment by utilising a wide array of chemical stimulus to improve the capability of plants to ward off stresses. Abiotic stress such as salinity, drought, heat and mineral toxicity increases the population of specific microbial symbionts associated with plants by the following mechanisms viz., (i) change in immunity and plant physiology, which modifies the secretion profile of exudations from the plant roots and favour the growth of specific microbes associated (ii) directly alter the properties of soil properties viz., pH, O₂ and nutrient levels and communities of microbes by accumulation of specific microorganisms in the roots of the plants e.g., drought and flooding events favour betaproteobacteria and actinobacteria in soils (121) (iii) increase the population density of specific microorganisms due to nutritional interdependence (122).

Alterations in the microbiome of roots generally improve tolerance of plants during stress period. For example, when leguminous plants were grown in limited nitrogen (N), plants released higher proportions of flavonoids which attracted N-fixing bacteria (123). Studies also revealed that 1-aminocyclopropane-1-carboxylate (ACC), an amino acid present in the plants which ameliorates the stress conditions, could reorganize microbiome population which enhanced tolerance of plants to the salinity stress conditions (124).

Various exudations from the roots attract microorganisms during the stress period. Microbes retort swiftly to variations in root exudations. Abiotic stresses like metal toxicity, poor nutrition, drought and light limitation alters the microbial community and the plant root metabolism. In some instances, increase in the specific population of microorganisms during the stress-induced environments could improve the stress tolerance of plants (125). Drought increases the population of actinobacteria in the

endosphere and/or rhizosphere of the plants. Likewise, limiting nitrogen/phosphate supply of improves the symbiosis of plants with advantageous microorganisms for acquisition of nutrients (8).

Temperature stress

Significant rise in temperatures could influence the plant physiology by increasing leaf transpiration rate and respiration and by varying the allocation of photosynthates. While, low temperature regimes reduce the metabolism of plants, which leads to reduction in the foliar growth, early senescence and reduced photosynthetic activity. Temperatures lower than freezing point hampers the development of buds because the growing and rehydrated buds get destroyed due to cold stress.

Microbial biostimulants boost the response of plants to heat stress. For instance, the production of ROS-damaging enzymes e.g. peroxidases, catalase, superoxide and dismutase, improve stress tolerance of plants to heat and it can be improved by colonizing with beneficial bacteria, such as *Pseudomonas* and *Bacillus* and mycorrhizal fungi such as *Septoglomus deserticola* and *Septoglomus constrictum* (126). For example, AM fungi enhance the growth under heat stress condition in cyclamen (127).

By utilizing microbes that inhibit ethylene synthesis helps the plants in escaping the damaging effects of heat stress (128). Specifically, bacteria which show activity of 1-aminocyclopropane-1-carboxylic acid (ACC), are known to be highly effective in surviving stress conditions such as *Bacillus subtilis* BERA 7, *Leclercia adecarboxylata* MO1, *Pseudomonas fluorescens* YsS6 and *Pseudomonas migulae* 8R6 (129).

For overcoming stress due to low temperatures, microbes which possess ice nucleating activity (INA⁺), can encourage production of ice cores in plants and the first bacterial strain that was stated to wield ice nucleating activity was *Pseudomonas syringae* (130). Other microbes which show ice nucleating activity are *Erwinia herbicola* (syn. *Pseudomonas agglomerans*), *Xanthomonas campestris* and *Lysinibacillus* sp. (131).

Salinity and drought stress

Drought and salinity induce water stress in plants which takes place when water losses due to leaf transpiration surpass the water absorbed by roots, resulting in reduced turgor in plant tissues. Stress

caused by the drought conditions highly influences the plant morphology and physiology, leading to buildup of harmful ROS (132), ethylene emission and reduced availability of mineral nutrient, absorption and transport (133). Various direct and plant-mediated mechanisms involving microbial biostimulants improve plants tolerance to drought and salinity stress such as, *Pseudomonas* spp. PF23 and *Pseudomonas putida* GAP-P45 help in producing bacterial exopolysaccharides which improve structure of soil by creating micro and macroaggregates, which in turn improve the development of plants in water stressed conditions (134).

Mycorrhizal fungi increase root biomass, improves water retention, reduces leaching of mineral nutrients (135) and increase the tolerance of plants to soil salinity. The improvement in nitrogen uptake is considered as the major mechanism which induces salt tolerance in plants (136). *Glomus* sp. produces a glycoprotein known as glomalin, which helps in improving plant tolerance to drought (137). Arbuscular mycorrhizal fungi protects the plants from drastic effects of drought which might be due to an increase in hydraulic conductivity of roots, transpiration rate or stomatal regulation, increased uptake of water at lower levels soil moisture because of highly protruding hyphae (138), osmotic adjustment which helps in maintaining turgor pressure of plants even at lower tissue water potential (9), proline and carbohydrate accumulation, increased photosynthetic activity and nutrient availability (139).

Glomus constrictum Trappe was reported to induce drought stress tolerance in marigold (*Tagetes erecta*) (140). Similar report on mycorrhizal symbiosis improving the photosynthetic yield and water status of rose plants under drought stress have been documented (9). Enhanced plant growth as well as yield following AM inoculation in marigold due to improved uptake of P and Cu under water-stressed conditions was also reported (141).

The genus *Pseudomonas* is well known for its ability to increase plant growth in low-nutrient and drought conditions. Other genera like *Pseudomonas*, *Bacillus*, *Enterobacter*, *Agrobacterium*, *Streptomyces*, *Klebsiella* and *Ochromobacter* improves the production of wide range of crops under saline conditions (142). Increased number of flowers and biomass in *Petunia hybrida* and *Impatiens walleriana* plants grown under low-nutrient and drought conditions with the inoculation of *Pseudomonas poae* 29G9 and *Pseudomonas fluorescens* 90F12-2 was reported (143). Similarly, increase in plant growth, shoot biomass and growth index in greenhouse-grown zinnia and petunia at low fertility conditions was also reported with the inoculation of *Bacillus subtilis* QST 713 (144).

Some bacteria protect plants against different types of stress, which might be due to physiological-adaptive responses such as producing osmoprotective metabolic compounds; producing hormones which promote root growth of plants under stress; producing bioactive and antioxidant molecules. *Bacillus licheniformis* SA03 isolated from chrysanthemum plants grown in saline-alkaline soil of China conferred increased salt tolerance in chrysanthemum (145).

A previous study reported that *Crassula ovata* plants inoculated with *Glomus* spp. show more resistance to salt stress than non-inoculated plants by limiting sodium ion uptake at the cellular level (146). She further stated that the protective effect of microorganisms against salt toxicity is achieved through production of exopolysaccharide matrices, production of osmolytes,

biostimulation of root development and production of ACC deaminase.

An earlier study reported that under high salinity levels (6 dS/m), *Entrophospora* sp. fungi could invade and inhabit the chrysanthemum plant roots effectively thereby significantly improving flowering and yield (147). Other mycorrhizal fungi like *Funnelliformis mosseae* and *Diversispora versiformis* improve shoot and root dry weight, root length and root nitrogen concentration in *Chrysanthemum morifolium* plants cultivated for five months under moderate salinity conditions (50 mM NaCl). As salinity increased to 200 mM NaCl, mycorrhizal dependence and mycorrhizal colonization decreased due to the adverse effects of salinity on spore germination or impaired hyphal growth (136). A previous findings suggested that rhizosphere microbiome promote growth and germination of *Hibiscus hamabo* in salinity stress (148).

Preparation of microbial formulation and its application

The microbes thrive in their natural environments as compared to the ones introduced to new environments because they cannot compete with the indigenous microbes. The competitiveness of the introduced strain is affected by the physiological activity as well as concentration or quantity of the strain of the strain that is introduced (149). To secure that the cells are delivered at the appropriate positions with extended shelf-life appropriate solutions should be made. For example, for a specific microorganism to perform biocontrol functions, it is of utmost importance that the colonized strain establishes itself in the same niches as of pathogenic organisms and show antagonistic activities (150).

Combination of microbes with various characteristics are now commonly used as some microbes complement one another to cooperate various processes required for specific results such as biocontrol of pathogens and plant growth enhancement. Many a times, it has been observed that by combining bacteria that exhibit no, or very minute plant growth promotion impacts when used alone can exhibit plant growth promotion results when used in combinations, varying from combining three different species of bacteria that thrive in one biofilm (2) to applying of the whole microbiota (151). Efficiency of microbial inoculants relies on the inoculation characteristics and the capability of the inoculum to surpass the native microbiota competition and establish itself in the rhizosphere.

Formulations ensure that the cells remain viable during the storage period and are also available to the plants grown in fields. However, viability of these formulations depends on the capability of these microbes to tolerate low humidity (115).

The bioinoculants which have greater capability to invade, set up themselves in the rhizosphere of the host, they also create structural changes in the microbial community of native species (152). When arbuscular mycorrhizal fungi were introduced in salvia (*Salvia officinalis*), it was shown to alter the makeup of the bacterial community associated with the rhizosphere (153).

Pyrosequencing the 16S rRNA gene reveals variation in bacterial community in the rhizosphere of chamomile. The use of distinct bioinoculants where the bacteria related to phylum Verrucomicrobia existed when rhizosphere was treated with *Stenotrophomonas rhizophila* (154).

Table 3. Role of microbiome in crop productivity of flower crops

Crop plant	Microorganism	Beneficial effect	Reference
<i>Zinnia elegans</i>	AMF	Growth and overall yield	(61)
Crossandra	AMF	Increase in weight of flowers	(68)
Rose	<i>Glomus mosseae</i>	Time to flowering reduced by one month	(73)
<i>Antirrhinum majus</i>	<i>Glomus deserticola</i>	Increase in flower size	(64)
Marigold	AMF	Increased the size of the flowers	(140)
<i>Antirrhinum majus</i> and <i>Hypericum perforatum</i>	AMF	Enhanced the number of flowers	(64, 65)
<i>Rosa hybrida</i>	<i>Azotobacter</i>	Increased leaf area and plant height	(50)
<i>Tagetes erecta</i>	<i>Azotobacter</i>	Maximize flower yield	(49)
<i>Petunia hybrida</i>	<i>Azospirillum</i> sp. + PSB	Improved flower quality and increase flower yield	(155)
Carnation	<i>Azospirillum</i> + PSB	Enhance flower fresh weight	(55)
<i>Tagetes erecta</i> cv. "Alcosa"	<i>Bacillus subtilis</i> and <i>Glomus fasciculatum</i>	Produced 14-24 % more inflorescences and flowers had higher fresh weight	(45)
<i>Chrysanthemum indicum</i> L.	<i>Acaulospora laevis</i> + <i>Pseudomonas fluorescens</i>	Increased plant height, fresh and dry root weight	(46)
Potted hyacinth (<i>Hyacinthus orientalis</i> L. Anna Marie) plants	<i>Funneliformis mosseae</i>	Greater raceme length and biomass production of single floret, raceme and flower stem	(29)
<i>Gazania rigens</i> (L.) Gaertn	<i>Funelliformis mosseae</i> , <i>Acaulospora laevis</i> and <i>Pseudomonas fluorescens</i>	Early flowering, improved flower head size, flower fresh and dry weight, total chlorophyll and carotene content	(47)
Tulip	<i>Azotobacter chroococcum</i> and <i>Azospirillum lipoferum</i>	Increased plant height, leaf area, tepal diameter, growth and quality of flower as well as bulb yield	(156)
<i>Gladiolus grandifloras</i>	<i>Azotobacter</i> + Phosphorus solubilizing bacteria	Early corm sprouting, increase in corm weight and number of corms/plants	(51)
<i>Polyanthes tuberosa</i>	<i>Azotobacter</i> + PSB	Increased bulb production	(52)
<i>Pelargonium peltatum</i> , <i>Chrysanthemum</i> sp.	<i>Pseudomonas putida</i> and <i>P. fluorescens</i> strains	Increases plant size and flower number of plants	(48)
<i>Petunia × hybrida</i>	<i>Azospirillum lipoferum</i>	Increases flower number	(56)
<i>Matthiola incana</i>	<i>Azotobacter chroococcum</i>	Increase plant growth and nutrient content	(53)
<i>Eustoma grandiflorum</i>	VAM + PSB + free-living nitrogen-fixing bacteria	Produced more flowers per plant, advanced flowering by 14 days	(71)

Conclusion

Microbiomes of the ornamental plants are highly diverse and several factors are responsible for their functioning. The microbes interact in an associated, epiphytic, or endophytic manner with the ornamental plants. These microorganisms possess high potential to increase productivity and plant growth (Table 3). Their abundance as well as composition in the environment are governed by abiotic as well as biotic factors. They also play a significant role in mitigation of abiotic and biotic stresses. The most common utilization of the plant microbiome finds place in biofertilization and disease control. Biofertilization improves the functionality and interaction of the plant microbiome. Moreover, the application of microbial formulations helps in maintaining the ecological balance by reducing the use of chemical fertilizers and pesticides for cultivating ornamental plants. For commercialization of beneficial microbes in cultivating ornamental plants, research on the suitability and functionality of specific microbe for specific crop is critical. Understanding the role of the plant microbiome in enhancing plant productivity by enhancing nutrient acquisition, biocontrol against pathogens and pests, stress tolerance enhancement and improved soil health is of paramount importance. Isolation, selection and quantification of efficient microbes with better performance in the field can contribute a huge in increasing the production potential of ornamental crops. Research on the factors influencing plant and soil microbiomes is progressing rapidly. However, a key challenge lies in translating microbiome composition and diversity into predictable functional outcomes because of the

complexity of interactions of microbes and dependency on environmental factors too.

Acknowledgements

The authors acknowledge the Division of Floriculture and Landscaping, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu and the Research Station Katrain, Kullu, for providing the necessary support to complete this review paper.

Authors' contributions

NL and AS conceptualized the work. NL, AS, AS and SS prepared the original draft. NL and SS contributed to writing, review and editing. AS, SK and KMG handled visualization. and RK, SG and RM performed the editing. All authors have read and approved the final version of the manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None

AI Declaration : During the preparation of this work the authors used 'Napkin.ai' tool in order to support the drafting and structuring of the manuscript. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

References

- Mendes R, Garbeva P, Raaijmakers JM. The rhizosphere microbiome: significance of plant beneficial, plant pathogenic and human pathogenic microorganisms. *FEMS Microbiol Rev.* 2013;37(5):634-63. <https://doi.org/10.1111/1574-6976.12028>
- Berendsen RL, Vismans GyuK, Song Y, Jonge R, Burgman WP, Burmölle M, et al. Disease-induced assemblage of a plant-beneficial bacterial consortium. *ISME J.* 2018;12(6):1496-507. <https://doi.org/10.1038/s41396-018-0093-1>
- Gu Y, Wei Z, Wang X, Friman V, Huang J, Wang X, et al. Pathogen invasion indirectly changes the composition of soil microbiome via shifts in root exudation profile. *Biol Fert Soils.* 2016;52(7):997-1005. <https://doi.org/10.1007/s00374-016-1136-2>
- Raaijmakers JM, Weller DM. Exploiting genotypic diversity of 2,4-diacetylphloroglucinol-producing *Pseudomonas* spp.: characterization of superior root-colonizing *P. fluorescens* strain Q8r1-96. *Appl Environ Microbiol.* 2001;67(6):2545-54. <https://doi.org/10.1128/AEM.67.6.2545-2554.2001>
- Suzuki M. Biosynthesis and secretion of mugineic acid family phytosiderophores in zinc-deficient barley. *Plant J.* 2006;48(1):85-97. <https://doi.org/10.1111/j.1365-3113X.2006.02853.x>
- Kumar A, Bhatti SK, Aggarwal A. Biodiversity of endophytic mycorrhiza in some ornamental flowering plants of Solan, Himachal Pradesh. *Biol Forum.* 2012;4(2):45-51.
- Heijden MGVD, Martin FM, Selosse MA, Sanders IR. Mycorrhizal ecology and evolution: the past, the present and the future. *New Phytol.* 2015;205(4):1406-23. <https://doi.org/10.1111/nph.13288>
- Fitzpatrick CR, Copeland J, Wang PW, Johnson MTJ. Assembly and ecological function of the root microbiome across angiosperm plant species. *Proc Natl Acad Sci USA.* 2018;115(6):1157-65. <https://doi.org/10.1073/pnas.1717617115>
- Auge RM, Schekel KA, Wample RL. Greater leaf conductance of well-watered VA mycorrhizal rose plants is not related to phosphorus nutrition. *New Phytol.* 1986;103(1):107-16. <https://doi.org/10.1111/j.1469-8137.1986.tb00600.x>
- Green CD, Stodola A, Auge RM. Transpiration of detached leaves from mycorrhizal and non-mycorrhizal cowpea and rose plants given varying abscisic acid, pH, calcium and phosphorus. *Mycorrhiza.* 1998;8(2):93-9. <https://doi.org/10.1007/s005720050218>
- Pinior A, Grunewaldt SG, Von AH, Strasser RJ. Mycorrhizal impact on drought stress tolerance of rose plants probed by chlorophyll a fluorescence, proline content and visual scoring. *Mycorrhiza.* 2005;15(8):596-605. <https://doi.org/10.1007/s00572-005-0001-1>
- Aboul-Nasr A. Effects of vesicular-arbuscular mycorrhiza on *Tagetes erecta* and *Zinnia elegans*. *Mycorrhiza.* 1996;6(1):61-4. <https://doi.org/10.1007/s005720050107>
- Ganadevi G, Haripriya K. Studies on screening efficient VAM fungi for chrysanthemum. *South Indian Hortic.* 1999;47(1-6):325-6.
- Haripriya K, Sriramachandrasekharan MV. Effect of VAM inoculation on growth and yield of chrysanthemum. *J Ecobiol.* 2002;14(1):39-42.
- Compant S, Mitter B, Colli-Mull JG, Gangl H, Sessitsch A. Endophytes of grapevine flowers, berries and seeds: identification of cultivable bacteria, comparison with other plant parts and visualization of niches of colonization. *Microb Ecol.* 2011;62(1):188-97. <https://doi.org/10.1007/s00248-011-9883-y>
- Finkel OM, Burch Adrien Y, Lindow SE, Post AF, Belkin S. Geographical location determines the population structure in phyllosphere microbial communities of a salt-excreting desert tree. *Appl Environ Microbiol.* 2011;77(21):7647-55. <https://doi.org/10.1128/AEM.05565-11>
- Heydenreich B, Bellinghausen I, König B, Becker WM, Grabbe S, Petersen A, et al. Gram-positive bacteria on grass pollen exhibit adjuvant activity inducing inflammatory T cell responses. *Clin Exp Allergy.* 2012;42(1):76-84. <https://doi.org/10.1111/j.1365-2222.2011.03888.x>
- Pusey PL, Stockwell VO, Mazzola M. Epiphytic bacteria and yeasts on apple blossoms and their potential as antagonists of *Erwinia amylovora*. *Phytopathology.* 2009;99(5):571-81. <https://doi.org/10.1094/PHYTO-99-5-0571>
- Groenewald M, Robert V, Smith MT. Five novel *Wickerhamomyces*- and *Metschnikowia*-related yeast species, *Wickerhamomyces chaumierensis* sp. nov., *Candida pseudofloscolorum* sp. nov., *Candida danieliae* sp. nov., *Candida robnettiae* sp. nov. and *Candida eppingiae* sp. nov., isolated from plants. *Int J Syst Evol Microbiol.* 2011;61(8):2015-22. <https://doi.org/10.1099/ijs.0.026062-0>
- Junker RR, Loewel C, Gross R, Dötterl S, Keller A, Blüthgen N. Composition of epiphytic bacterial communities differs on petals and leaves. *Plant Biol.* 2011;13(6):918-24. <https://doi.org/10.1111/j.1438-8677.2011.00454.x>
- Pozo MI, Lachance MA, Herrera CM. Nectar yeasts of two southern Spanish plants: the roles of immigration and physiological traits in community assembly. *FEMS Microbiol Ecol.* 2012;80(2):281-93. <https://doi.org/10.1111/j.1574-6941.2011.01286.x>
- Bouffaud ML, Poirier MA, Muller D, Moenne-Loccoz Y. Root microbiome relates to plant host evolution in maize and other Poaceae. *Environ Microbiol.* 2014;16(9):2804-14. <https://doi.org/10.1111/1462-2920.12442>
- Jangid K, Williams MA, Franzluebbers AJ, Schmidt TM, Coleman DC, Whitman WB. Land-use history has a stronger impact on soil microbial community composition than aboveground vegetation and soil properties. *Soil Biol Biochem.* 2011;43(11):2184-93. <https://doi.org/10.1016/j.soilbio.2011.06.022>
- Besmer YL, Koide RT. Effect of mycorrhizal colonization and phosphorus on ethylene production by snapdragon (*Antirrhinum majus* L.) flowers. *Mycorrhiza.* 1999;9(3):161-6. <https://doi.org/10.1007/s005720050301>
- Sohn BK, Kim KY, Chung SJ, Kim WS, Park SM, Kang JG, et al. Effect of the different timing of AMF inoculation on plant growth and flower quality of chrysanthemum. *Sci Hortic.* 2003;98(2):173-83. [https://doi.org/10.1016/S0304-4238\(02\)00210-8](https://doi.org/10.1016/S0304-4238(02)00210-8)
- Bednarek P. Chemical warfare or modulators of defence responses – the function of secondary metabolites in plant immunity. *Curr Opin Plant Biol.* 2012;15(4):407-14. <https://doi.org/10.1016/j.pbi.2012.03.002>
- Sugiyama Y, Ueda Y, Zushi T, Takase H, Yazaki K. Changes in the bacterial community of soybean rhizospheres during growth in the field. *PLoS One.* 2014;9(7):e100709. <https://doi.org/10.1371/journal.pone.0100709>
- Harmsen G, Jager G. Determination of the quantity of carbon and nitrogen in the rhizosphere of young plants. *Nature.* 1962;195(4846):1119-20. <https://doi.org/10.1038/1951119a0>
- Xie MM, Wu QS. Arbuscular mycorrhizal fungi regulate flowering of *Hyacinth orientalis* L. Anna marie. *Emirates J Food Agric.* 2018;30(2):144-9. <https://doi.org/10.9755/ejfa.2018.v30.i2.1614>
- Fierer N. Embracing the unknown: disentangling the complexities of the soil microbiome. *Nat Rev Microbiol.* 2017;15(9):579-90. <https://doi.org/10.1038/nrmicro.2017.87>
- Hardoim PR, van Overbeek LS, Berg G, Pirttilä AM, Compant S, Campisano A, et al. The hidden world within plants: ecological and evolutionary considerations for defining functioning of microbial endophytes. *Microbiol Mol Biol Rev.* 2015;79(3):293-320. <https://doi.org/10.1128/MMBR.00050-14>
- Navarrete AA, Cannavan FS, Taketani RG, Tsai SM. A molecular survey of the diversity of microbial communities in different Amazonian agricultural model systems. *Diversity.* 2010;2(5):787-809. <https://doi.org/10.3390/d2050787>
- Hartmann A, Schmid M, van Tuinen D, Berg G. Plant-driven selection of microbes. *Plant Soil.* 2009;321(1-2):235-75. <https://doi.org/10.1111/j.1365-2222.2011.03888.x>

- doi.org/10.1007/s11104-008-9814-y
34. Tripathi BM, Moreenyane I, Sherman C, Lee YK, Adams JM, Steinberger Y. Trends in taxonomic and functional composition of soil microbiome along a precipitation gradient in Israel. *Microb Ecol*. 2017;74(1):168-76. <https://doi.org/10.1007/s00248-017-0931-0>
 35. Kumar P, Thakur S, Dhingra GK, Singh A, Pal M, Harshvardhan K, et al. Inoculation of siderophore producing rhizobacteria and their consortium for growth enhancement of wheat plant. *Biocatal Agric Biotechnol*. 2018;15:264-9. <https://doi.org/10.1016/j.bcab.2018.06.019>
 36. De Vries FT, Griffiths RI, Knight CG, Nicolitch O, Williams A. Harnessing rhizosphere microbiomes for drought-resilient crop production. *Science*. 2020;368(6488):270-4. <https://doi.org/10.1126/science.aaz5192>
 37. de Bello F, Lavorel S, Díaz S, Harrington R, Cornelissen JHC, Bardgett RD, et al. Towards an assessment of multiple ecosystem processes and services via functional traits. *Biodivers Conserv*. 2010;19(4):2873-93. <https://doi.org/10.1007/s10531-010-9850-9>
 38. Hardoim PR, van Overbeek LS, Van EJD. Properties of bacterial endophytes and their proposed role in plant growth. *Trends Microbiol*. 2008;16(9):463-71. <https://doi.org/10.1016/j.tim.2008.07.008>
 39. Spaepen S, Dobbelaere S, Croonenborghs A, Vanderleyden J. Effects of *Azospirillum brasilense* indole-3-acetic acid production on inoculated wheat plants. *Plant Soil*. 2008;312(1-2):15-23. <https://doi.org/10.1007/s11104-008-9560-1>
 40. Oblisami G. The inter-relationships between ornamental plants and soil microorganisms. 1. The rhizosphere microflora of marigold. *South Indian Hortic*. 2008;21(2):59-62.
 41. Oblisami G, Onkarayya H. The inter-relationships between ornamental plants and soil microorganisms. 2. The rhizosphere microflora of balsam and English daisy and enzyme activities. *South Indian Hortic*. 1973;21(3):83-8.
 42. Bhattacharya P, Mishra UC. Biofertilizers for flowers and ornamental plants. In: Singh T, editor. A book on biofertilizer to extension worker. UP: National Biofertilizer Development Centre; 1995. p. 102.
 43. Marulanda A, Azcón R, Chaumont F, Lozano JMR, Aroca R. Regulation of plasma membrane aquaporins by inoculation with a *Bacillus megaterium* strain in maize (*Zea mays* L.) plants under unstressed and salt-stressed conditions. *Planta*. 2010;232(2):533-43. <https://doi.org/10.1007/s00425-010-1196-8>
 44. Rillig MC, Mummey DL. Mycorrhizas and soil structure. *New Phytol*. 2006;171(1):41-53. <https://doi.org/10.1111/j.1469-8137.2006.01750.x>
 45. Flores CA, Estrada LAA, Olalde PV. Yield and quality enhancement of marigold flowers by inoculation with *Bacillus subtilis* and *Glomus fasciculatum*. *J Sustain Agric Environ*. 2007;31(1):21-31. https://doi.org/10.1300/J064v31n01_04
 46. Prasad K, Aggarwal A, Yadav K, Tanwar A. Impact of different levels of superphosphate using arbuscular mycorrhizal fungi and *Pseudomonas fluorescens* on *Chrysanthemum indicum* L. *J Soil Sci Plant Nutr*. 2012;12(3):451-62. <https://doi.org/10.4067/S0718-95162012005000007>
 47. Saini I, Aggarwal A, Kaushik P. Inoculation with mycorrhizal fungi and other microbes to improve the morpho-physiological and floral traits of *Gazania rigens* (L.) Gaertn. *Agriculture*. 2019;9(3):51. <https://doi.org/10.3390/agriculture9030051>
 48. Gore ME, Altin N. Growth promoting of some ornamental plants by root treatment with specific fluorescent pseudomonads. *J Biol Sci*. 2006;6(3):610-15. <https://doi.org/10.3923/jbs.2006.610.615>
 49. Syamal MM, Dixit SK, Kumar S. Effect of biofertilizers on growth and yield in marigold. *J Ornamental Hortic*. 2006;9(4):304-5.
 50. Singh AK. Effect of FYM, *Azotobacter* and nitrogen on leaf nutrient composition, growth, flowering and yield in rose. *Indian J Hortic*. 2006;63(1):62-5.
 51. Dubey RK, Misra RL. Study of chemical and biofertilizers on gladiolus. *Progressive Hortic*. 2006;38(2):125-8.
 52. Yadav BS, Gupta AK, Singh S. Studies on the effect of nitrogen, plant spacing and bio-fertilizers on growth parameters in tuberose cv. Double. *Haryana J Hortic Sci*. 2005;34(1-2):78-80.
 53. Eid AR, Awad MN, Hamouda HA. Evaluate effectiveness of bio and mineral fertilization on the growth parameters and marketable cut flowers of *Matthiola incana* L. *Am Eurasian J Agric Environ Sci*. 2009;5(4):509-18.
 54. Preethi TL, Pappiah CM, Anbu S. Studies on the effect of *Azospirillum* sp., nitrogen and ascorbic acid on the growth and flowering of Edward rose (*Rosa bourboniana* Desp.). *South Indian Hortic*. 1999;47(1-6):106-10.
 55. Bhalla R, Sandeep D, Dhiman SR, Ritu J. Effect of biofertilizers and biostimulants on growth and flowering in standard carnation (*Dianthus caryophyllus* Linn.). *J Ornament Hortic*. 2006;9(4):282-4.
 56. Hoda EEM, Mona S. Effect of bio and chemical fertilizers on growth and flowering of *Petunia hybrida* plants. *Am J Plant Physiol*. 2014;9(2):68-77. <https://doi.org/10.3923/ajpp.2014.68.77>
 57. Mosse B. Advances in study of vesicular arbuscular mycorrhiza. *Ann Rev Phytopathol*. 1981;19:171-98.
 58. Perner H, Schwarz D, Bruns C, Mäder P, George E. Effect of arbuscular mycorrhizal colonization and two levels of compost supply on nutrient uptake and flowering of *Pelargonium* plants. *Mycorrhiza*. 2007;17(5):469-74. <https://doi.org/10.1007/s00572-007-0116-7>
 59. Meir D, Pivonia S, Levita R, Dori I, Ganot L, Meir S, et al. Application of mycorrhizae to ornamental horticultural crops: *lisianthus* (*Eustoma gradiflorum*) as a test case. *Spanish J Agric Res*. 2010;8(S1):5-10. <https://doi.org/10.5424/sjar/201008S1-1221>
 60. Scagel CF. Inoculation with vesicular-arbuscular mycorrhizal fungi and rhizobacteria alters nutrient allocation and flowering of harlequin flower. *HortTechnology*. 2004;14(1):39-48. <https://doi.org/10.21273/HORTTECH.14.1.0039>
 61. Saini I, Yadav K, Aggarwal A. Effect of bioinoculants on morphological and biochemical parameters of *Zinnia elegans* Jacq. *J Appl Hortic*. 2017;19(3):167-72. <https://doi.org/10.37855/jah.2017.v19i02.31>
 62. Janowska B, Andrzejak R. Effect of mycorrhizal inoculation on development and flowering of *Tagetes patula* L. 'YellowBoy' and *Salvia splendens* Buc'hoz ex Et. 'Saluti Red'. *Acta Agrobot*. 2017;70(1):1703. <https://doi.org/10.5586/aa.1703>
 63. Gaur A, Adholeya A. Diverse response of five ornamental plant species to mixed indigenous and single isolate arbuscular-mycorrhizal inocula in marginal soil amended with organic matter. *J Plant Nutr*. 2005;28(4):707-23. <https://doi.org/10.1081/PLN-200052647>
 64. Asrar AA, Abdel-Fattah GM, Elhindi KM. Improving growth, flower yield and water relations of snapdragon (*Antirrhinum majus* L.) plants grown under well-watered and water-stress conditions using arbuscular mycorrhizal fungi. *Photosynthetica*. 2012;50(2):305-16. <https://doi.org/10.1007/s11099-012-0024-8>
 65. Lazzara S, Militello M, Carrubba A, Napoli E, Saia S. Arbuscular mycorrhizal fungi altered the hypericin, pseudohypericin and hyperforin content in flowers of *Hypericum perforatum* grown under contrasting P availability in a highly organic substrate. *Mycorrhiza*. 2017;27(4):1-10. <https://doi.org/10.1007/s00572-016-0756-6>
 66. Nowak J. Effects of arbuscular mycorrhizal fungi and organic fertilization on growth, flowering, nutrient uptake, photosynthesis and transpiration of geranium (*Pelargonium hortorum* L.H. Bailey 'Tango Orange'). *Symbiosis*. 2004;37(1-4):259-66.
 67. Attia M, Eid RA. Effect of inoculation timing with arbuscular mycorrhizal fungi on growth and flowering of micropropagated

- Chrysanthemum morifolium*. Arab Univ J Agric Sci. 2005;13(3):677-88. <https://doi.org/10.21608/ajs.2005.15269>
68. Vaingankar JD, Rodrigues BF. Effect of arbuscular mycorrhizal inoculation on growth and flowering in *Crossandra infundibuliformis* (L.) Nees. J Plant Nutr. 2015;38(9):1478-88. <https://doi.org/10.1080/01904167.2014.957398>
 69. Tarafdar JC, Gharu A. Mobilization of organic and poorly soluble phosphates by *Chaetomium globosum*. Appl Soil Ecol. 2006;32(3):273-83. <https://doi.org/10.1016/j.apsoil.2005.08.005>
 70. Chandra S, Srivastava N, Narayan R, Sharma N. Efficacy of microbial inoculants on reducing the phosphatic fertilizer input in chrysanthemum. Indian J Hortic. 2009;66(1):83-7.
 71. Lin CS, Lin SY, Guang WC. Microbial inoculation effect on the growth and phosphorus concentration of *Eustoma grandiflorum*. J Agric Res China. 2001;50(4):66-73.
 72. Scagel CF. Inoculation with arbuscular mycorrhizal fungi alters nutrient allocation and flowering of *Freesia × hybrida*. J Environ Hortic. 2003;21(4):196-205.
 73. Garmendia I, Mangas VJ. Application of arbuscular mycorrhizal fungi on the production of cut flower roses under commercial-like conditions. Spanish J Agric Res. 2012;10(1):166-74. <https://doi.org/10.5424/sjar/2012101-156-11>
 74. Dubsky M, Sramek F. Inoculation of cyclamen (*Cyclamen persicum*) and poinsettia (*Euphorbia pulcherrima*) with arbuscular mycorrhizal fungi and *Trichoderma harzianum*. Rostlinna Vyroba. 2002;48(2):63-8. <https://doi.org/10.17221/4361-PSE>
 75. Adholeya A. Diverse response of five ornamental plant species to mixed indigenous and single isolate arbuscular mycorrhizal inocula in marginal soil amended with organic matter. J Plant Nutr. 2005;28(4):707-23. <https://doi.org/10.1081/PLN-200052647>
 76. Jin Z, Li J, Li Y. Interactive effects of arbuscular mycorrhizal fungi and copper stress on flowering phenology and reproduction of *Elsholtzia splendens*. PLoS One. 2015;10(12):e0145793. <https://doi.org/10.1371/journal.pone.0145793>
 77. Banla EM, Banito A, Sogbedji JM. Effects of arbuscular mycorrhizal fungi on the production of tomato in Togo. Int J Biol Chem Sci. 2015;9(3):1270-6. <https://doi.org/10.4314/ijbcs.v9i3.12>
 78. Soroa MR, Cortes SL, Hernandez A. Study of the effect of application of biofertilizers on some variables of growth and yield in *Gerbera jamesonii* cv. Bolus. Cultiv Trop. 2003;24(2):15-7.
 79. Schroth MN, Hildebrand DC. Influence of plant exudates on root-infecting fungi. Ann Rev Phytopathol. 1964;2:101-32. <https://doi.org/10.1146/annurev.py.02.090164.000533>
 80. Heath MC. A generalized concept of host-parasite specificity. J Phytopathol. 1981;71(12):1121-3. <https://doi.org/10.1094/Phyto-71-1121>
 81. Mazumdar N. Epidemiological factors in relation to development of *Alternaria* leaf blight of marigold and fungicidal control. Plant Disease Res. 2000;15(1):28-33.
 82. Agrios GN. Plant pathology. 5th ed. New York: Academic Press; 2005.
 83. Saroj A, Kumar A, Saeed ST, Samad A, Alam M. First report of *Tagetes erecta* damping off caused by *Ceratobasidium* sp. from India. Plant Dis. 2017;97(9):1251. <https://doi.org/10.1094/PDIS-02-13-0145-PDN>
 84. Trolinger JC, Strider DL. Botrytis diseases. In: Strider DL, editor. Diseases of floral crops. Vol. 1. New York: Praeger Special Studies; 1985. p. 17-101.
 85. Khatun S, Chakraborti J, Santra S, Banerjee S, Chatterjee S, Dutta S, et al. Black spot disease of rose (*Rosa centifolia*) – a new record from India. Int J Plant Sci. 2008;3(1):294.
 86. Shukla A, Thakur R. First report of septoria leaf spot on marigold (*Tagetes erecta* L.) from Himachal Pradesh, India. Int J Curr Microbiol Appl Sci. 2018;7(1):1744-8. <https://doi.org/10.20546/ijcmas.2018.701.211>
 87. Minuto A, Gullino ML, Garibaldi A. *Gerbera jamesonii*, *Osteospermum* sp. and *Argyranthemum frutescens*: new hosts of *Fusarium oxysporum* f. sp. chrysanthemi. J Phytopathol. 2007;155(6):373-6. <https://doi.org/10.1111/j.1439-0434.2007.01248.x>
 88. Troisi M, Gullino ML, Garibaldi A. *Gerbera jamesonii*, a new host of *Fusarium oxysporum* f. sp. tracheiphilum. J Phytopathol. 2010;158(1):8-14. <https://doi.org/10.1111/j.1439-0434.2009.01551.x>
 89. Padghan PR, Gade RM. Biomanagement of root rot complex of gerbera (*Gerbera jamesonii* Bolus). Ann Plant Prot Sci. 2006;14(1):134-8.
 90. Ortu G, Bertetti D, Gullino ML, Garibaldi A. New formae speciales of *Fusarium oxysporum* on ornamental plants. Acta Hortic. 2014;1044:89-92. <https://doi.org/10.17660/ActaHortic.2014.1044.9>
 91. Uroz S, Courty PE, Oger P. Plant symbionts are engineers of the plant-associated microbiome. Trends Plant Sci. 2019;24(10):905-16. <https://doi.org/10.1016/j.tplants.2019.06.008>
 92. Berg M, Koskella B. Nutrient- and dose-dependent microbiome-mediated protection against a plant pathogen. Curr Biol. 2018;28(15):2487-92.e3. <https://doi.org/10.1016/j.cub.2018.05.085>
 93. Hunter P. Plant microbiomes and sustainable agriculture. EMBO Rep. 2016;17(9):1-4. <https://doi.org/10.15252/embr.201643476>
 94. Ellis JG. Can plant microbiome studies lead to effective biocontrol of plant diseases? Mol Plant-Microbe Interact. 2017;30(3):190-3. <https://doi.org/10.1094/MPMI-12-16-0252-CR>
 95. Trivedi PM, Delgado-Baquerizo Trivedi C, Hamonts K, Anderson IC, Singh BK. Keystone microbial taxa regulate the invasion of a fungal pathogen in agroecosystems. Soil Biol Biochem. 2017;111:10-14. <https://doi.org/10.1016/j.soilbio.2017.03.013>
 96. Singh P, Vijay K. Biological control of *Fusarium* wilt of chrysanthemum with *Trichoderma* and botanicals. Int J Agric Technol. 2011;7(6):1603-13.
 97. Maya MA, Matsubara Y. Tolerance to *Fusarium* wilt and anthracnose diseases and changes of antioxidative activity in mycorrhizal cyclamen. Crop Prot. 2013;47:41-8. <https://doi.org/10.1016/j.cropro.2013.01.007>
 98. Hassan N, Elsharkawy MM, Shimizu M, Hyakumachi M. Control of root rot and wilt diseases of roselle under field conditions. Mycobiology. 2014;42(4):376-84. <https://doi.org/10.5941/MYCO.2014.42.4.376>
 99. David BV, Chandrasehar G, Selvam PN. *Pseudomonas fluorescens*: a plant-growth-promoting rhizobacterium (PGPR) with potential role in biocontrol of pests of crops. In: Sharma M, Kumar R, Singh P, editors. Crop improvement through microbial biotechnology. Amsterdam: Elsevier; 2018. p. 221-43. <https://doi.org/10.1016/B978-0-444-63987-5.00010-4>
 100. Zhang Z, Kong X, Jin D, Yu H, Zhu X, Su X, et al. *Euonymus japonicus* phyllosphere microbiome is significantly changed by powdery mildew. Arch Microbiol. 2019;201(8):1099-109. <https://doi.org/10.1007/s00203-019-01683-3>
 101. Kumar V, Chandel S. Management of rose powdery mildew (*Podosphaera pannosa*) through ecofriendly approaches. Indian Phytopathol. 2018;71(3):393-7. <https://doi.org/10.1007/s42360-018-0050-y>
 102. Janice FE, Haselhan C, Punja ZK. Evaluation of biological control agents for control of botrytis blight of geranium and powdery mildew of rose. Canadian J Plant Pathol. 2011;33(4):499-505. <https://doi.org/10.1080/07060661.2011.630758>
 103. Tyagi A, Raj H. Integration of soil solarization with bio-control agents for the management of stem rot of chrysanthemum. J Pharmacogn Phytochem. 2021;10(1):2468-71. <https://doi.org/10.22271/phyto.2021.v10.i1ai.13734>
 104. Chandel S, Sharma S. Botanicals, biofumigants and antagonists application in managing stem rot disease caused by *Rhizoctonia*

- solani* Kuhn in carnation. J Biopestic. 2014;7(1):3-10. <https://doi.org/10.57182/jbiopestic.7.1.3-10>
105. Vavre KB, Kakade DS, Sawne SS, Khaire PB. In vitro efficacy of bio-control agents against *Fusarium oxysporum* f. sp. *gladioli*. J Pharmacogn Phytochem. 2021;10(1):590-3.
 106. Atakan A, Ozkaya HO. Induced resistance to Fusarium wilt in carnation with mixture of mycorrhizal fungi. Fresen Environ Bull. 2021;30(4A):4217-27.
 107. Chandel S. Organic amendment, biocontrol agents and soil solarization practice in management of fusarium wilt of carnation caused by *Fusarium oxysporum* Schlecht. f. sp. *dianthi* (Prill. and Del.) Snyder. Indian J Plant Prot. 2015;8(1):130-3. <https://doi.org/10.15740/HAS/IJPP/8.1/130-133>
 108. Mahalakshmi P, Ahiladevi P. Ecofriendly management of fusarium wilt of carnation (*Dianthus caryophyllus* L.) by application of organic amendments. Adv Appl Sci Res. 2020;12(1):7-11. <https://doi.org/10.5958/2349-2104.2020.00002.9>
 109. Rajendran L, Raja P, Vellaichamy J, Pushparaj V, Selvaraj NK. *Pseudomonas fluorescens* and *Trichoderma viride* enriched bioconsortium for the management of Fusarium wilt in carnation and gerbera under protected cultivation. Indian Phytopathol. 2014;67(1):77-81.
 110. Salma Z, Sindhu SS, Ahlawat VP. Suppression of Fusarium wilt disease in gladiolus by using rhizobacterial strains. J Crop Weed. 2014;10(2):466-71.
 111. Kumar S, Tomar KS, Shakywar RC, Mahesh P. Integrated management of fusarium wilt of gladiolus (*Gladiolus hybridus* Hort.) in Arunachal Pradesh conditions. J Ornament Hort. 2013;16(1):52-6.
 112. Shanmugam V, Kanoujia N, Singh M, Singh S, Prasad R. Biocontrol of vascular wilt and corm rot of gladiolus caused by *Fusarium oxysporum* f. sp. *gladioli* using plant growth promoting rhizobacterial mixture. Crop Prot. 2011;30(7):807-13. <https://doi.org/10.1016/j.cropro.2011.02.033>
 113. Beale R, Pitt D. Biological and integrated control of Fusarium basal rot of Narcissus using *Minimedusa polyspora* and other micro-organisms. Plant Pathol. 1990;39(3):477-88. <https://doi.org/10.1111/j.1365-3059.1990.tb02524.x>
 114. Tatagiba JS, Maffia LA, Barreto RW, Alfenas AC, Sutton JC. Biological control of *Botrytis cinerea* in residues and flowers of Rose (*Rosa hybrida*). Phytoparasitica. 1998;26(1):8-19. <https://doi.org/10.1007/BF02981261>
 115. Kohl J, Postma J, Nicot P, Ruocco M, Blum B. Stepwise screening of microorganisms for commercial use in biological control of plant pathogenic fungi and bacteria. Biol Control. 2011;57(1):1-12. <https://doi.org/10.1016/j.biocontrol.2010.12.004>
 116. Romina D, Massa N, Amalero EG, Gostino GD, Ampò SS, Berta G, et al. Preliminary results on the evaluation of the effects of elicitors of plant resistance on chrysanthemum yellows phytoplasma infection. Bull Insectol. 2012;60(2):317-18.
 117. Rout MK, Mishra B. Studies on collar rot disease of marigold and its management. J Mycopathol Res. 2008;46(2):283-4.
 118. Widmer TL, Brousseau SSJ, Kosta K, Ghosh S, Schweigkofler W, Sharma SK. Remediation of *Phytophthora ramorum*-infested soil with *Trichoderma asperellum* isolate under ornamental nursery conditions. Biol Control. 2018;118:67-73. <https://doi.org/10.1016/j.biocontrol.2017.12.007>
 119. Elhady A, Adss S, Hallmann J, Heuer H. Rhizosphere microbiomes modulated by pre-crops assisted plants in defense against plant-parasitic nematodes. Front Microbiol. 2018;9:1133. <https://doi.org/10.3389/fmicb.2018.01133>
 120. Ritpitakphong U, Falquet L, Vimoltust A, Berger A, Métraux JP, Haridon FL. The microbiome of the leaf surface of *Arabidopsis* protects against a fungal pathogen. New Phytol. 2016;210(3):1033-43. <https://doi.org/10.1111/nph.13808>
 121. Naylor D, de Graaf S, Purdom E, Derr DC. Drought and host selection influence bacterial community dynamics in the grass root microbiome. ISME J. 2017;11(12):2691-704. <https://doi.org/10.1038/ismej.2017.118>
 122. Niu B, Paulson JN, Zheng X, Kolter R. Simplified and representative bacterial community of maize roots. Proc Natl Acad Sci USA. 2017;114(12):E2450-9. <https://doi.org/10.1073/pnas.1616148114>
 123. Hassan S, Mathesius U. The role of flavonoids in root-rhizosphere signalling: opportunities and challenges for improving plant-microbe interactions. J Exp Bot. 2012;63(9):3429-44. <https://doi.org/10.1093/jxb/err430>
 124. Liu H, Khan MY, Carvalhais LC, Baquerizo MD, Yan L, Crawford M, et al. Soil amendments with ethylene precursor alleviate negative impacts of salinity on soil microbial properties and productivity. Sci Rep. 2019;9(1):6892. <https://doi.org/10.1038/s41598-019-43305-4>
 125. Xu L, Derr DC. Causes and consequences of a conserved bacterial root microbiome response to drought stress. Curr Opin Microbiol. 2019;49:1-6. <https://doi.org/10.1016/j.mib.2019.07.003>
 126. Duc NH, Csintalan Z, Posta K. Arbuscular mycorrhizal fungi mitigate negative effects of combined drought and heat stress on tomato plants. Plant Physiol Biochem. 2018;132:297-307. <https://doi.org/10.1016/j.plaphy.2018.09.011>
 127. Maya MA, Ito M, Matsubara Y. Tolerance to heat stress and anthracnose in mycorrhizal *Cyclamen*. Acta Hort. 2014;(1025):371-8. <https://doi.org/10.17660/ActaHortic.2014.1025.21>
 128. Glick BR. Bacteria with ACC deaminase can promote plant growth and help to feed the world. Microbiol Res. 2014;169(1):30-9. <https://doi.org/10.1016/j.micres.2013.09.009>
 129. Mosqueda COM, Glick BR, Santoyo G. ACC deaminase in plant growth-promoting bacteria (PGPB): an efficient mechanism to counter salt stress in crops. Microbiol Res. 2020;235:126439. <https://doi.org/10.1016/j.micres.2020.126439>
 130. Arny D, Lindow S, Upper C. Frost sensitivity of *Zea mays* increased by application of *Pseudomonas syringae*. Nature. 1976;262(5566):282-4. <https://doi.org/10.1038/262282a0>
 131. Failor K, Schmale DG, Vinatzer BA, Monteil CL. Ice nucleation active bacteria in precipitation are genetically diverse and nucleate ice by employing different mechanisms. ISME J. 2017;11(12):2740-53. <https://doi.org/10.1038/ismej.2017.124>
 132. Smirnov N. The role of active oxygen in the response of plants to water deficit and desiccation. New Phytol. 1993;125(1):27-58. <https://doi.org/10.1111/j.1469-8137.1993.tb03863.x>
 133. Rouphael Y, Cardarelli M, Schwarz D, Franken P, Colla G. Effects of drought on nutrient uptake and assimilation in vegetable crops. In: Aroca R, editor. Plant responses to drought stress. Berlin: Springer; 2012. p. 171-95. https://doi.org/10.1007/978-3-642-32653-0_7
 134. Sangiorgio D, Cellini A, Donati I, Pastore C, Onofrietti C, Spinelli F. Facing climate change: application of microbial biostimulants to mitigate stress in horticultural crops. Agronomy. 2020;10(6):794. <https://doi.org/10.3390/agronomy10060794>
 135. Santoyo G, Gamalero E, Glick BR. Mycorrhizal-bacterial amelioration of plant abiotic and biotic stress. Front Sustain Food Syst. 2021;5:139. <https://doi.org/10.3389/fsufs.2021.672881>
 136. Wang Y, Wang M, Li Y, Wu A, Huang J. Effects of arbuscular mycorrhizal fungi on growth and nitrogen uptake of *Chrysanthemum morifolium* under salt stress. PLoS One. 2018;13(4):e0196408. <https://doi.org/10.1371/journal.pone.0196408>
 137. Gong M, Tang M, Chen H, Zhang Q, Feng X. Effects of two *Glomus* species on the growth and physiological performance of *Sophora davidii* seedlings under water stress. New For. 2013;44(3):399-408. <https://doi.org/10.1007/s11056-012-9349-1>
 138. Fagbola O, Osonubi O, Mulongoy K, Odunfa S. Effects of drought stress and arbuscular mycorrhiza on the growth of *Gliricidia sepium* (Jacq.) Walp. *Leucaena leucocephala* (Lam). De wit. in simulated

- eroded soil conditions. *Mycorrhiza*. 2001;11(5):215-23. <https://doi.org/10.1007/s005720100114>
139. Scheilenbaum L, Schuepp H, Wiemken A, Boller T. Effect of drought, transgenic expression of a fructan synthesizing enzyme and of mycorrhizal symbiosis on growth and soluble carbohydrate pools in tobacco plants. *New Phytol*. 1999;142(1):67-77. <https://doi.org/10.1046/j.1469-8137.1999.00376.x>
 140. Asrar AWA, Elhindi KM. Alleviation of drought stress of marigold (*Tagetes erecta*) plants by using arbuscular mycorrhizal fungi. *Saudi J Biol Sci*. 2011;18(1):93-8. <https://doi.org/10.1016/j.sjbs.2010.06.007>
 141. Al-Karaki GN, Clark RB. Growth, mineral acquisition and water use by mycorrhizal wheat grown under water stress. *J Plant Nutr*. 1998;21(2):263-76. <https://doi.org/10.1080/01904169809365401>
 142. Jha CK, Saraf M. Plant growth promoting rhizobacteria (PGPR): a review. *J Agric Res Dev*. 2015;5(1):108-19.
 143. Nordstedt NP, Chapin LJ, Taylor CG, Jones ML. Identification of *Pseudomonas* spp. that increases ornamental crop quality during abiotic stress. *Front Plant Sci*. 2020;10:1754. <https://doi.org/10.3389/fpls.2019.01754>
 144. Lin Y, Jones ML. Evaluating the growth-promoting effects of microbial biostimulants on greenhouse floriculture crops. *HortScience*. 2021;57(1):97-109. <https://doi.org/10.21273/HORTSCI16149-21>
 145. Zhou C, Zhu L, Xie Y, Li F, Xiao X, Ma Z, et al. *Bacillus licheniformis* SA03 confers increased saline-alkaline tolerance in Chrysanthemum plants by induction of abscisic acid accumulation. *Front Plant Sci*. 2017;8:1143. <https://doi.org/10.3389/fpls.2017.01143>
 146. Prisa D. Role of microorganisms in horticulture to improve plant quality. *Karbala Int J Modern Sci*. 2023;9(1):34-42. <https://doi.org/10.33640/2405-609X.3284>
 147. Kumbhar I, Patil CP, Kulkarni BS, Shiragur M, Shirol AM. Efficacy of *Entrophospora* sp. (VA mycorrhiza) on salt tolerance and flower yield and quality of Chrysanthemum var. Marigold (*Dendranthema grandiflora* Tzvelev.). *Int J Curr Microbiol Appl Sci*. 2017;6(10):4769-77. <https://doi.org/10.20546/ijcmas.2017.610.443>
 148. Yuan Y, Brune C, van Kleunen M, Li J, Jin Z. Salinity-induced changes in the rhizosphere microbiome improve salt tolerance of *Hibiscus hamabo*. *Plant Soil*. 2019;443(1):525-37. <https://doi.org/10.1007/s11104-019-04258-9>
 149. Samad A, Antonielli L, Sessitsch A, Compant S, Trognitz F. Comparative genome analysis of the vineyard weed endophyte *Pseudomonas viridiflava* CDRTc14 showing selective herbicidal activity. *Sci Rep*. 2017;7(1):17336. <https://doi.org/10.1038/s41598-017-16495-y>
 150. Chowdhury SP, Hartmann A, Gao XW, Rainer B. Biocontrol mechanism by root-associated *Bacillus amyloliquefaciens* FZB42 - a review. *Front Microbiol*. 2015;6:780. <https://doi.org/10.3389/fmicb.2015.00780>
 151. Raaijmakers JM, Paulitz TC, Steinberg C, Alabouvette C, Moënne-Loccoz Y. The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial microorganisms. *Plant Soil*. 2009;321(1-2):341-61. <https://doi.org/10.1007/s11104-008-9568-6>
 152. Ardanov P, Lyastchenko S, Karpinen K, Häggman H, Kozyrovska N, Pirttilä AM. Effects of *Methylobacterium* sp. on emergence, yield and disease prevalence in three cultivars of potato (*Solanum tuberosum* L.) were associated with the shift in endophytic microbial community. *Plant Soil*. 2016;405(1):299-310. <https://doi.org/10.1007/s11104-015-2500-y>
 153. Rodríguez-Caballero G, Caravaca F, Fernández-González AJ, Alguacil MM, Fernández-López M, Roldán A. Arbuscular mycorrhizal fungi inoculation mediated changes in rhizosphere bacterial community structure while promoting revegetation in a semiarid ecosystem. *Sci Total Environ*. 2017;584-5:838-48. <https://doi.org/10.1016/j.scitotenv.2017.01.128>
 154. Schmidt R, Köberl M, Mostafa A, Ramadan EM, Monschein M, Jensen KB, et al. Effects of bacterial inoculants on the indigenous microbiome and secondary metabolites of chamomile plants. *Front Microbiol*. 2014;5:1-11. <https://doi.org/10.3389/fmicb.2014.00064>
 155. Moghadam AR, Shoor M. Effect of different organic and inorganic fertilizers on growth and flowering of African violet (*Saintpaulia ionantha* Wendl.). *Int J Agric Crop Sci*. 2013;5(17):1930-6.
 156. Khan N, Bano A, Jameel M, Zahid I. Effect of plant growth promoting rhizobacteria (PGPR) on growth, physiology and yield of wheat under drought. *J Basic Appl Sci*. 2009;5(4):112-17.
 157. Yasin M, Ahmed N. Response of tuberose (*Polianthes tuberosa* Linn.) to combined application of bio-organo-mineral fertilizers. *J Plant Nutr*. 2016;39(12):1748-59.
 158. Sant S, Singh SK, Khan M. Effect of different phosphorus levels and PSB on growth, flowering and bulb production of tuberose (*Polianthes tuberosa* L.) cv. Double. *J Ornament Hort*. 2010;13(1):76-8.
 159. Negi A, Raj S. Effect of inorganic fertilizers and biofertilizers on growth, flowering and yield of *Gaillardia pulchella* Foug. var. 'Local Red'. *Int J Pharm Sci Rev Res*. 2013;23(2):35-9.

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc
See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.