



REVIEW ARTICLE

Organic compounds as biostimulants: Cow derivatives, botanicals and biofertilizers enhancing seed germination, vigour and yield in paddy

M Subhiksha¹, N Punithavathi^{2*}, A Sabir Ahamed³, S Elamathi⁴ & S Geethanjali⁵

¹Department of Seed Science and Technology, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

²Department of Seed Science and Technology, Krishi Vigyan Kendra, Tamil Nadu Agricultural University, Sirugamani, Trichy 639 115, Tamil Nadu, India

³Department of Vegetable Science, Horticultural College and Research Institute for Women, Trichy 620 027, Tamil Nadu, India

⁴Department of Agronomy, Tamil Nadu Rice Research Institute, Aduthurai 612 101, Tamil Nadu, India

⁵Department of Crop Physiology and Bio Chemistry, Anbil Dharmalingam Agricultural College and Research Institute, Trichy 620 027, Tamil Nadu, India

*Correspondence email - punithavathi.n@tnau.ac.in

Received: 23 March 2025; Accepted: 21 June 2025; Available online: Version 1.0: 19 July 2025

Cite this article: Subhiksha M, Punithavathi N, Sabir AA, Elamathi S, Geethanjali S. Organic compounds as biostimulants: Cow derivatives, botanicals and biofertilizers enhancing seed germination, vigour and yield in paddy. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.8468>

Abstract

Organic agriculture employs various natural techniques to cultivate crops, emphasizing crop rotation, composting and biological pest control. In modern intensive agriculture, chemical fertilizer plays a major role. Continuous application of these chemicals leads to the accumulation of residues, resulting in soil hardening and degradation of soil structure. Although chemically treated seeds and inorganic inputs can enhance crop yield initially, their excessive use may negatively affect productivity and result in the accumulation of toxins, rendering the produce unfit for human consumption. Organic farming addresses many of the challenges faced by modern agriculture and food production. The idea behind organic farming is to use affordable, easily accessible inputs. Organic farmers pay significantly more for labour and feed also spend less on inputs. Organic farming utilizes farmyard manure, compost, green manures, farm waste, vermicompost and crop residues that increases the crop yield and restore soil fertility. In this paper, the reasons for using organic compounds are reviewed and their roles are analysed, particularly focusing on their effects as seed treatments and foliar applications on seed germination, growth and yield in paddy. In future, new research pathways emerge, offering the chance to improve organic practices and ensure healthy environment.

Keywords: foliar application; germination; organic compounds; organic farming; seed treatment

Introduction

Rice serves as a staple food for more than half of the global population and is cultivated in more than 100 countries, with Asia accounting for 90 % of the total production (1). The Indian subcontinent consumes more than 40 % of its calories from rice (2). According to USDA, global rice output and yield are projected to increase by 3.8, 7.2 and 11.3 % by 2025-2026. China produces the most rice among the top ten rice-producing nations (3). In the year 2023-24, the area under rice cultivation in India was 47.82 million hectares, with a total production of 137.83 million tonnes and a productivity of 2882 kilograms per hectare (Department of Agriculture and Farmers Welfare). In India, West Bengal is the top rice-producing state, with an acreage of 5.80 million ha and the highest production of 15.5 million tons, yielding a productivity of 2.79 million tons (4). In conventional farming, the farming community have identified major issues such as quality seed insecurity, declining soil health, productivity stagnation and environmental risks. The use of easily accessible organic materials, such as FYM, compost, biogas slurry, etc., led to a significant increase in interest in

organic agriculture (Fig. 1) (5). The basic principles of organic agriculture are the ecology, fairness and environment (6-8). One of the main limitations of organic farming is lower yield compared to conventional farming. To meet the growing food demand of the increasing population, more land must be brought under organic cultivation to achieve comparable yields (9). The incorporation of augmented sludge, biochar in combination with organic fertilizers, biofertilizers, organo-minerals and advanced digital technologies is essential to mitigate the constraints and obstacles associated with organic agriculture (7). As organic farming employs a systems approach to control fertility, weeds and pests rather than depending on off-farm inputs, it also requires a high level of experience (7). To protect future generation, we need transition to organic farming and restore soil fertility.

Reason for using organic compounds in seed treatment

The use of chemical seed treatment products has a negative impact on the environment and human health. Before planting, the seeds should be treated with organic products to ensure high yield and good quality of seeds and thereby




Fig. 1. Types of organic compounds.

reducing the losses due to diseases. Organic seed treatment is currently being used in place of chemical seed treatment (10). For the conventional production of seeds, chemical fertilizers, insecticides, fungicides and herbicides are needed. The main reason in favour of utilizing organic materials for cultivating organic crops is that they are naturally occurring and has lower environmental impacts. In the process of producing organic seeds, various methods such as cropping pattern, organic manures, biofertilizers, cultural practices and biopesticides made from plant materials were used to maintain soil fertility, crop yield and pest control (Fig. 2) (11). Organic products have higher nutrient content and are often free of additives and pesticide residues (6). The use of organic products can improve soil pH, increase soil organic matter and enrich the soil with nutrients (12-14).

Cow products as organic compounds

Cow products include panchagavya, jeevamritta and beejamrita. In organic fertilizer, nitrogen percentage is increased due to the use of cow dung and cow urine as it contains rich source of amino acids. Cow urine, comprises of 95 % water, 25 % urea and 25 % essential minerals, hormones and enzymes (15). Other cow products serve as a rich source of nitrogen, phosphorus, potassium, calcium, magnesium, chloride and sulfate, contributing significantly to soil fertility and plant growth. It also enhances the total phenolic content, control pathogenic infection, improve soil health and plant growth (16). The cow based formulation contain majority of microbial sources like rhizobacteria that promote plant growth in paddy (Table 1) (17).




Fig. 2. Seed treatment with organic compounds.

Table 1. Effect of cow-based organic treatments on paddy seeds

S. No	Crop	Method of treatment	Result	Reference
1.	Rice CO 43	Seeds soaked in 4 % panchagavya for 16 hr	Significant rise in seedling growth parameters	(19)
2.	Rice ADT 43	Seeds treated with neem cake + panchagavya	Increased in seed quality and plant growth parameters	(67)
3.	Rice	Seeds treated with 10 % concentration of panchagavya	Germination decreases with increase in panchagavya Recorded maximum seedling length, fresh and dry weight	(20)
4.	Rice DRR Dhan 39	Seeds treated with 50 % vermicompost + jeevamrith 5 % + panchagavya 3 %	Significant increase in grain and straw yield	(79)

Panchagavya is a mixture of five by products of cow include cow dung, cow urine, milk, curd and ghee. These compounds aid in plant growth and have antimicrobial and nutritional quality (18). In Panchagavya, significant amount of nutrients, minerals and growth hormones are present, which enhances plant metabolic activity and promotes seed germination. In addition, panchagavya contains beneficial bacteria that possess biocontrol properties and produce substance essential for plant growth (19). Compared to vermicompost and farmyard manure, panchagavya contains higher levels of organic carbon and total nitrogen. Application of panchagavya significantly increase the chlorophyll content in plants. Wide range of organic C, N, P and K present in Panchagavya may be the cause for higher root and shoot growth (20). Biochemical properties of panchagavya showed that it contains almost all of the major nutrients, including N, P and K, as well as micronutrients and growth hormones, like gibberellic acid (GA) and indole acetic acid (IAA), which are essential for crop growth. It also contains a high concentration of fermentative microorganisms, such as *Lactobacillus*, *Azotobacter*, *Phosphobacteria* and yeast (21). The ammonia uptake and overall N supply are enhanced by the chemolithotrophs and autotrophic nitrifiers (ammonifiers and nitrifiers) found in panchagavya which proliferate on leaves (22).

To enhance crop nutrition, on-site produced Jeevamrit is essential, as it boosts soil microbial activity and improves soil fertility (23). Jeevamrit promotes enormous biological activity in the soil, thereby making nutrients more readily available to crops. Jeevamrit is a nutritious liquid bio-enhancer that improves soil productivity, growth and yield of plants (24). The soil microflora and soil enzymes are increased due to the use of beejamrit and jeevamrit which have microbial properties. Jeevamrit enhances the development of N fixing bacteria on the substrate like FYM and compost (25).

Beejamrit is a type of organic liquid manure that is used to treat crop seeds. It is a powerful organic mixture that protects the seed from various kinds of bacterial and fungal pathogens. Bijamrit or Beejamrita is a pre-sowing treatment for plants, seedlings or any other planting material. In addition to

protecting against soil-borne and seed-borne diseases that commonly affect crops after the monsoon season, it effectively protects young roots from fungal infection. Beejamrita is prepared using a variety of microorganisms, such as bacteria, fungi and nitrogen-fixing organisms, which improves seed protection. Beejamrita, enriched with diverse microorganisms like bacteria, fungi and nitrogen-fixers, enhances seed protection by fostering beneficial microbial activity. Its rich microbial composition safeguards seeds against various pests and insect infestations (26).

Botanicals - plant extracts

Botanicals have antioxidant properties and they are rich in cytokinin and zeatin, which increases plant growth (27). Alkaloids, protein, quinine, cytokinin, antioxidants and macro- and micronutrients are all present in *Moringa oleifera* leaves. These biochemicals support the growth, defence and metabolic processes of plants (Table 2) (28). Neem leaf extracts has antifungal properties and is used as a tool for controlling infection caused by seeds and promotes seed germination and seedling growth (27). In seed treatment, botanical agents are utilized to treat several phytopathological diseases and common insect pests that threaten seed storage, which improves the overall storability of seeds (29).

Seaweed extract contains antibiotics, vitamins, macro and micronutrients. This product when sprayed on the vegetative system, plays a vital role in controlling cell components and enhances the plant's resistance to environmental stresses like salt stress and drought stress. It also enhances leaf efficiency and speeds up carbon metabolism (30). In addition, the auxin and cytokinin present in seaweed extract promote elongation and enlargement of internodes and accelerate internal cell division, resulting in taller plants (31). When applied exogenously, seaweed extracts provide a range of plant growth hormones, polyamines, growth promoters, trace elements, vitamins, amino acids, antibiotics and micronutrients (32). Seaweed extract promotes tolerance against environmental challenges and increases plant nutrient uptake from soil (33).

Table 2. Effect of plant-based extracts on paddy seed germination and growth

S. No	Crop	Method of treatment	Result	Reference
1.	Rice	Seeds treated with 5 % solution of <i>Moringa oleifera</i> extract Seeds treated with 5 and 15 % solution of vermiwash	Increase germination, seedling vigour and biochemical content in seed	(18)
2.	Rice	Foliar spray of 3 % horse gram sprout extract and 2 % cowpea sprout extract	Observed higher seed germination, number of productive tillers/hill and seed yield	(74)
3.	ADT 43	Application of seaweed sap <i>Kappaphycus alvarezii</i> and <i>Gracilaria</i> sp. at 15 % + 100 % recommended dose of fertilizer	Increase in chlorophyll content, quality and yield	(80)
4.	ADT 53	Soil application of seaweed extract @ 12.5 kg/ha along with the foliar spray of seaweed extract liquid @ 0.5 % twice at tillering and panicle initiation stage	Recorded the highest value of growth and yield parameters	(81)

Bio-fertilizers and bio-control agents

Biofertilizers enhance soil fertility by supplying essential nutrients through nitrogen fixation, phosphate and potassium solubilization, mineralization and organic matter decomposition. They also promote plant growth by releasing phytohormones, synthesizing antibiotics and improving soil microbiota (34). Application of biofertilizers to seeds, seedlings, plants or soil helps mobilize plant nutrients for crop growth through biological nitrogen fixation, phosphorus solubilization or mobilization of any other plant nutrient required for crop growth. Biofertilizers are enriched with beneficial live microorganisms and formulated using sterilized carriers like lignite or talc, enhancing soil fertility and plant growth. To control a variety of root-borne infections, several biofertilizers also function as efficient biocontrol agents. They increase the fertility of the soil and make plants resistant to diseases (35). Bio-fertilizers establishes symbiotic relationship with plants and they are inoculated with live microorganisms that enhances major nutritional supply to plants and maintain soil fertility (36). They include *Azospirillum*, *Phosphobacteria*, *Rhizobium*, *Pseudomonas* and *Trichoderma* species. The use of biofertilizers increases plant height, boosts nitrogen use efficiency and increases the chlorophyll content in rice leaf tissue (Table 3) (37). Predominant PGPRs thriving in minimally or non-tilled soils include *Azotobacter*, *Azospirillum*, *Rhizobium*, cyanobacteria, phosphate solubilizers, potassium mobilizers and mycorrhizae, all of which contribute to improved soil fertility and plant growth (38). Productivity in rice is increased due to the application of *Azospirillum* (36). Benefits of applying biofertilizer include increased nutrient and water uptake from roots and also involve in biological nitrogen fixation (39).

Trichoderma is well known for its ability to promote growth and actively control plant diseases (40). It alters physiological processes in plants, enhancing the efficient use of water, light and nutrients (41). *Rhizobium* seed inoculation enhances nodulation, boosts seed yield, lowers fertilizer costs and helps mitigates soil degradation and environmental pollution caused by excessive chemical fertilizer use (35).

In many crop species, including rice, soil bacteria like *Pseudomonas* and *Azospirillum* are known to reduce abiotic stressors. One type of free-living fungi that interacts extensively with roots, soil and leaves is *Trichoderma* (42). It enhances nutrient absorption, stimulates seed germination even under adverse soil conditions and strengthens tolerance to abiotic stress. It secretes phosphate-solubilizing enzymes and phytohormones, inhibits harmful root microflora and promotes root growth, seedling vigour, plant development and overall yield improvement (43). By improving the availability of

minerals, particularly P and Zn to plants, *Trichoderma* stimulates growth and improves rice grain and vegetable yield even with minimal mineral fertilizer use (44). *Trichoderma* combined with 70 % NPK and burnt rice straw increased soil fertility, benefit-cost ratio and crop yield while lowering the cost of inorganic fertilizer (45). *Trichoderma* species that stimulate the production of auxins and gibberellins improve the performance of rice seedlings during seed germination (46). Increased nutrient uptake by *Trichoderma*-treated plants led to improved root and shoot growth as well as increased plant vigour, which ensured quick growth and increased plant greenness because of higher photosynthetic rates, which in turn raised the production of biomass and carbohydrates (47).

Azotobacter fixes atmospheric nitrogen in the rhizosphere, making it available to plants. In addition to nitrogen fixation, *Azotobacter* produces gibberellin, thiamine, riboflavin, nicotine and indole acetic acid. These compounds enhance seed germination and help reduce plant diseases. Seeds with low germination rates can be inoculated with *Azotobacter* to increase germination by 20-30 % (48). *Azospirillum* enhances plant growth by promoting development, enhancing nutrient absorption, improving biomass accumulation, increasing water uptake and ultimately boosting yield (49).

Organic manures

The integrated use of organic manures such as FYM, vermicompost, poultry manures and castor cake contain macro and micronutrients. They enhance the crop productivity and soil fertility (Table 4) (50). The application of organic manures improves soil health by enhancing its physical, chemical and biological properties (51). The main forms of organic matter that enhance soil fertility and help plants make better use of fertilizers are farmyard manure, press mud (a sugar waste product), green manure (*Sesbania aculeata*) and wheat residue. Farmyard manure boosts microbial activity, root penetration, water and nutrient retention and nutrient exchange capacity, thereby significantly improving soil fertility (52). The seeds treated with organic manure showed higher percentage of nitrogen in rice (53). Organic carriers, such as FYM, cow and poultry manures, improve plant development by changing the biological and physico-chemical properties of the soil (54). Due to its excellent capacity to store water, nutrients and buffers, peat has been utilized effectively as an organic carrier (55). Due to the presence of major nutrients and other essential elements like P and K, vermicompost promotes the growth of plants (56). Vermicompost, a potent organic fertilizer, enhances soil fertility and promotes sustainable crop growth. Its rich microbiota and bioactive compounds aid in suppressing pests and diseases, improving overall plant resilience (57).

Table 3. Effect of biofertilizers and biocontrol agents on paddy seed treatment

S. No	Crop	Method of treatment	Result	Reference
1.	Rice	Seeds treated with <i>Azospirillum brasilense</i> + <i>Bacillus megaterium</i> @ 4 kg/ha	Recorded higher seedling growth parameters	(82)
2.	Rice	Seeds treated with <i>Azospirillum</i> + <i>Phosphobacteria</i> + VAM at 600g/ha	Recorded higher values in growth and yield parameters	(76)
3.	Organic rice	<i>Azotobacter</i> (12.5 g/kg) + <i>Azospirillum</i> (12.5 g/kg) + FYM (12 t/ha)	Increased the yield and yield components of organic rice	(49)
4.	Rice	Foliar spray of <i>P. fluorescens</i> + butter milk (0.5 %) or <i>P. fluorescens</i> (0.5 %)	Reduced the incidence of sheath rot, false smut and grain discolouration in rice	(27)

Table 4. Impact of organic manure on paddy growth and yield

S. No	Crop	Method of treatment	Result	Reference
1.	Rice	Foliar application of vermiwash at 6 % twice during the active tillering and 50 % flowering stage	Increases productivity and achieves greater yield	(71)
2.	Rice	Foliar spray of egg amino acid 1 % + recommended dose of fertilizer	Recorded high leaf area index, soluble protein and yield	(65)
3.	Rice	Application of fish amino acid with a dose of 1mL	Increases the plant height, number of tillers and N uptake in plant leaves	(83)
4.	Bali red rice	Application of 2.5 % fish waste fertilizer	Increases the growth parameters in rice	(84)
5.	Rice	FYM + goat manure + compost manure @ 16.5 t/ha	Improves the grain quality	(85)
6.	Rice	Foliar application of vermiwash @ 5 %	Increases the growth parameters and nutrient uptake	(86)

Vermicompost, derived from earthworms and its liquid extract, vermiwash, enhance soil properties by increasing electrical conductivity, water retention and moisture content. It promotes superior plant growth and productivity (58). The physical, chemical and biological qualities of soil are enhanced by vermicompost (59). Vermiwash contains plant growth hormones such as auxins and cytokinins (58). The application of vermiwash may increase the level of chlorophyll and cytokinin, which prevent leaf senescence and extend the retention of effective photo assimilatory surface (60).

Humic substances, key sources of organic nitrogen and carbon, are naturally derived compounds formed through the decomposition of plant matter from terrestrial or marine origins (61). Their structure contains several phytohormones and exogenous auxins that induce plant growth (62). The stimulating effect observed in the humic substance extract is attributed to the presence of humic acid and fulvic acid. Due to their multifunctionality and diverse structure, humic substances include components that promote the activity of certain enzymes, as well as minerals, amino acids, phytohormones and beneficial microbes (63). Humic substances enhance plasma membrane H⁺-H-ATPase activity, facilitating H⁺ release into the soil, thereby reducing soil and root surface pH. This process optimizes nutrient solubility and absorption, improving overall plant nutrition (64).

Liquid organic manure generated from fish waste is known as fish amino acid. Fish contain various minerals and amino acids that are valuable to plants and microbes for their growth. Foliar and soil application of fish amino acid enhances nutrient uptake, providing the plant with adequate nitrogen and chlorophyll to support its growth and health (65). Fish

waste typically contains a variety of nutrients, including potassium (K), phosphorus (P) and nitrogen (N), making it a valuable component of organic fertilizers (66).

Effect of organic compounds on seed germination in paddy

Organic seed treatment in rice will increase germination rates and promote healthy seedling growth (Table 5). In Rice cultivar CO 43, the rate of imbibition was low in organic solutions of panchagavya, vermiwash, starter solution, cow's urine and coconut water when compared to water soaking (30.34 %) (19). On comparing the organic combination of neem cake + panchagavya, the inorganic treated seeds exhibited 5–8 % lower germination. The application of neem cake + panchagavya treatment in rice enhances germination and seed filling. The improvement in seed filling is reflected in a higher seed-to-husk ratio and increased 100-seed weight (67).

The seeds treated with various organic solutions at different levels will have significant rise in the vigour and germination compared to the control (19). Treating the seeds with different extracts of organic emulsion like *Parthenium hysterophorus*, *Moringa oleifera* leaves, vermiwash, panchagavya, *Ascophyllum nodosum* and *Medicago denticulate* at 5, 10, 15 % concentration increases germination of rice seeds (18). Application of organic amendments increases seed germination rate when compared to antibiotic control in plants, as demonstrated by the results of the Tukey's HSD test (68).

Rice seeds imbibed in humic substance derived from bovine manure vermicompost for 24 hr at 2 different dilutions recorded no effect on germination percentage but increased in seedling growth. This is because of humic substances on increasing the percentage of seeds that successfully germinate.

Table 5. Effect of organic seed treatment and foliar spray on paddy

S. No	Crop	Method of treatment	Result	Reference
1.	Rice	Application of rice husk, FYM and poultry litre mix with soil at 5 g/kg. After 2 weeks, antibiotics applied in aqueous solution form at 10 mg/kg.	Improves physiological growth parameters in rice	(68)
2.	Rice Sukhadhan 3, Tilkidhan	Seeds treated with 20 t FYM/ha. Before transplanting, the seedlings were dipped in bio inoculation of <i>Trichoderma</i> for 10-20 min.	Recorded maximum grain yield	(78)
3.	RiceBlackgram sequence	Application of vermicompost at 4 t/ha + foliar spray of panchagavya at 3 % at tillering and panicle initiation stage.	Recorded maximum plant height, number of tillers/ hills. Increases in dry matter production and nutrient uptake.	(86)
4.	Rice ADT 45	Application of neem cake + foliar spray of panchagavya @ 3 %	Increased the growth and seed quality parameters.	(87)
5.	Rice NDR 359	Application of vermicompost 5 t/ha + panchagavya 3 %	Recorded the highest plant growth and yield parameters.	(88)

Germination involves complex cellular and metabolic activities, starting from water absorption and culminating in the emergence of the radicle, without encompassing seedling growth (69).

In rice variety (Keteki Joha), the seeds are treated with five organic bioformulations namely organic metajal, organic trichoja, organic beauverijal, *Azospirillum* sp. and BIO phos liquid PSB formulation. Among all bioformulations, organic metajal showed higher seed germination percentage and seedling vigour (70).

Effect of organic compounds on growth and yield in paddy

Foliar application of jeevamrutha @ 10 % or 20 %, panchagavya @ 3 %, vermiwash @ 6 % and cow urine @ 100 % increases the productivity and economical returns in cereals and millets (71). When panchagavya and egg lime mix were applied to crops such as rice, wheat, bananas, vegetables, greens and fruit trees, the plants' longevity and productivity increased (72). Foliar spray alone did not provide enough nutrients for crop growth in sodic soil. However, when combined with the required fertilizer dosage, a 1 % foliar spray of fish and egg amino acids increased yield, soluble protein content and leaf area index (65).

In rice plants, foliar spray of horse gram and cowpea sprout extract at critical stages increases seed filling, productive tillers, number of panicles and seed yield (73). Foliar spray of horse gram sprout extract and cowpea sprout extract of different concentration @ 1, 2, 3 and 4 % recorded higher seed germination, number of productive tillers/hill and seed yield (74). In rice-blackgram sequence, the basal incorporation of green manure at 6.25 t ha⁻¹ along with vermicompost in four equal splits and foliar spray of panchagavya twice enhance growth and yield parameters (75).

When the rice seeds are inoculated with *Azospirillum* + *Phosphobacteria* + VAM, most of the growth and yield parameters are increased (76). The most effective pre-sowing treatment for enhancing seed germination and seedling vigor is to treat rice seeds with 4 % *P. fluorescens* for 12 hrs (77).

In rice variety ADT 45, seeds treated with various organic inputs recorded higher values of growth, yield and seed quality

parameters (5). In a pot experiment, soil was amended with organic inputs at 5 g/kg and after two weeks, antibiotics (ciprofloxacin, levofloxacin, ofloxacin, amoxicillin and ampicillin) were applied as an aqueous solution at 10 mg/kg. The study confirmed that the application of rice husk combined with antibiotics significantly improved physiological growth parameters in rice (Fig. 3) (68).

In the organic SRI method of rice cultivation, *Trichoderma* treatment showed greater yield, grain weight, plant height and more panicle per square meter. When *Trichoderma* inoculation and traditional cultivation techniques were used, the average yield increased by 12.5 % than untreated plot (78).

Conclusion

The transition to organic farming is a viable option for enhancing resilience to environmental pressures. Organic farming practices play high priority on biodiversity, soil health and preservation of natural resources. Reduced dependency on external inputs makes the farming system more self-sufficient and economically viable. This not only lowers production costs for farmers but also improves food security and promotes eco-friendly agriculture. Embracing organic farming fosters rural livelihood and community development, benefiting both farmers and society. By investing in research, education and policy support, we can accelerate the shift towards organic farming, fostering sustainable and fair agricultural sector that benefits future generation.

Authors' contributions

MS contributed to writing original draft preparation. NP was responsible for conceptualization. ASA and SE supervised the work and were involved in drafting and reviewing the manuscript. SG participated in sequence alignment, editing and visualization. All authors have read and approved the final version of the manuscript for publication.




Fig. 3. Effect of organic compounds on growth and yield of paddy.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

References

- Fukagawa NK, Ziska LH. Rice: Importance for global nutrition. *Journal of Nutritional Science and Vitaminology*. 2019;65 (Supplement): S2–3. <https://doi.org/10.3177/jnsv.65.S2>
- Shankar T, Banerjee M, Malik GC, Dutta S, Maiti D, Maitra S, et al. The productivity and nutrient use efficiency of rice-rice-black gram cropping sequence are influenced by location specific nutrient management. *Sustainability*. 2021;13(6):3222. <https://doi.org/10.3390/su13063222>
- Unicef, Organization WH. The state of food security and nutrition in the world 2017: Building resilience for peace and food security. 2017.
- Mondal R, Goswami S, Mandi SK, Goswami S. Quality seed production of rice (*Oryza sativa* L.) as influenced by nutrient management during kharif season in the lower Indo-Gangetic plains. *Environment and Ecology*. 2019;37(1A):274–80.
- Bhavani S, Saravanan K, Kumar S. Organic seed management techniques to improve seed yield and quality in rice. *Journal of Pharmacognosy and Phytochemistry*. 2017;6(6S):1158–63.
- Hammed TB, Oloruntoba EO, Ana G. Enhancing growth and yield of crops with nutrient-enriched organic fertilizer at wet and dry seasons in ensuring climate-smart agriculture. *International Journal of Recycling of Organic Waste in Agriculture*. 2019;8:81–92. <https://doi.org/10.1007/s40093-019-0274-6>
- Gamage A, Gangahagedara R, Gamage J, Jayasinghe N, Kodikara N, Suraweera P, et al. Role of organic farming for achieving sustainability in agriculture. *Farming System*. 2023;1(1):100005. <https://doi.org/10.1016/j.farsys.2023.100005>
- Mondal S, Mallikarjun M, Ghosh M, Ghosh DC, Timsina J. Effect of integrated nutrient management on growth and productivity of hybrid rice. *Journal of Agricultural Science and Technology*. 2015;5 (5):297–308. <https://doi.org/10.17265/2161-6264/2015.05.001>
- Azarbad H. Conventional vs. organic agriculture—which one promotes better yields and microbial resilience in rapidly changing climates? *Frontiers in Microbiology*. 2022;13:903500. <https://doi.org/10.3389/fmicb.2022.903500>
- Lamichhane JR, Corrales DC, Soltani E. Biological seed treatments promote crop establishment and yield: a global meta-analysis. *Agronomy for Sustainable Development*. 2022;42(3):45. <https://doi.org/10.1007/s13593-022-00761-z>
- Nabooji A, Veeranna HK, Nadukeri SV, Santosh SM, Shilpa HD. Organic seed production: Concept and practices. *The Pharma Innovation Journal*. 2022;11(10):1070–4.
- Huang QH, Li DM, Liu KL, Yu XC, Ye HC, Hu HW, et al. Effects of long-term organic amendments on soil organic carbon in a paddy field: A case study on red soil. *Journal of Integrative Agriculture*. 2014;13(3):570–6. [https://doi.org/10.1016/S2095-3119\(13\)60714-5](https://doi.org/10.1016/S2095-3119(13)60714-5)
- Qaswar M, Jing H, Ahmed W, Dongchu L, Shujun L, Lu Z, et al. Yield sustainability, soil organic carbon sequestration and nutrients balance under long-term combined application of manure and inorganic fertilizers in acidic paddy soil. *Soil and Tillage Research*. 2020;198:104569. <https://doi.org/10.1016/j.still.2019.104569>
- Wang H, Xu J, Liu X, Zhang D, Li L, Li W, et al. Effects of long-term application of organic fertilizer on improving organic matter content and retarding acidity in red soil from China. *Soil and Tillage Research*. 2019;195:104382. <https://doi.org/10.1016/j.still.2019.104382>
- Pradhan SS, Verma S, Kumari S, Singh Y. Bio-efficacy of cow urine on crop production. *International Journal of Chemical Studies*. 2018;6(3):298–301.
- Nautiyal CS, Mehta S, Singh HB, Mansinghka SB, Dawle SH, Rajhans NE, et al. Synergistic fermented plant growth promoting, bio-control composition. Council of Scientific, Industrial Research CSIR, assignee. United States patent US 7,297,659. 2007.
- Sreenivasa MN, Nagaraj MN, Bhat SN. Beejamruth: A source for beneficial bacteria. *Karnataka Journal of Agricultural Sciences*. 2010;17(3):72–7.
- Banjare S, Tripathi V, Dixit A, Verma A. Effect of selected weeds and organic emulsions on seed germination, seedling growth and biochemical content in rice (*Oryza sativa*). *The Indian Journal of Agricultural Sciences*. 2024;94(2):211–5. <https://doi.org/10.56093/ijas.v94i2.143627>
- Kumar GD, Narayanan GS. Efficacy of organic seed invigoration on seed germination and vigour in rice cv. Co 43. *Journal of Environment and Bio-Sciences*. 2014;28(2):245–8.
- Jain P, Sharma RC, Bhattacharyya P, Banik P. Effect of new organic supplement (Panchgavya) on seed germination and soil quality. *Environmental Monitoring and Assessment*. 2014;186:1999–2011. <https://doi.org/10.1007/s10661-013-3513-8>
- Golakiya P, Kaswala A, Dubey P, Sisodiyaan R, Neethu T. Panchagavya-cow-based bio-enhancers and botanicals for organic farming. *International Journal of Chemical Studies*. 2019;7(3):2741–6.
- Papen H, Geßler A, Zumbusch E, Rennenberg H. Chemolithoautotrophic nitrifiers in the phyllosphere of a spruce ecosystem receiving high atmospheric nitrogen input. *Current Microbiology*. 2002;44:56–60. <https://doi.org/10.1007/s00284-001-0074-9>
- Saharan BS, Tyagi S, Kumar R, Vijay, Om H, Mandal BS, et al. Application of jeevamrit improves soil properties in zero budget natural farming fields. *Agriculture*. 2023;13(1):196. <https://doi.org/10.3390/agriculture13010196>
- Kumar S, Trivedi H, Sah R, Verma AK, Yadav A. Effect of different bioenhancers on growth and yield of cauliflower [*Brassica oleracea* (L.) var. Botrytis]. *Journal of Pharmacognosy and Phytochemistry*. 2018;1:769–72.
- Devakumar N, Rao G, Shubha S. Evaluation of locally available media for the growth and development of nitrogen fixing microorganisms. In: *Proceedings of the Third Scientific Conference of the International Society of Organic Agriculture Research (ISO FAR), Korean Organizing Committee (KOC)*. Namyangju, Korea Republic; 2011. p. 504–9.
- Shyamsunder B, Menon S. Study of traditional organic preparation beejamrita for seed treatment. *International Journal of Modern Agriculture*. 2021;10(2):1823–8.
- Jeyalakshmi C, Renuka R, Ramanadane T. Evaluation of foliar sprays of biocontrol agents, botanicals and fungicides for the production of healthy rice seed in the coastal ecosystem. *Journal of Cereal Research*. 2021;13(S1):77–83. <https://doi.org/10.25174/2582-2675/2021/115310>
- Paikra BK, Gidwani B. Phytochemistry and pharmacology of *Moringa oleifera* Lam. *Journal of Pharmacopuncture*. 2017;20 (3):194. <https://doi.org/10.3831/KPI.2017.20.022>
- Umesha U, Gupta N, Bhanuprakash K, Sharma V. Seed treatment with botanicals: an indigenous technology. *Indian Farming*. 2018;68(6).
- Khuit SA, Ahmed AS, Radhi DA. The partial replacement of seaweed extract and humic acid instead of chemical fertilizers and its effect on the growth and yield of rice cultivars. *NeuroQuantology*. 2022;20(10):1580.

31. Mahdi DH, Madlol FM, Hashim MA. Effect of humic acid spraying on growth and yield of wheat (*Triticum aestivum* L.). Plant Archives. 2019;19(1):1291–3.
32. Panda D, Pramanik K, Nayak B. Use of sea weed extracts as plant growth regulators for sustainable agriculture. International Journal of Bio-resource and Stress Management. 2012;3(3):404–11.
33. Yadav S, Verma A, Nepalia V. Effect of phosphorus, sulphur and seaweed sap on growth, yield and nutrient uptake of chickpea (*Cicer arietinum* L.). Research on Crops. 2016;17(3):496–502. <https://doi.org/10.5958/2348-7542.2016.00082.6>
34. Sinha RK, Valani D, Chauhan K, Agarwal S. Embarking on a second green revolution for sustainable agriculture by vermiculture biotechnology using earthworms: reviving the dreams of Sir Charles Darwin. Journal of Agricultural Biotechnology and Sustainable Development. 2010;2(7):113.
35. Dubey M, Verma V, Barpete R, Verma N. Effect of biofertilizers on growth of different crops: a review. Plant Archives. 2019;19:1083–6.
36. Roychowdhury D, Paul M, Banerjee SK. A review on the effects of biofertilizers and biopesticides on rice and tea cultivation and productivity. International Journal of Science, Engineering and Technology. 2014;2(8):96–105.
37. Khan HI. Appraisal of biofertilizers in rice: To supplement inorganic chemical fertilizer. Rice Science. 2018;25(6):357–62. <https://doi.org/10.1016/j.rsci.2018.10.006>
38. Dogan K, Celik I, Gok M, Coskan A. Effect of different soil tillage methods on rhizobial nodulation, biyomas and nitrogen content of second crop soybean. African Journal of Microbiology Research. 2011;5(20):186–94. <https://doi.org/10.5897/AJMR11.165>
39. Naseri R, Mirzaei A. Response of yield and yield components of Safflower (*Carthamus tinctorius* L.) to seed inoculation with *Azotobacter* and *Azospirillum* and different nitrogen levels under dry land condition. American-Eurasian Journal of Agricultural & Environmental Sciences. 2010;9(4):445–9.
40. Diánez Martínez F, Santos M, Carretero F, Marín F. *Trichoderma saturnisporum*, a new biological control agent. Journal of the Science of Food and Agriculture. 2016;96(6):1934–44. <https://doi.org/10.1002/jsfa.7301>
41. Doni F, Zain CR, Isahak A, Fathurrahman F, Sulaiman N, Uphoff N, et al. Relationships observed between *Trichoderma* inoculation and characteristics of rice grown under System of Rice Intensification (SRI) vs. conventional methods of cultivation. Symbiosis. 2017;72:45–59. <https://doi.org/10.1007/s13199-016-0438-3>
42. Vejan P, Abdullah R, Khadiran T, Ismail S, Nasrulhaq Boyce A. Role of plant growth promoting rhizobacteria in agricultural sustainability-a review. Molecules. 2016;21(5):573. <https://doi.org/10.3390/molecules21050573>
43. Harman GE, Howell CR, Viterbo A, Chet I, Lorito M. *Trichoderma* species-opportunistic, avirulent plant symbionts. Nature Reviews Microbiology. 2004;2(1):43–56. <https://doi.org/10.1038/nrmicro797>
44. Rahman M, Ali J, Masood M. Seed priming and *Trichoderma* application: a method for improving seedling establishment and yield of dry direct seeded boro (winter) rice in Bangladesh. University Journal Agriculture. 2015;3(2):59–67. <https://doi.org/10.13189/ujar.2015.030205>
45. Debnath S, Chakraborty G, Dutta SS, Chaudhuri SR, Das P, Saha AK. Potential of *Trichoderma* species as biofertilizer and biological control on *Oryza sativa* L. cultivation. Biotechnología Vegetal. 2020;20(1):1–16.
46. Doni F, Anizan I, Radziah CC, Salman AH, Rodzihan MH, Yusoff WM. Enhancement of rice seed germination and vigour by *Trichoderma* spp. Research Journal of Applied Sciences, Engineering and Technology. 2014;7(21):4547–52. <https://doi.org/10.19026/rjaset.7.832>
47. Saba H, Vibhash D, Manisha M, Prashant K, Farhan H, Tauseef A. *Trichoderma*—a promising plant growth stimulator and biocontrol agent. Mycosphere. 2012;3(4):524–31. <https://doi.org/10.5943/mycosphere/3/4/14>
48. Gayatri AK, Shri Om Narayan Verma and Uttara Sahu. Effect of *Azotobacter* and phosphate solubilizing bacteria on the yield of different wheat (*Triticum aestivum* L.) Cultivar. The Pharma Innovation Journal. 2022;11(1):1629–33.
49. Mugalavai VK, Aduol KO, Onkware AO. Nutritional characteristics of rice (*Oryza sativa* L.) composite flours obtained by food fortification. European Journal of Agriculture and Food Sciences. 2021;3(1). <https://doi.org/10.24018/ejfood.2021.3.1.224>
50. Arya R, Niranjana K, Singh A, Singh J. Production potential and sustain ability of food-fodder alley cropping system under rainfed conditions. The Indian Journal of Agricultural Sciences. 2000;70(2).
51. Damse D, Bhalekar M, Pawar P. Effect of integrated nutrient management on growth and yield of garlic. The Bioscan. 2014;9(4):1557–60.
52. Yazdanpanah. CO₂ emission and structural characteristics of two calcareous soils amended with municipal solid waste and plant residue. Solid Earth. 2016;7(1):105–14. <https://doi.org/10.5194/se-7-105-2016>
53. Ashrafi R, Biswas M, Rahman G, Khatun R, Islam M. Effect of organic manure on nutrient contents of rice grown in an arsenic contaminated soil. Bangladesh Journal of Scientific and Industrial Research. 2010;45(3):183–8. <https://doi.org/10.3329/bjsir.v45i3.6526>
54. Ebaid R, El-Refaei I. Utilization of rice husk as an organic fertilizer to improve productivity and water use efficiency in rice fields. African Crop Science Conference Proceedings. 2007;8:1923–8.
55. Saha AK, Deshpande MV, Kapadnis BP. Studies on survival of *Rhizobium* in the carriers at different temperatures using green fluorescent protein marker. Current Science. 2001;669–71.
56. Fernández-Luqueño F, Reyes-Varela V, Martínez-Suárez C, Salomón-Hernández G, Yáñez-Meneses J, Ceballos-Ramírez J, et al. Effect of different nitrogen sources on plant characteristics and yield of common bean (*Phaseolus vulgaris* L.). Bioresource Technology. 2010;101(1):396–403. <https://doi.org/10.1016/j.biortech.2009.07.058>
57. Rakesh J, Jaswinder Singh JS, Vig A. Vermicompost as an effective organic fertilizer and biocontrol agent: effect on growth, yield and quality of plants. Reviews in Environmental Science and Bio/Technology. 2015;14(1):137–59. <https://doi.org/10.1007/s11157-014-9347-1>
58. Tharmaraj K, Ganesh P, Kolanjinathan K, Suresh K, Anandan A. Influence of vermicompost and vermiwash on physico chemical properties of rice cultivated soil. Current Botany. 2011;2(3):18–21.
59. Ansari A. Urban Planning and Environment—Strategies and Challenges. New Delhi: Macmillan India Ltd; 2007. p. 277–9.
60. Nath G, Singh K. Effect of foliar spray of biopesticides and vermiwash of animal, agro and kitchen wastes on soybean (*Glycine max* L.) crop. Botany Research International. 2011;4(3):52–7.
61. Madende M, Hayes M. Fish by-product use as biostimulants: An overview of the current state of the art, including relevant legislation and regulations within the EU and USA. Molecules. 2020;25(5):1122. <https://doi.org/10.3390/molecules25051122>
62. Nardi S, Pizzeghello D, Schiavon M, Ertani A. Plant biostimulants: physiological responses induced by protein hydrolyzed-based products and humic substances in plant metabolism. Scientia Agricola. 2016;73:18–23. <https://doi.org/10.1590/0103-9016-2015-0006>
63. García AC, Izquierdo FG, Balmori DM, de Armas MM, Barrueta MA, Rebato SM, et al. Vermicompost-derived liquid humus in low-input and small-scale farming. In: Bates L, editor. Humic Substances and Natural Organic Matter. New York: Nova Publishers; 2015. p. 1–18.

64. Canellas LP, Olivares FL. Physiological responses to humic substances as plant growth promoter. *Chemical and Biological Technologies in Agriculture*. 2014;1:1–11. <https://doi.org/10.1186/2196-5641-1-3>
65. Priyanka B, Ramesh T, Rathika S, Balasubramaniam P. Foliar application of fish amino acid and egg amino acid to improve the physiological parameters of rice. *International Journal of Current Microbiology and Applied Sciences*. 2019;8(2):3005–9. <https://doi.org/10.20546/ijcmas.2019.802.351>
66. Saputra R, Sari N, editors. Ameliorant engineering to elevate soil pH, growth and productivity of paddy on peat and tidal land. IOP Conference series: earth and environmental science; IOP Publishing; 2021. <https://doi.org/10.1088/1755-1315/648/1/012183>
67. Vijayan R, Krishnasamy V. Impact of organic techniques of seed crop management on seed yield and quality in rice cv. ADT 43. *Scientific Research and Essays*. 2014;9(13):611–8. <https://doi.org/10.5897/SRE2013.5606>
68. Mukhtar A, Manzoor M, Gul I, Zafar R, Jamil HI, Niazi AK, et al. Phytotoxicity of different antibiotics to rice and stress alleviation upon application of organic amendments. *Chemosphere*. 2020;258:127353. <https://doi.org/10.1016/j.chemosphere.2020.127353>
69. Galbán-Méndez JM, Martínez-Balmori D, González-Viera D, San José de las Lajas M. Effect of extracts of humic substances on germination and growth of rice plant (*Oryza sativa* L.), cv. INCA LP-5. *Cultivos Tropicales*. 2021;42(1):e05.
70. Nasrin S, Deka S, Sharma P, Medhi T, Das K, Dutta P. Performance evaluation of pre-sowing seed treatments using bio-agents in transplanted aromatic rice under organic conditions. *Oryza*. 2023;60(3):397–405. <https://doi.org/10.35709/ory.2023.60.3.3>
71. Raman, Krishnamoorthy. Impact of organic foliar nutrition and its efficacy on sustainable production of rice. *Acta Scientific Agriculture*. 2019;3(10):61–3. <https://doi.org/10.31080/ASAG.2019.03.0647>
72. Prabu MJ. Muttai rasam: An excellent plant growth promoter. *Sci & Tech*. Tamil nadu The Hindu. 2008;18:2008.
73. Kavipriya R, Dhanalakshmi P, Jayashree S, Thangaraju N. Seaweed extract as a biostimulant for legume crop, green gram. *Journal of Ecobiotechnology*. 2011;3(8):16–9.
74. Veerappan V, Rangnathan U, Mannar J. Effect of organic foliar spray with pulse sprout extract on seed yield and quality of rice (*Oryza sativa*). *Journal of Plant Nutrition*. 2019;42(8):900–14. <https://doi.org/10.1080/01904167.2019.1567764>
75. Thiagarajan M, Somasundaram E. Residual effect of organic manures on growth and yield of black gram in rice blackgram sequence. *Madras Agricultural Journal*. 2019;106(4–6):369. <https://doi.org/10.29321/MAJ.2019.000276>
76. Anand S, Kamaraj A. Influence of pre sowing biofertilizer seed treatment on growth and yield parameters in rice (*Oryza sativa* L.). *Plant Archives*. 2017;17(2):1377–80.
77. Somasundaram G, Bhaskaran M. Effect of seed priming on germination and vigour in low and high longevity rice genotypes. *International Journal of Agricultural Science and Research*. 2017;7(2):373–80.
78. Khadka RB, Uphoff N. Effects of *Trichoderma* seedling treatment with system of rice intensification management and with conventional management of transplanted rice. *Peer J*. 2019;7:e5877. <https://doi.org/10.7717/peerj.5877>
79. Sharada P, Sujathamma P. Effect of organic and inorganic fertilizers on the quantitative and qualitative parameters of rice (*Oryza sativa* L.). *Current Agriculture Research Journal*. 2018;6(2):166–74. <https://doi.org/10.12944/CARJ.6.2.05>
80. Devi NL, Mani S. Effect of seaweed saps *Kappaphycus alvarezii* and *Gracilaria* on growth, yield and quality of rice. *Indian Journal of Science and Technology*. 2015;8(19):1–6. <https://doi.org/10.17485/ijst/2015/v8i19/47610>
81. Deepana P, Bama KS, Santhy P, Devi TS. Effect of seaweed extract on rice (*Oryza sativa* var. ADT53) productivity and soil fertility in Cauvery delta zone of Tamil Nadu, India. *Journal of Applied and Natural Science*. 2021;13(3):1111. <https://doi.org/10.31018/jans.v13i3.2906>
82. K Sankar A. Effect of FYM, N, P fertilizers and biofertilizers on germination and growth of paddy (*Oryza sativa* L.). *International Letters of Natural Sciences*. 2013.
83. Jumar J, Saputra RA, Jannah SR. Effect of fish amino acid application on growth and n-uptake in plants rice using the system of rice intensification method. *Tropical Wetland Journal*. 2021;7(1):25–30. <https://doi.org/10.20527/twj.v7i1.91>
84. Kusuma ID, Suriani NL, Ramona Y. The use of fish waste based organic fertilizer to improve the growth of balinese red rice (*Oryza sativa* L. Cv. Barak Cenana). *Asian Journal of Applied Research for Community Development and Empowerment*. 2021;5(2):13–8. <https://doi.org/10.29165/ajarcde.v5i2.67>
85. Punitha M, Jeyamangalam F, Selvam V. The effect of farm yard manure combinations with goat manure and compost manure on yield and nutrient uptake of rice. *International Journal of Mechanical Engineering*. 2021;6:285–90.
86. Saranraj T, Imayavaramban V. Effect of liquid organic manures on growth attributes and nutrient uptake of rice (*Oryza sativa*). *Trends in Biosciences*. 2015;8(5):1216–9.
87. Sathiyarayan G, Mallavarapu Geetha L, Satya Bhavani S. Organic seed management techniques to improve seed yield and quality in rice. *Journal of Pharmacognosy and Phytochemistry*. 2017;SP1:1158–63.
88. Ganesh KS, Mehera B, Kumar P. Impact of organic nutrient sources on growth and yield of rice varieties (*Oryza sativa* L.) and yield validation using SPSS model. *International Journal of Environment and Climate Change*. 2023;13(7):252–8. <https://doi.org/10.9734/ijecc/2023/v13i71874>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc. See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.