



REVIEW ARTICLE

# Phytochemical and pharmacological advancement on *Tridax procumbens* L.: A traditional medicinal plant with multiple health benefits: A review

Harpreet Kaur<sup>1</sup>, Agam Bori<sup>1</sup>, Kuldip Bora<sup>1</sup>, Kangkana Medhi<sup>1</sup>, Gongwna Basumatary<sup>1</sup>, Natesan Balasubramanian<sup>1\*</sup>, Jagdish Kumar Sahu<sup>2</sup> & Gomathinayagam Subramanian<sup>3</sup>

<sup>1</sup>Department of Botany, Tripura University (A Central University), Suryamaninagar, Agartala 799 022, Tripura, India

<sup>2</sup>Department of Pharmacy, Tripura University (A Central University), Suryamaninagar, Agartala 799 022, Tripura, India

<sup>3</sup>Division of Agriculture, University of Guyana, Berbice Campus, Berbice, Guyana, South America

\*Correspondence email - [balasubramanian.natesan@tripurauniv.ac.in](mailto:balasubramanian.natesan@tripurauniv.ac.in)

Received: 10 April 2025; Accepted: 14 October 2025; Available online: Version 1.0: 11 March 2026; Version 2.0: 01 April 2026

**Cite this article:** Harpreet K, Agam B, Kuldip B, Kangkana M, Gongwna B, Natesan B, Jagdish KS, Gomathinayagam S. Phytochemical and pharmacological advancement on *Tridax procumbens* L.: A traditional medicinal plant with multiple health benefits: A review. *Plant Science Today*. 2026; 13(2): 1-12. <https://doi.org/10.14719/pst.8791>

## Abstract

*Tridax procumbens* L. is a persistent herb typically found as an agronomic weed that grows on wasteland or invades bare soil. Indigenous medicine has employed several extracts of *T. procumbens* to treat an array of illnesses in both humans and animals. The primary objective of this study is to comprehensively review the phytochemical constituents and medicinal potential of *T. procumbens*, highlighting its traditional and pharmacological uses with updated literature. The study aims to consolidate scientific literature on *T. procumbens*, its biological activities and therapeutic applications to provide a valuable ethnomedicinal reference. Despite the traditional and ethnomedicinal significance of *T. procumbens*, several critical research gaps remain, including a lack of in-depth phytochemical characterization. Bridging these gaps is essential for advancing *T. procumbens* from traditional use to evidence-based therapeutic applications. The literature review confirms the presence of several crucial phytochemicals in *T. procumbens* such as alkaloids, tannins, flavonoids, phenols, proteins, anthocyanins and amino acids. These compounds are associated with various biological activities including wound healing, antioxidant and antibacterial effects, anti-inflammatory properties, antifungal, antiparasitic, insecticidal and immunomodulatory therapeutic effects. *Tridax procumbens* demonstrates significant potential as a medicinal plant with a broad range of biological activities attributed to its diverse phytochemical profile. However, further in-depth pharmacological studies, clinical investigations and toxicological assessments are essential to validate its ethnomedicinal claims and develop it into a reliable therapeutic agent. This review underscores the need for advanced research to bridge the gap between traditional knowledge and modern evidence-based medicine.

**Keywords:** ethnomedicinal; pharmacological; phytochemical; medicinal plant; *Tridax procumbens*

## Introduction

Nature has generously provided us with various therapeutic plants from which many contemporary medications have already been extracted (1). According to World Health Organization (WHO), over 80 % of people practice some form of traditional medicine and herbal medicine for their basic well-being (2). Since pre-Hispanic times, indigenous doctors have utilized medicinal plants, which are a part of human traditional practice (3). Plant secondary metabolites with pharmacological qualities are helpful; some are essential drug components for antitumor, antivirals, antiepileptics, antibiotics and more (4,5).

The *Tridax procumbens* is also recognized as “coat buttons,” a perennial plant belonging to the family Asteraceae, indigenous to South and Central America (6, 7). *Tridax* species have been utilized in Indian Ayurveda since ancient times (8). *Tridax procumbens* are found throughout India, especially in Maharashtra, Madhya Pradesh, Chhattisgarh, Tamil Nadu, Kerala, Rajasthan and Uttar Pradesh and

more regions as a weed. *Tridax procumbens* has various pharmacological properties like antimicrobial, immunomodulatory, antioxidant, antidiabetic, analgesic and antifungal activities (9).

Developed and developing nations frequently employ herbal remedies or traditional medicines to treat various illnesses. Synthetic medication side effects account for 8 % hospital admissions in the USA and these toxicities cause around 100000 deaths annually (9, 10). Therefore, natural medications that minimise synthetic side effects and toxicities and simultaneously increase therapeutic potential to cure various diseases are increasingly being explored. The key objective of this study is to comprehensively review the phytochemical constituents and medicinal potential of *T. procumbens* and highlight its traditional and pharmacological uses with updated literature. Few manuscripts have been published in *T. procumbens* related to phytochemicals and pharmacological activity; however, the present review covers all aspects of the plant's biological activity with updated information. This review bridges the gap between traditional knowledge and

modern evidence-based medicine. Finally, this review provides significant information on this species. It specifies phytochemicals for specific treatments for safe and low-cost treatment of some ailments, especially in tropical and sub-tropical areas. The taxonomical classification of *T. procumbens* is presented below to provide systematic and botanical details of the species (11).

### Taxonomical classification of *T. procumbens*

Kingdom	: Plantae
Phylum	: Streptophyta
Class	: Equisetopsida
Subclass	: Magnoliidae
Order	: Asterales
Family	: Asteraceae
Genus	: <i>Tridax</i>
Species	: <i>procumbens</i>

### Distribution

The *T. procumbens* belongs to the family Asteraceae, native to Central and South America and is an extensively distributed herb that generally grows like grass (12). This genus includes about 26 species (13). *Tridax procumbens* mainly originated in Central America but now can be found throughout the tropical and subtropical areas of the world growing with annual crops, along roadsides, pastures, fallow land, dikes, railroads, riverbanks, meadows, dunes and waste areas (9). *Tridax procumbens* is perennial, blossoming and producing fruit usually named in Hindi 'Ghamra', in English 'Wild daisy' and 'Coat buttons' and in Sanskrit as 'Jayanti-veda' based on the presence of the flower (14) (Table 1 & Fig. 1). *Tridax procumbens* is extensively spread in India and this plant thrives in various soil types, including dry and moist conditions (12, 15).

### Traditional and ethnomedicinal applications

Coat Button (*T. procumbens*) has been used in folk remedies and practices for various health benefits. The multiple studies highlight the widespread use of Coat Button in traditional medicine, which suggests it may hold promise as a potential analgesic agent (19). The plant is prevalent throughout India and is utilised in indigenous medicine for various ailments. In Ayurvedic medicine, it is widely used due to its medicinal properties, including reducing blood

pressure, bronchial catarrh, malaria, dysentery, diarrhoea, stomach-ache, wound healing, headaches and preventing hair loss and haemorrhage from cuts and bruises (19). This versatile herb has also been associated with treating various diseases, including malaria, leishmaniasis and dysentery. Up-to-date recent research studies have noted that lactating pregnant women in Chiquimula, Guatemala, suffering from anemia could alleviate their symptoms by using *T. procumbens* (6). Aqueous extracts of Coat Button exhibit significant activity against *Trypanosoma brucei*, along with antibacterial and wound-healing properties (20). The flowers and leaves possess antiseptic, insecticidal and parasiticidal properties and the plant exhibits various pharmacological activities such as immunomodulatory, antidiabetic, antihepatotoxic, antioxidant, anti-inflammatory, analgesic, anticancer and marked depressant action on respiration (21). It plays a crucial role in defluoridation water, providing an inexpensive method in regions where the natural level of fluoride minerals is high in groundwater (22).

Typhoid fever, cough, fever, backache, diarrhoea and epilepsy are treated with the entire plant of *T. procumbens* in Nigeria; besides, this plant is used by African farmers to treat livestock (6). A study in Tamil Nadu, India, revealed that native inhabitants apply the leaf juice to heal wounds. In Udaipur in Rajasthan, India, people traditionally consume powdered *T. procumbens* leaves mixed with other herbs to manage diabetes. In some other parts of the world, traditional healers and local communities also use the leaves of this plant as a treatment for conjunctivitis (9).

### Phytochemical screening

Phytochemicals are chemical compounds naturally present in plants and are credited with positive or adverse health effects (23). Phytochemical screening identifies and analyses the chemical compounds or phytochemicals, present in plant extracts (24). These compounds, often secondary metabolites, can be investigated for their potential medicinal or biological activities. The screening involves various qualitative and quantitative tests to detect the presence and estimate the amounts of different phytochemicals (25). The phytochemical components of the plants have healing qualities (26). Flavonoids, alkaloids, tannins, phenolics, steroids, saponins, terpenes, glycosides and other vital phytochemicals are found in different plant parts. Besides, fatty acid derivatives, lipid constituents—luteolin, glucoluteolin and quercetin are described (27, 28). In *T. procumbens*, diverse phytochemicals were reported

**Table 1.** List of common names of *Tridax procumbens* throughout the world

Country/region	Common names	References
English	Coat buttons, Tridax daisy	(7, 16)
Chinese	Kotobukigiku	
Sanskrit	Kumminnippacha	
Latin	<i>Tridax procumbens</i>	
Kannada	Jayanthi	
Japanese	Kotobukigiku	
Spanish	Cadillochisaca	
Gujrati	Ghaburi	
Telegu	Gaddi chemanthi	
Tamil	Thatapoodu	
United States	<i>Tridax</i> daisy	(9)
Assamese	Bikhalyakarani	
Hindi	Ghamra	
Bengali	Tridhara	
Oriya	Bishalyakarani	
Malayalam	Chiravanak	
French	Herbecailli	
Marathi	Kambarmodi, Jakhamjudi and Tantani	
Thai	Tintúkkæ	
Nigeria	Igbalobe, muwagun, muriyampachila, jayanti, vettukkaaya-thala	(17)
Guatemala	Bull grass, bull's herb	(18)



**Fig. 1.** Morphology of Leaves, bud, flower and stem of *Tridax procumbens*.

**Table 2.** Different solvent extraction using different parts of *Tridax procumbens*

Extraction	Plant part	Compounds	References
Chloroform extract and Chloroform water extract	Leaves	Steroid, saponin, coumarins, amino acids, diterpenes, phenol and flavonoids, amino acids, phlobatannin	
Acetone-water extract and acetone extract	Leaves	Steroid, tannin, saponin, anthocyanin, coumarins, alkaloids, diterpenes, phenol and flavonoids, proteins, carbohydrate, antioxidant property	
Aqueous extract	Leaves	Blood clotting properties	(9)
Acetone extracts	Roots, leaves	Antibacterial activity	
Methanol extract	Leaves	Alkaloids, tannin, anthocyanins, proteins, saponin, steroid, phlobatannin, terpenoids, flavonoids, amino acids, phenols and cardiac glycosides, antibacterial activity, antioxidant properties	(29)
Ethanol extract	Whole plant	Flavonoid, quercetin, alkaloids, tannins, flavonoids, saponins and phenolic compounds	(30)
Petroleum ether and ethanolic extract	Whole plant	Antibacterial activity against <i>B. faecalis</i> due to presence of alkaloids, tannins, steroids, purines, carbohydrates, proteins	(31)
Chloroform extract	Whole plant	Against <i>B. faecalis</i> and <i>E. coli</i>	
Aqueous extract	Leaves	DPPH radical scavenging activity	(32)
Ethanol extract	Leaves	Polyphenol content, flavonoids, antibacterial activity against <i>Pseudomonas aeruginosa</i>	(32, 33)
Methanol extract fractionated with Dichloromethane (DCM)	Aerial parts of plant	9, 12-octadecadienoic acid ethyl ester (18.04 %), 5-cholestane (12.42 %), hexadecanoic acid ethyl ester (4.86 %) and 9-octadecenoic acid ethyl ester (4.72 %). Cholestane glycosides and rhamnosides are known for their potent cytotoxicity against malignant tumor cells.	(34)

from different parts of the plants with various biological activities (Table 2).

### Flavonoids

Flavonoids are a large group of naturally occurring plant compounds in the polyphenol class. It can be detected by the Alkaline reagent test,  $\text{NH}_4\text{OH}$  test, Mg turning test and Zn test (35). Flavonoids constitute a major and repeatedly reported class in *T. procumbens*. A novel flavonoid glycoside named procumbenetin (3,6-dimethoxy-5,7,2',3',4'-pentahydroxyflavone 7

-O- $\beta$ -D glucopyranoside) was isolated and fully characterised from aerial parts, an essential structural finding that demonstrates the plant's unique flavonoid profile. Other flavonoids reported in various surveys include apigenin, quercetin, luteolin and related derivatives. These polyphenolic molecules are frequently linked to antioxidant and anti-inflammatory properties (36, 37).

### Alkaloids

Alkaloids are a large group of naturally occurring nitrogen-containing organic compounds that plants produce as secondary

metabolites. Dragendroff's, Mayer, Wagner and Hager's tests were used to detect alkaloids (35). Alkaloids have been reported in aqueous, ethanolic, methanolic and chloroform extracts of leaves, stems and flowers, confirmed their presence in both aqueous and ethanolic extracts of leaves and identified nitrogen-containing peaks in GC-MS profiles of methanolic extracts, suggestive of heterocyclic alkaloids (38, 39). Alkaloids may contribute to the plant's antimicrobial and wound-healing activities.

### Tannins

Tannins are a group of naturally occurring polyphenolic compounds found in many plants. The Lead acetate  $[Pb(CH_3COO)_2]$  and Iron(III) chloride  $[FeCl_3]$  (5 %) test can detect it (35). Tannins are consistently reported in qualitative screenings of *T. procumbens* leaves, stems and flowers. Tannins exhibit antibacterial and antifungal properties, enhancing the plant's traditional medicinal use in treating infections. They also help scavenge free radicals, thereby protecting cells from oxidative damage and may reduce inflammation, supporting their role in wound healing and the management of various inflammatory conditions (40) (Table 3 & Fig. 2a, b).

### Terpenoids

Terpenoids (also called isoprenoids) are a large and diverse group of naturally occurring organic compounds which can be detected by mixing about 5 mL of each extract in 2 mL of chloroform and adding 3 mL of concentrated  $H_2SO_4$ , which forms a reddish-brown precipitate indicator (41). Pentacyclic triterpenes such as lupeol, oleanolic acid and taraxasteryl acetate have been isolated from *T. procumbens*. Terpenoids are also detected in preliminary phytochemical screenings of methanolic, ethanolic and hexane extracts. These compounds have been linked to anti-inflammatory, hepatoprotective and antimicrobial effects (42).

### Saponins

Saponins are a group of naturally occurring plant compounds known for their soap-like properties. It can be detected by the Foam Test and the Haemolysis Test (35). Saponins have been detected in leaves and whole plants' aqueous, methanolic and ethanolic extracts. These glycosides possess surfactant properties and contribute to antimicrobial and anti-inflammatory actions. Their foaming ability in aqueous extracts is a typical qualitative test result for *T. procumbens* (43).

### Phenolic Compounds

Phenolic compounds are a large and diverse group of plant chemicals (phytochemicals) that have one or more phenol groups in their structure, which can be detected by the formation of intense colour on the addition of 0.5 mL of  $FeCl_3$  (w/v) solution into 2 mL of test solution, indicating the presence of phenols (44). Phenolics, including simple phenols and complex polyphenols, are abundant in *T. procumbens* extracts. They are major contributors to the plant's antioxidant potential. Quantitative assays such as the Folin-Ciocalteu method have shown significant phenolic content, with flavonoids making up a large proportion. Phenolic compounds act synergistically with flavonoids and tannins to provide free radical scavenging and anti-inflammatory effects (45) (Table 3 & Fig. 2a, b).

### Pharmacological activity

The therapeutic potential of the various solvent extracts of *T. procumbens* has been thoroughly investigated for their antifungal, antibacterial, immunomodulatory, wound healing, anticancer, antioxidant, larvicidal, anticoagulating, homeostatic activities and more (Fig. 3). This review discuss the pharmacological properties of *T. procumbens* extracts and their metabolites both *in vitro* and *in vivo* (Table 4 & Fig. 3).



### Anti-bacterial activity


**Table 3.** List of different phytochemicals detected in *Tridax procumbens* and their activity

Sl. No.	Phytochemical	Compounds	Mol. formula	Pharmacological Activity	References
1.	Alkaloid	Tryptanthrin	$C_{15}H_{18}N_2O_2$	Anticancer, antimicrobial, anti-inflammatory, antioxidant.	(46, 47)
		Betulinic Acid	$C_{30}H_{48}O_3$	Anticancer, anti-inflammatory, hepatoprotective, antioxidant	(48, 49)
		Stigmasterol	$C_{29}H_{48}O$	Anti-inflammatory, hypocholesterolemic, antidiabetic, antioxidant, anticancer.	(50, 51)
		Quercetin	$C_{15}H_{10}O_7$	Potent antioxidant, anti-inflammatory, anticancer, cardioprotective, neuroprotective, antihypertensive.	(52, 53)
2.	Flavonoids	Luteolin	$C_{15}H_{10}O_6$	Antioxidant, anti-inflammatory, anticancer, neuroprotective, analgesic/antiallergen	(54, 55)
		Catechin	$C_{15}H_{14}O_6$	Antioxidant/free-radical scavenger, cardioprotective, anti-inflammatory, antimicrobial, anti-cancer	(56, 57)
		Kaempferol	$C_{15}H_{10}O_6$	Antioxidant, anti-inflammatory, anticancer, neuroprotective, hepatoprotective, antidiabetic	(58, 59)
3.	Tannins	Apigenin	$C_{15}H_{10}O_5$	Antioxidant, anti-inflammatory, anticancer, neuroprotective, anxiolytic/sedative properties	(60, 61)
		Ellagic acid	$C_{14}H_6O_8$	Antioxidant, anti-inflammatory, antimutagenic/anticancer, hepatoprotective, cardioprotective, neuroprotective.	(37, 62)
4.	Terpenoids	Gallic acid	$C_7H_6O_5$	Antioxidant, anti-inflammatory, antimicrobial, anticancer, neuroprotective, hepatoprotective.	(63, 64)
		Eugenol	$C_{10}H_{12}O_2$	Antioxidant, anti-inflammatory	(65)
5.	Saponins	Disogenin	$C_{27}H_{42}O_3$	Anticancer/antiproliferative, anti-inflammatory, hypolipidemic, antidiabetic, osteoprotective, cardioprotective.	(66, 67)
		Oleanolic acid	$C_{30}H_{48}O_3$	Hepatoprotective, anti-inflammatory, antioxidant, anticancer, antiviral, cardioprotective, hypoglycemic	(68, 69)
		Hederagenin	$C_{30}H_{48}O_4$	Anti-inflammatory, antimicrobial, anticancer, neuroprotective, metabolic benefits	(70)
		Campesterol	$C_{28}H_{48}O$	Phytosterol- cholesterol-lowering (reduces intestinal cholesterol uptake), anti-inflammatory, antioxidant, potential chemopreventive effects	(71)
		Caffeic acid	$C_9H_8O_4$	Antioxidant, anti-inflammatory, anticancer (modulates signalling), antiviral/antimicrobial, neuro- and hepatoprotective.	(72, 73)
6.	Phenolic Compound	Chlorogenic acid	$C_{16}H_{18}O_9$	Antioxidant, anti-inflammatory, hepatoprotective, cardioprotective, antidiabetic (glucose-metabolism modulating), neuroprotective, antimicrobial	(74, 75)
		Ferulic acid	$C_{10}H_{10}O_4$	Antioxidant, anti-inflammatory, neuroprotective, cardioprotective/antithrombotic, anticancer (chemo-preventive), skin-protective	(37, 76)

**Table 4.** Different phytochemicals in different parts of *Tridax procumbens* and their biological activity

Activity	Part used	Solvent	Phytochemical	Reference
Anti-inflammatory	Aerial parts	Ethanol	Cent aureidin, Centaurein and Bergenin Luteolin-4'-O-B-D-glucopyranoside	(77)
Anti-bacterial	Aerial part	Ethanol	3-O-methylquercetin-4'-O-β-D-glucopyranoside 3-O-methylquercetin-4'-O-β-D-glucopyranoside	(78, 79)
Anti-fungal			5,8,3-dihydroxy-3,7,4-trimethoxy-6-O-β-D-glucopyranosyl flavones, 6,8,3-trihydroxy-3,7,4-trimethoxyflavone, puerarin, esculetin, oleanolic acid, betulinic acid, centaurein, bergenin and centaureidin	(34)
Anti-cancer	Aerial part	Ethanol	(3S,5R,6S,7E)-5,6 epoxy 3-hydroxy-7-megastigmene-9-one Icaricide B1	(78, 80)
Anti-leishmanial	Leaves	Methanol	(3S)-16-17-Didehydrofalcarinol or Oxylinpin R= Gu 8,3'-dihydroxy-3,7,4'-trimethoxy-6-O-B-D glucopyranosyl flavone	(78, 81)
Anti-oxidant	Aerial parts	Ethanol	Quercetangatin-3,6,4'-trihydroxy-7-O-neohesperidoside 3-O-methylquercetin-4'-O-B-D-glucopyranoside Quercetangatin-3,6,4'-trihydroxy-7-O-β-D-glucopyranoside	(78)
Wound healing			Caffeic acid, Epicatechin, Kaempferol and Tannic acid,	(72)

**Fig. 2a.** Different secondary metabolites present in *Tridax procumbens* extracts.**Fig. 2b.** Elucidation of secondary metabolites from *Tridax procumbens*.



**Fig. 3.** Pharmacological activity of *Tridax procumbens* extracts.

The antibacterial assay of *T. procumbens*' various solvent extracts against different bacterial species was tested using disc diffusion and well diffusion methods. Disc diffusion technique revealed that *T. procumbens* methanolic extract showed a zone of inhibition (ZOI) in *Escherichia coli*. However, the ethyl acetate extract demonstrated a larger ZOI in *Bacillus cereus*, *Staphylococcus aureus* and *Salmonella typhi* (82). For *Pseudomonas vulgaris*, ethyl alcohol extract of *T. procumbens* leaf is the most potent antibacterial agent (9). Tannins and flavonoids with multiple mechanisms of action, including inhibition of cytoplasmic membrane function, energy metabolism and DNA gyrase, further ethanolic extract demonstrated antibacterial activity against multidrug-resistant *Pseudomonas* from nosocomial infections (33, 83, 84).

*Tridax procumbens* ethanolic extract displayed antibacterial action on nosocomial strain of *Pseudomonas aeruginosa* from pneumonia patient secretions, including bronchoalveolar lavage and tracheal secretions (33). Further, the methanolic extract of *T. procumbens* flower was found to inhibit *P. aeruginosa*, *E. coli* and *B. cereus*. In contrast, the methanolic extract of the leaf demonstrated the maximum zone against *Pseudomonas fluorescens* (85). In addition, the extract of *T. procumbens* leaf with methanol, ethanol and ethyl acetate against *Bacillus subtilis*, *Enterobacter aerogenes*, *S. aureus* and *E. coli* showed good activity (86).

Four bacterial strains, *Proteus mirabilis*, *B. subtilis*, *S. aureus* and *E. coli* were tested against various *T. procumbens* extracts. The ethyl acetate extract had the most excellent potent repressing effects on all the tested strains of bacteria. Meanwhile, hexane extract inhibits *S. aureus* and *E. coli*, while chloroform extract only inhibits *S. aureus* (87). Ethanolic extract of *T. procumbens* at 200 mg/mL demonstrated broad-spectrum activities on every tested bacterium; *E. coli* had the least inhibitory effect, while *S. aureus* had the maximum inhibition (88).

Petroleum ether, hexane extracts of *T. procumbens* leaf, flower and stem showed antibacterial efficacy against gram-

negative (*E. aerogenes*) and gram-positive (*S. aureus* and *B. subtilis*) bacteria. However, the flower's hexane extract showed potent antibacterial activity against *E. aerogenes*, with a 20 mm zone of inhibition. The test bacteria were moderately inhibited by the leaf, followed by the stem. Petroleum ether had a moderate effect on the test bacteria, while chloroform extracts were found to be the least active (86). Furthermore, aqueous extract of *T. procumbens* was used to test the antibacterial assay against *E. coli*, *Enterococcus faecalis*, *B. subtilis* and *P. aeruginosa* (89). Studies on *T. procumbens* different plant parts with different solvents demonstrated that extracts have rich active phytochemicals that can suppress the microbes by static and cidal action (Table 4 & Fig. 3).

#### Anti-fungal activity

The *T. procumbens* can control fungal growth from the presence of phenols, flavonoids, saponins, sterols and fatty acids (9, 90). Significant inhibitory action against *Candida albicans* was demonstrated by *T. procumbens* methanol extract from various leaves, stems, flower and roots, showing 8-13 mm inhibition. Additionally, root methanol extract demonstrated intense action against *Candida tropicalis* and *Candida glabrata*, indicating that *T. procumbens* bioactive components could be employed as potent anti-candidal medications (91).

The *T. procumbens* showed activity against both *Aspergillus turcosus* and *Aspergillus parasiticus*. Significant inhibition zone was observed in ethanol, aqueous, chloroform and methanol extracts at different concentration range between 1 to 5 mg/mL; however, the highest inhibition was observed in aqueous and chloroform extracts (20 mm at 5 mg/mL) against *A. parasiticus* (92). *Curvularia eragrostidis* was less effective than *Bipolaris sorokiniana* when treated with *T. procumbens* aqueous alcohol extract (93). Further, *T. procumbens* ethanolic extract showed maximum antifungal activity towards *A. flavus* and *C. albicans* (94).

The *T. procumbens* methanolic extract on *A. fumigatus*, *A. niger*, *C. albicans* inhibition range was 9-15 mm (82). Additionally, in comparison to ciprofloxacin, *T. procumbens* methanolic extracts

demonstrated efficient suppression against *A. niger* and *A. ocraceus* (95). Furthermore, *T. procumbens* methanolic leaf extract at 0.1 % and 0.01 % concentrations was tested against *Helminthosporium oryzae*, *Rhizoctonia solani* and *Pyricularia oryzae* at 0.1 % and 0.01 % concentrations; the comparative magnitude of percent inhibition was found to be higher in *H. oryzae* and *R. solani* than in the other two pathogens over their respective controls (96).

#### Anti-hepatotoxic or hepatoprotective activity

An antihepatotoxic agent protects the liver from damage and can prevent or reduce the damaging effects of toxins and diseases on the liver. The alcoholic extract of *T. procumbens* triggers the cholinergic receptors, which protect the liver through the vagus nerve; furthermore, this extract can repair damage from free radicals and increase the activity of the liver antioxidant defence mechanism (9). This plant is used to make "Bhringraj," a well-known ayurvedic medication for liver problems (9). In the rat model of hepatitis induced by D-galactosamine/lipopolysaccharide (D-GalN/LPS), the ethanolic extract of *Tridax procumbens* significantly elevated the levels of alanine transaminase, aspartate transaminase, lactate dehydrogenase, alkaline phosphatase and bilirubin, restoring them toward normal values. (7). Furthermore, *T. procumbens* ethanolic extract showed hepatoprotective effect against CCl<sub>4</sub>, restoring the liver to its normal look and resembling the normal liver cells observed in rats (97).

Various dosages of *T. procumbens* ethanolic extract administered orally demonstrated the antioxidative and hepatoprotective effectiveness against liver damage caused by paracetamol (9). Additionally, male Wistar albino rats were protected against hepatotoxicity induced by D-galactosamine by methanol, chloroform and water extracts of *T. procumbens* flowers; the methanolic extract showed the highest efficacy because of its high phenolic contents. However, aqueous extract leaves have antioxidant properties, demonstrating hepatoprotective effects in rats (9, 28). Besides, the aerial parts of *T. procumbens* showed hepatoprotective effects and essential protection in alleviating D-galactosamine/lipopolysaccharide (98).

#### Anti-inflammatory activity

Numerous investigations have documented the anti-inflammatory properties of various *T. procumbens* extracts attributed to their active chemical components (78). Anti-inflammatory properties of *T. procumbens* extracts derived from numerous parts of the plant are well documented due to their active chemical ingredients. Anti-inflammatory activity, induced by carrageenan, correlated with diclofenac sodium in various experiments (78). *In vivo*, different authors have observed the anti-inflammatory effects of *T. procumbens* methanol extract (78, 99). *Tridax* leaves aqueous extract exhibited contractile action, in contrast to the control group. The aqueous *tridax procumbens* leaf extract (ATPLE)-treated group exhibited considerably greater contraction in response to Ca<sup>2+</sup> activated K<sup>+</sup> channel (KCl), H1 receptor subtype activation and the M3 receptor acetylcholine, which relates to Gq and PKC (9). *In vivo* anti-inflammatory activity of methanol extract of *T. procumbens* leaves was shown in xylene-induced ear, carrageenan-induced paw, egg albumin and formalin-induced inflammation, in mice at 25–100 µg/kg. The authors observed good anti-inflammatory effects in mice BALB/c (20–30g) (99). Various animal studies have shown that several extracts of *T. procumbens* have anti-inflammatory properties.

#### Anti-cancer activity

Cancer is one of the most prevalent diseases that can start in almost any organ or tissue of the body when abnormal cells grow uncontrollably. Treatments can include surgery, radiation therapy, chemotherapy, targeted therapy, immunotherapy and hormone therapy. However, naturally derived compounds from medicinal plants and their properties make them targets for potential anticancer treatments (100). In traditional medicine, *T. procumbens* has been used to treat various illnesses. However, there is no evidence regarding its historical use for cancer; nevertheless, recently, several new publications suggest that *T. procumbens* contains many potent anticancer compounds, including flavonoids, tannins and saponins (78). Using the 9(3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay (MTT assay), anticancer activity of 5,6-epoxy-3-hydroxy-7-megastigmene-9-one was examined in leukemia (HL-60), gastric cells (SGC-7901) (58). The MTT test was used to determine the cytotoxicity of chemicals obtained from plants against human lung cancer cells (35).

*In vitro* MTT assay was used to assess the essential oil extracted from *T. procumbens* leaves for anticancer activity in the MCF-7 cell line. The results displayed that the essential oil had a strong anticancer effect, which could be explained by the occurrence of crucial terpenes like α-pinene and β-pinene (9). *Tridax procumbens* essential oil demonstrated excellent cytotoxicity, killing 70.28 % of cancer cells in 24 hr at a dose of 50 µg. This indicates that the oil effectively stops lung metastases of the B16F-10 cells in mice (78). Anticancer action by acetone and water extracts of *T. procumbens* flowers was examined. Contrary to the water extract (6.6 % cell death), the acetone extract displayed intense anticancer action against PC3 cell lines using the MTT test at a dose of 250 µg/mL (82.28 % cell death) (101).

The *T. procumbens* leaf acetone and aqueous extract demonstrated anticancer efficacy in PC3 cancer cells *in vitro* using the MTT and trypan blue exclusion assays. While the aqueous extract showed little anticancer effect, the acetone extract had intense anticancer action, resulting in 93 % cell death (78). *Tridax procumbens* leaves ethanol, acetone extracts (10–50 µg/mL) revealed anticancer action in lung cancer cells A-549, while water extract exhibited no anticancer activity (101). Further, the cytotoxicity impact of *T. procumbens* leaf methanol extract at 250 µg/mL was assessed in human lung cancer A-549 cells at 84 ± 2.8 % and human breast MCF-7 cells at 68 ± 3.1 % (86). Different cancer cell studies have demonstrated that the active chemicals in medicinal plants can prevent the proliferation of cancer cells.

#### Anti-diabetic activity

Diabetes mellitus is a chronic metabolic disorder characterised by hyperglycemia due to insufficient insulin production, impaired insulin action or both. It is a global health concern affecting millions of people and it's associated with serious complications. Synthetic drugs help lower blood sugar levels and improve insulin sensitivity but may also have side effects. Hence, herbal and natural anti-diabetics-based treatment is gaining interest in integrative medicine.

Ethanolic extract of *T. procumbens* showed antidiabetic action in Wistar rats induced by streptozotocin-nicotinamide (30). On the other hand, methanolic leaf and flower extract of *T. procumbens* revealed effective antidiabetic activity (85). Besides, *T. procumbens* leaf aqueous and alcoholic extracts dramatically lowered blood glucose levels in rats with alloxan-induced diabetes (98). Water,

chloroform, alcoholic and petroleum ether extracts of *T. procumbens* leaves were administered orally for seven days at 200 mg/kg. The results showed that these extracts significantly reduced blood glucose in rats with alloxan-induced diabetes, but petroleum extract showed very little antidiabetic activity (16).

The *T. procumbens* extract was extracted using 50 % methanol and tested for acute and sub chronic antihyperglycemic effects in alloxan-induced diabetic rats. Rats with diabetes revealed a decrease in fasting blood glucose when oral *T. procumbens* extract was administered at acute and sub chronic levels (250, 500 mg/kg b.wt.), but not in untreated rats (102). The antihyperglycemic effects were dose-independent, Oral Glucose Tolerance Test (OGTT) and body weight-maintained by antihyperglycemic action (102). Diabetics' chronic disease is inadequately controlled by insulin; hence, various plant extracts have shown a reduction in blood glucose levels.

### Immunomodulatory activity

The immune system is critical in maintaining physiological homeostasis by defending the body against pathogens, removing damaged cells and controlling inflammation. Immunomodulatory activity refers to the ability of a substance to modify, regulate or normalise the immune response. This modulation can either enhance (immunostimulation) or suppress (immunosuppression) specific components of the immune system, depending on the therapeutic need (103).

Albino rats orogastrically dosed with *P. aeruginosa* and treated with ethanolic extracts of *T. procumbens* leaves have an immunomodulatory impact (98, 104). In animal models, plant leaf ethanol decoctions have been found to have immunomodulatory effects. Additionally, an increase in hemagglutination antibody and immunological response stimulation were observed (105). The humoral immune response was stimulated by the ethanol-insoluble fraction of *T. procumbens* extract, which also markedly enhanced leukocyte count, phagocytic index and splenic antibody-secreting cells (35). Further, *T. procumbens* ethanol-insoluble water extract revealed IP administration of TPEIF (0.25, 0.5 g/kg body weight) showed increased leucocyte count and splenic antibody-secreting cells (56). Similarly, ethanol-insoluble water extract has been shown to significantly raise phagocytic index and leukocyte count (84).

### Anti-oxidant activity

2, 2-Diphenyl-1-Picrylhydrazyl (DPPH) and a vitamin as a positive control in the *in vitro* antioxidant assay of *Tridax* were assessed against three isolated flavonoids: 6-4'-trihydroxy-7-O- $\beta$ -D-glucopyranoside, 3-O-methyl quercetin-4'-O- $\beta$ -D-glucopyranoside, Quercetangatin-3. The antioxidant capacity of compounds 3-O-methyl quercetin-4'-O- $\beta$ -D-glucopyranoside and 6-O-trihydroxy-7-O-neohesperidoside was restricted, with  $IC_{20} > 100 \mu\text{mol/L}$ , whereas compound 6-O-trihydroxy-7-O- $\beta$ -D-glucopyranoside exhibited little activity, with  $IC_{20} > 100 \mu\text{mol/L}$  (78, 80).

The *T. procumbens* has a high concentration of flavonoids, phenols, anthraquinones, vitamins A and C and carotenoids, which account for its intense antioxidant action (6, 106). Using the DPPH activity, *T. procumbens* essential oils have demonstrated antioxidant activity by lowering oxidative stress levels (6). *Tridax procumbens* leaves aqueous and ethanol extracts were examined using DPPH antioxidant assay. At a 200  $\mu\text{g/mL}$  dose, the ethanol extract demonstrated better antioxidant activity (86). Another study assessed the antioxidant assay of two compounds of *T. procumbens*,

6,8,3'-trihydroxy-3,7,4'-trimethoxy flavone and 3'-dihydroxy-3,7,4'-trimethoxy-6-O- $\beta$ -D-glucopyranosyl flavone. Comparing these two compounds to the standard control, their  $IC_{50}$  values were 0.03180 and 0.04618 mg/mL, respectively, indicating strong antioxidant activity,  $IC_{50}$  for Trolox = 0.00948 mg/mL (77).

### Anti-parasitic activity

*In vitro*, the methanolic extract of *T. procumbens* exhibited 50 % inhibition at 3  $\mu\text{g/mL}$  and reduced promastigote growth of *Leishmania mexicana*, which causes leishmaniasis illness (9, 81). Furthermore, *in vivo* experiments showed that *T. procumbens* has significant activity in *L. mexicana* and is a candidate for leishmanicidal activity. In *T. procumbens*, oxylipin, isocordoin, 20,40-dihydroxy-30-(g,g-dimethylallyl)-dihydrochalcone, cholestra-4,20,24-trien-3-one or pentalinosterol and 24-methylcholesta-4-24(28)-dien-3-one, cholest-4-en-3-one, 6,7-dihydroneeridenone, neridenone and cholest-5,20,24-trien-3b-ol, metabolites showed an  $IC_{50}$  at 30 mg/mL against *L. mexicana* (107).

Water and ethanol decoctions of *T. procumbens* offer anti-plasmodial qualities that combat *Plasmodium falciparum*, which is resistant to chloroquine (105). *Tridax procumbens* demonstrated leishmanial action when crude extracts of the entire plant were used (60). Red blood cells were shielded from *P. falciparum* damaged by *T. procumbens* aqueous and ethanol extract, demonstrated by the tetrazolium-based colourimetric technique (108). The antiplasmodial properties of water, ethyl acetate, ethanol and chloroform extracts of *T. procumbens* flowers, leaves and stems were examined in Ghana (6). The water and ethanol infusions exhibit little toxicity to human red blood cells and possess anti-plasmodial qualities against the chloroquine-resistant parasite *P. falciparum* (35). In addition, *Tridax*, entire plant in methanol extracts were evaluated for leishmanicidal assay against promastigotes of *L. mexicana* in an *in vitro* assay (56). Thus, active phytochemicals can control parasitic growth and proliferation.

### Insect repellent activity

Repellents act locally or at a distance, deterring an arthropod from flying to, landing on or biting human or animal skin (109). Insect repellents act primarily by interfering with the insect's olfactory or sensory receptors, thereby modifying their behaviour and keeping them away from the treated surface, human, animal or plant (110). The *T. procumbens* leaves produce some essential oils, which act as a source of insect repellent (35). *Tridax procumbens* contains a significant chemical constituent,  $\alpha$ -terpinene,  $\alpha$ -terpineol and  $\beta$ -pinene, which provides a repellency against *A. stephensi* (12). The plant essential oils do not cause skin irritation, hot sensations or rashes on the arms and can be used directly (9). In mosquito cages, essential oils extracted from leaves by condensation, their ability to locally resist the malaria parasite *A. stephensi* was examined (105). *Tridax procumbens* leaf extracts in hexane, chloroform, ethyl acetate and acetone have adulticidal activity against *A. stephensi* (111). On the contrary, chloroform and acetone extracts of *T. procumbens* have the maximum adulticidal activity (78, 112).

### Hemostatic activity

*In vitro*, the hemostatic properties of the plant leaves in various solvent extracts were determined by employing Lee-White's method (105). When the hemostatic activity of several extracts from *T. procumbens* leaves was examined, the ethanol extract consistently shortened the clotting time in the blood samples (16). Further, *T. procumbens* leaves water extract showed significant

blood coagulation activity; for those reasons, it may be used as a potent hemostatic agent (9). Plant-based flavonoids, which stimulate osteoblasts and inhibit osteoclasts, can control bone homeostasis. *Tridax procumbens* fraction (TPF) can potentially treat disorders such as osteoporosis associated with bone loss by blocking receptor activator of nuclear factor- $\kappa$ B ligand (RANKL)-induced osteoclast growth and pit formation in primary osteoclastic cells. In mice, TPF boosted bone formation indices and inhibited the growth of osteoclasts (35).

### Wound healing activity

Studies have shown that *T. procumbens* ethanolic leaf extracts can heal wounds in diabetic and non-diabetic animals. Animals treated with 5% and 2.5% (w/w) extract showed a noticeable difference in wound index, epithelization period and wound contraction (113). In experimental male Wistar rats, water extract of *T. procumbens* leaves accelerated healing and overcame steroid-depressed healing (56). *Tridax procumbens* has been used traditionally to promote wound healing and has strong anti-inflammatory, anti-microbial and anticancer effects (114). Mice treated with *T. procumbens* leaf juice at a 1 mg/g concentration exhibited increased vascularity, fibroblast number, collagen content, re-epithelization and wound contraction parameters. Additionally, it was noted that high dosages cause inflammation, oedematous tissue and vascularity and it was determined that this plant extract's pro-healing properties depend on dosage (115). In a different study, mice with excision wounds showed improved results after 15 days using an ointment formulation containing hydroalcoholic leaf extract of *T. procumbens* instead of the blank ointment model.

### Conclusion

In conclusion, this review highlights *in vitro* and *in vivo* the significant therapeutic potential of *T. procumbens* which is traditionally used for treating ailments such as anti-diabetic, anti-ulcer, anti-cancer etc. *Tridax procumbens* is one of India's most widely recognized herbal remedies. It contains diverse bioactive compounds, including alkaloids, phenols, flavonoids and terpenoids, that have shown promising pharmacological activities such as antioxidant, anti-inflammatory, antimicrobial and more. Despite its traditional uses and known benefits, further research must identify and characterize the specific bioactive compounds responsible for these effects and mechanism. Advanced extraction methods and compound profiling are essential to better understanding the plant's therapeutic mechanisms. This review highlights the historical significance and the need for continued investigation into *T. procumbens*, as it holds promise for uncovering additional pharmacological properties yet to be fully explored to bridge the gap. Future studies targeting the isolation and functional analysis of its active constituents may contribute significantly to developing novel, plant-based pharmaceuticals and pave the way to creating more advanced conventional medications.

### Acknowledgements

The authors thank the Head, Department of Botany, Tripura University, for providing laboratory facilities. We thank Mr. Rajdeep Dhar, Research scholar, for helping to make phytochemical structures.

### Authors' contributions

HK carried out the literature collection and drafted the manuscript. AB, KB, KM and GB participated and carried out the table and figure preparation, NB conceived the idea and critical review of the manuscript. JKS and GS participated and improved the manuscript. All authors read and approved the final manuscript.

### Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interest to declare.

**Ethical issues:** None

### References

1. Chaachouay N, Zidane L. Plant-derived natural products: a source for drug discovery and development. *Drugs Drug Candidates*. 2024;3(1):184-207. <https://doi.org/10.3390/ddc3010011>
2. World Health Organization. WHO global report on traditional and complementary medicine. Geneva: World Health Organization; 2019.
3. Heinrich M, Ankli A, Frei B, Weimann C, Sticher O. Medicinal plants in Mexico: healers consensus and cultural importance. *Soc Sci Med*. 1998;47:1859-87. [https://doi.org/10.1016/S0277-9536\(98\)00181-6](https://doi.org/10.1016/S0277-9536(98)00181-6)
4. Alonso-Castro AJ, Villarreal ML, Salazar-Olivo LA, Gomez-Sanchez M, Dominguez F, Garcia-Carranca A. Mexican medicinal plants used for cancer treatment: pharmacological, phytochemical and ethnobotanical studies. *J Ethnopharmacol*. 2011;133(3):945-72. <https://doi.org/10.1016/j.jep.2010.11.055>
5. Rhun E, Devos P, Bourg V, Darlix A, Lorgis V, Ahle G, et al. Complementary and alternative medicine use in glioma patients in France. *J Neurooncol*. 2019;145(3):487-99. <https://doi.org/10.1007/s11060-019-03315-8>
6. Beck S, Mathison H, Todorov T, Calderón-Juárez EA, Kopp O. A review of medicinal uses and pharmacological activities of *Tridax procumbens* (L.). *J Plant Stud*. 2018;7(1):19-35. <https://doi.org/10.5539/jps.v7n1p19>
7. Ravikumar V, Shivashangari KS, Devaki T. Hepatoprotective activity of *Tridax procumbens* against d-galactosamine/lipopolysaccharide-induced hepatitis in rats. *J Ethnopharmacol*. 2005;101(1-3):55-60. <https://doi.org/10.1016/j.jep.2005.03.019>
8. Kethamakka SRP, Deogade MS. Jayanti veda (*Tridax procumbens*)—unnoticed medicinal plant by Ayurveda. *J Ind Syst Med*. 2014;2(1):6-22. <https://doi.org/10.5005/JISM-11051-02103>
9. Dattaray D. Traditional uses and pharmacology of plant *Tridax procumbens*: a review. *Syst Rev Pharm*. 2022;13(5):511-7.
10. Karimi A, Majlesi M, Rafieian-Kopaei M. Herbal versus synthetic drugs; beliefs and facts. *J Nephropharmacol*. 2015;4(1):27.
11. Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. <https://powo.science.kew.org/>
12. Gubbiveeranna V, Nagaraju S. Ethnomedicinal, phytochemical constituents and pharmacological activities of *Tridax procumbens*: a review. *Int J Pharm Sci*. 2016;8(2):1-7.
13. Shabana HA, Gairola S, Mahmoud T. *Tridax procumbens* L. (Asterales: Asteraceae), a new record to the flora of the United Arab Emirates. *Biodivers J*. 2020;11(4):889-96. <https://doi.org/10.31396/Biodiv.Jour.2020.11.4.889.896>
14. Nalage AA, Kambale HV, Satpute VM, Waghmare SA. Pharmacology of *Tridax procumbens*. *Int J Novel Res Dev*. 2023;8(3):a751-60.
15. Poll E. Medicinal and aromatic plants of Guatemala and the need for their conservation. In: *Proc III WOCMAP Congr Med Aromat Plants*. 2003;167-70.

16. Bhagwat DA, Killedar SG, Adnaik RS. Anti-diabetic activity of leaf extract of *Tridax procumbens*. *Int J Green Pharm*. 2008;2(2):125-8. <https://doi.org/10.4103/0973-8258.41188>
17. Sureshkumar J, Silambarasan R, Ayyanar M. An ethnopharmacological analysis of medicinal plants used by the Adivyan community in Wayanad district of Kerala, India. *Eur J Integr Med*. 2017;12:60-73. <https://doi.org/10.1016/j.eujim.2017.04.006>
18. Gamboa-Leon R, Vera-Ku M, Peraza-Sanchez SR, Ku-Chulim C, Horta-Baas A, Rosado-Vallado M. Antileishmanial activity of a mixture of *Tridax procumbens* and *Allium sativum* in mice. *Parasite*. 2014;21:15. <https://doi.org/10.1051/parasite/2014016>
19. Torres RD, Gamit AM, Peria JT. Formulation and evaluation of coat button (*Tridax procumbens*; Asteraceae) leaf extract simple syrup for pain management. *Int J Res Public Rev*. 2025;6(7):401-12.
20. Agyare C, Boakyee YD, Bekoe EO, Hensel A, Dapaah SO, Appiah T. African medicinal plants with wound healing properties. *J Ethnopharmacol*. 2016;177:85-100. <https://doi.org/10.1016/j.jep.2015.11.008>
21. Ingle NA, Dubey HV, Kaur N, Gupta R. *Tridax procumbens*: a multiuseful weed—a review. *J Adv Oral Res*. 2014;5(1):14-6. <https://doi.org/10.1177/2229411220140104>
22. Shaikh JR, Patil M. Qualitative tests for preliminary phytochemical screening: an overview. *Int J Chem Stud*. 2020;8(2):603-8. <https://doi.org/10.22271/chemi.2020.v8.i2i.8834>
23. NdezoBisso B, NjikangEpieNkwelle R, TchuenguemTchuenteu R, Dzoyem JP. Phytochemical screening, antioxidant and antimicrobial activities of seven under investigated medicinal plants against microbial pathogens. *Adv Pharmacol Pharm Sci*. 2022;2022:1998808. <https://doi.org/10.1155/2022/1998808>
24. Adil M, Filimban FZ, Ambrin, Quddoos A, Sher AA, Naseer M. Phytochemical screening, HPLC analysis, antimicrobial and antioxidant effect of *Euphorbia parviflora* L. (Euphorbiaceae Juss.). *Sci Rep*. 2024;14(1):5627. <https://doi.org/10.1038/s41598-024-55905-w>
25. Ezeonu CS, Ejikeme CM. Qualitative and quantitative determination of phytochemical contents of indigenous Nigerian softwoods. *New J Sci*. 2016;2016:5601327. <https://doi.org/10.1155/2016/5601327>
26. Ikewuchi JC, Ikewuchi CC. Comparative study of the mineral element composition of some common Nigerian medicinal plants. *Pac J Sci Technol*. 2009;10(1):362-6.
27. Ujowundu CO, Igwe CU, Enemor VHA, Nwaogu LA, Okafor OE. Nutritive and anti-nutritive properties of *Boerhavia diffusa* and *Commelina nudiflora* leaves. *Pak J Nutr*. 2008;7(1):90-2. <https://doi.org/10.3923/pjn.2008.90.92>
28. Dhanabalan R. In vitro phytochemical screening and antibacterial activity of aqueous and methanolic leaf extracts of *Tridax procumbens* against bovine mastitis isolated *Staphylococcus aureus*. *Ethnobot Leaflets*. 2008;12:1090-5.
29. Petchi RR, Vijaya C, Parasuraman S. Antidiabetic activity of polyherbal formulation in streptozotocin–nicotinamide induced diabetic Wistar rats. *J Tradit Complement Med*. 2014;4(2):108-17. <https://doi.org/10.4103/2225-4110.126174>
30. Christudas S, Kulathivel TM, Agastian P. Phytochemical and antibacterial studies of leaves of *Tridax procumbens* L. *Asian Pac J Trop Biomed*. 2012;2(1):S159-61. [https://doi.org/10.1016/S2221-1691\(12\)60149-X](https://doi.org/10.1016/S2221-1691(12)60149-X)
31. Singh P, Jain K, Khare S, Shrivastav P. Evaluation of phytochemical and antioxidant activity of *Tridax procumbens* extract. *Pharm Biosci J*. 2017;5:41-7. <https://doi.org/10.20510/ukjpb/5/i6/166569>
32. Pai C, Kulkarni U, Borde M, Murali S, Mrudula P, Deshmukh Y. Antibacterial activity of *Tridax procumbens* with special reference to nosocomial pathogens. *Br J Pharm Res*. 2011;1(4):164-73. <https://doi.org/10.9734/BJPR/2011/763>
33. Policegoudra RS, Chattopadhyay P, Aradhya SM, Shivaswamy R, Singh L, Veer V. Inhibitory effect of *Tridax procumbens* against human skin pathogens. *J Herb Med*. 2014;4(2):83-8. <https://doi.org/10.1016/j.hermed.2014.01.004>
34. Pinjari R, Vaishnav S, Gangurde AB, Shevale P, Khyati S, Bairagi V. A review of *Tridax procumbens*: medicinal uses, pharmacological activity and phytochemical screening. *Int J Res Anal Rev*. 2024;11(3):1-10.
35. Ali M, Ravinder E, Ramachandram R. A new flavonoid from the aerial parts of *Tridax procumbens*. *Fitoterapia*. 2001;72(3):313-5. [https://doi.org/10.1016/S0367-326X\(00\)00296-3](https://doi.org/10.1016/S0367-326X(00)00296-3)
36. Ikewuchi JC, Ikewuchi CC, Ifeanacho MO. Phytochemical composition of *Tridax procumbens* Linn. leaves: potential as a functional food. *Food Nutr Sci*. 2015;6:992-1004. <https://doi.org/10.4236/fns.2015.611103>
37. Agme VN, Agme RN. Phytochemical analysis of *Tridax procumbens* L. *Am Int J Res Formal Appl Nat Sci*. 2022;6(2):95-7.
38. Petchi RR, Vijaya C, Parasuraman S. Anti-arthritis activity of ethanolic extract of *Tridax procumbens* (Linn.) in Sprague Dawley rats. *Pharmacogn Res*. 2013;5(2):113-7. <https://doi.org/10.4103/0974-8490.110541>
39. Ghuge SG, Gawale VG, Patil SP. A review on pharmacological activities of *Tridax procumbens*. *Int J Trend Sci Res Dev*. 2024;8(6):839-55.
40. Dhandapani R, Sabna B. Phytochemical constituents of some Indian medicinal plants. *Anc Sci Life*. 2008;27(1):1-8.
41. Pandey A, Tripathi S. A review on pharmacognosy, pre-phytochemistry and pharmacological analysis of *Tridax procumbens*. *Pharma Tutor*. 2014;2(4):78-86.
42. Chaudhari HC, Patil KP. A review on medicinal importance of *Tridax procumbens* Linn. *Res Rev Pharm Pharm Sci*. 2022.
43. Kushwaha P, Yadav SS, Singh V, Dwivedi LK. Phytochemical screening and GC-MS studies of the methanolic extract of *Tridax procumbens*. *Int J Pharm Sci Res*. 2019;10(5):2492-6.
44. Saxena M, Mir AH, Sharma M, Malla MY, Qureshi S, Mir MI, et al. Phytochemical screening and in vitro antioxidant activity of isolated bioactive compounds from *Tridax procumbens* Linn. *Pak J Biol Sci*. 2013;16(24):1971-7. <https://doi.org/10.3923/pjbs.2013.1971.1977>
45. Kaur M, Chopra DS. Green synthesis of iron nanoparticles for biomedical applications. *Glob J Nanomed*. 2018;4(4):68-76.
46. Zhou X. Recent advances of tryptanthrin and its derivatives as potential anticancer agents. *RSC Med Chem*. 2024;15(4):1127-47. <https://doi.org/10.1039/D3MD00698K>
47. Xu R, Zhang J, Yuan K. Two new flavones from *Tridax procumbens* Linn. *Molecules*. 2010;15(9):6357-64. <https://doi.org/10.3390/molecules15096357>
48. Lou H, Li H, Zhang S, Lu H, Chen Q. A review on preparation of betulinic acid and its biological activities. *Molecules*. 2021;26(18):5583. <https://doi.org/10.3390/molecules26185583>
49. Kale MA, Dhake AS. Anti-inflammatory activity of phytosterols and their derivatives isolated from the leaves of *Tridax procumbens* Linn. *J Curr Pharm Res*. 2013;4(1):1049-54. <https://doi.org/10.33786/JCPR.2013.v04i01.003>
50. Bakrim S, Benkhaira N, Bourais I, Benali T, Lee LH, El Omari N, et al. Health benefits and pharmacological properties of stigmasterol. *Antioxidants*. 2022;11(10):1912. <https://doi.org/10.3390/antiox11101912>
51. Pande PS, Mishra MN. Isolation and identification of quercetin and rutin from leaves of *Tridax procumbens* Linn. by HPLC analysis. *Int J Chem Phys Sci*. 2015;4:65-8.
52. Mirza MA, Mahmood S, Hilles AR, Ali A, Khan MZ, Zaidi SAA, et al. Quercetin as a therapeutic product: evaluation of its pharmacological action and clinical applications—a review. *Pharmaceuticals*. 2023;16(11):1631. <https://doi.org/10.3390/ph16111631>

53. Lakhera S, Rana M, Devlal K, Celik I, Yadav R. A comprehensive exploration of pharmacological properties, bioactivities and inhibitory potentiality of luteolin from *Tridax procumbens* as an anticancer drug by in silico approach. *Struct Chem*. 2022;33(3):703-19. <https://doi.org/10.1007/s11224-022-01882-7>
54. Nadalin P, Kim JK, Park SU. Recent insights into luteolin and its biological and pharmacological activities. *EXCLI J*. 2024;23:787-94.
55. Bhagat VC, Kondawar MS. Antitubercular potential of *Dendrophthoe falcata* (L.) and *Tridax procumbens* (L.) plant extracts against H37Rv strain of *Mycobacterium tuberculosis*. *Int J Pharm Sci Res*. 2019;10:51-9.
56. Chen B, Zhang W, Lin C, Zhang L. A comprehensive review on beneficial effects of catechins on secondary mitochondrial diseases. *Int J Mol Sci*. 2022;23(19):11569. <https://doi.org/10.3390/ijms231911569>
57. Kaushik D, Tanwar A, Davis J. Ethnopharmacological and phytochemical studies of *Tridax procumbens* Linn.: a popular herb in Ayurveda medicine. *Int J Eng Res Technol*. 2020;9:758-68. <https://doi.org/10.17577/JERTV9IS090426>
58. Alrumaihi F, Almatroodi SA, Alharbi HOA, Alwanian WM, Alharbi FA, Almatroudi A, et al. Pharmacological potential of kaempferol, a flavonoid in the management of pathogenesis via modulation of inflammation and other biological activities. *Molecules*. 2024;29(9):2007. <https://doi.org/10.3390/molecules29092007>
59. Jindal A, Kumar P. Antimicrobial flavonoids from *Tridax procumbens*. *Nat Prod Res*. 2012;26(22):2072-7.
60. Salehi B, Venditti A, Sharifi-Rad M, Kregiel D, Sharifi-Rad J, Durazzo A, et al. The therapeutic potential of apigenin. *Int J Mol Sci*. 2019;20(6):1305. <https://doi.org/10.3390/ijms20061305>
61. Sharifi-Rad J, Quispe C, Castillo CMS, Caroca R, Lazo-Vélez MA, Antonyak H, et al. Ellagic acid: a review on its natural sources, chemical stability and therapeutic potential. *Oxid Med Cell Longev*. 2022;2022:3848084. <https://doi.org/10.1155/2022/3848084>
62. Habila JD, Bello IA, Dzikwi AA, Musa H, Abubakar N. Total phenolics and antioxidant activity of *Tridax procumbens* Linn. *Afr J Pharm Pharmacol*. 2010;4(3):123-6.
63. Kahkeshani N, Farzaei F, Fotouhi M, Alavi SS, Bahramsoltani R, Naseri R, et al. Pharmacological effects of gallic acid in health and diseases: a mechanistic review. *Iran J Basic Med Sci*. 2019;22(3):225-37.
64. Akinola AO, Adelowo FE. Chromatographic and spectrophotometric determination of some phenolic compounds in *Tridax procumbens* Linn. stem. *Chem Afr*. 2021;4(1):103-13. <https://doi.org/10.1007/s42250-020-00195-8>
65. Nwauche KT, Anacletus FC, Ighorodje-Monago CC. Assessment of fatty acid, proximate and quantitative phytochemical compositions of matured stem of *Costus afer*. *J Drug Deliv Ther*. 2018;8(6):217-24. <https://doi.org/10.22270/jddt.v8i6.2057>
66. Semwal P, Painuli S, Abu-Izneid T, Rauf A, Sharma A, Daştan SD, et al. Diosgenin: an updated pharmacological review and therapeutic perspectives. *Oxid Med Cell Longev*. 2022;2022:1035441. <https://doi.org/10.1155/2022/1035441>
67. Thalkari AB, Karwa PN, Shinde PS, Gawli CS, Chopane PS. Pharmacological actions of *Tridax procumbens* L.: a scientific review. *Res J Pharmacogn Phytochem*. 2020;12(1):27-30. <https://doi.org/10.5958/0975-4385.2020.00006.0>
68. Wasim M, Bergonzi MC. Unlocking the potential of oleanolic acid: integrating pharmacological insights and advancements in delivery systems. *Pharmaceutics*. 2024;16(6):692. <https://doi.org/10.3390/pharmaceutics16060692>
69. Nwodo NJ, Ibezim A, Ntie-Kang F, Adikwu MU, Mbah CJ. Anti-trypanosomal activity of Nigerian plants and their constituents. *Molecules*. 2015;20(5):7750-71. <https://doi.org/10.3390/molecules20057750>
70. Agrawal S, Mohale D, Talele GS. Pharmacological activities of *Tridax procumbens* (Asteraceae). *Med Plants*. 2010;2(2):73-8. <https://doi.org/10.5958/j.0975-4261.2.2.012>
71. Jeevitha M, Kaarthikeyan G, Yadalam PK, Pradeep Kumar R, Kumar JK. Exploration of wound healing activity of phytochemicals from *Tridax procumbens* using computational approach. *J Pioneer Med Sci*. 2023;20-24. <https://doi.org/10.61091/jpms20231235>
72. Espindola KMM, Ferreira RG, Narvaez LEM, Silva Rosario ACR, da Silva AHM, Silva AGB, et al. Chemical and pharmacological aspects of caffeic acid and its activity in hepatocarcinoma. *Front Oncol*. 2019;9:541. <https://doi.org/10.3389/fonc.2019.00541>
73. Chinnappan BA, Krishnaswamy M, Bal T, Rajora AD. *In vitro* and *in vivo* wound healing efficacy of *Tridax procumbens* extract-loaded carboxymethylcellulose film. *Int J Biol Macromol*. 2023;253(1):126695. <https://doi.org/10.1016/j.ijbiomac.2023.126695>
74. Nguyen V, Taine EG, Meng D, Cui T, Tan W. Chlorogenic acid: a systematic review on the biological functions, mechanistic actions and therapeutic potentials. *Nutrients*. 2024;16(7):924. <https://doi.org/10.3390/nu16070924>
75. Purushothaman JR, Rizwanullah M. Ferulic acid: a comprehensive review. *Cureus*. 2024;16(8):e68063. <https://doi.org/10.7759/cureus.68063>
76. Jachak SM, Gautam R, Selvam C, Madhan H, Srivastava A, Khan T. Anti-inflammatory, cyclooxygenase inhibitory and antioxidant activities of standardized extracts of *Tridax procumbens* L. *Fitoterapia*. 2011;82(2):173-177. <https://doi.org/10.1016/j.fitote.2010.08.016>
77. Ingole W, Mhaske PC, Katade SR. Phytochemistry and pharmacological aspects of *Tridax procumbens* (L.): a systematic and comprehensive review. *Phytomed Plus*. 2022;2(1):100199. <https://doi.org/10.1016/j.phyplu.2021.100199>
78. Mecina GF, Chia MA, Cordeiro-Araujo MK, Bittencourt-Oliveira MDC, Varela RM, Torres A, et al. Effect of flavonoids isolated from *Tridax procumbens* on the growth and toxin production of *Microcystis aeruginosa*. *Aquat Toxicol*. 2019;211:81-91. <https://doi.org/10.1016/j.aquatox.2019.03.011>
79. Chen WH, Ma XM, Wu QX, Shi YP. Chemical constituent diversity of *Tridax procumbens*. *Can J Chem*. 2008;86:892-98. <https://doi.org/10.1139/v08-097>
80. Martin-Quintal Z, Moo-Puc R, Gonzalez-Salazar F, Chan-Bacab MJ, Torres-Tapia LW, Peraza-Sanchez SR. *In vitro* activity of *Tridax procumbens* against promastigotes of *Leishmania mexicana*. *J Ethnopharmacol*. 2009;122(3):463-467. <https://doi.org/10.1016/j.jep.2009.01.037>
81. Sarkar SL, Saha P, Sultana N. *In vitro* evaluation of phytochemical components and antimicrobial activity of the methanolic extract of *Tridax procumbens* L. against pathogenic microorganisms. *J Pharmacogn Phytochem*. 2016;5(5):42-46.
82. Jhample SB, Gajdhane SB, Kasabe PJ, Bhagwat PK, Dandge PB. Phytochemical screening and *in vitro* antimicrobial activity of *Tridax procumbens* L. *Res J Life Sci Bioinform Pharm Chem Sci*. 2015:42-51.
83. Kumar S, Prasad A, Iyer SV, Vaidya S. Pharmacognostical, phytochemical and pharmacological review on *Tridax procumbens* Linn. *Int J Pharm Biol Arch*. 2012;3(4):747-751.
84. Kumar S, Karl PN, Samuel J, Selvakumar M, Shalini K. Antioxidant, antidiabetic, antimicrobial and hemolytic activity of *Tridax procumbens*. *J Chem Pharm Res*. 2016;8(8):808-812.
85. Syed A, Benit N, Alyousef AA, Alqasim A, Arshad M. *In vitro* antibacterial, antioxidant potentials and cytotoxic activity of the leaves of *Tridax procumbens*. *Saudi J Biol Sci*. 2020;27(2):757-761. <https://doi.org/10.1016/j.sjbs.2019.12.031>
86. Andriana Y, Xuan TD, Quy TN, Minh TN, Van TM, Viet TD. Antihyperuricemia, antioxidant and antibacterial activities of *Tridax procumbens* L. *Foods*. 2019;8(1):21. <https://doi.org/10.3390/foods8010021>

87. Owoyemi OO, Oladunmoye MK. Phytochemical screening and antibacterial activities of *Bidens pilosa* L. and *Tridax procumbens* L. on skin pathogens. *Int J Mod Biol Med*. 2017;8(1):24–46.
88. Tejaswini K, Vishwanath Pradeep B, Rudrama Devi K, Shylaja S, Jyothsna K. Phytochemical screening and antimicrobial activities of plant extract of *Tridax procumbens*. *Bioscan*. 2011;6(2):321–3.
89. Joshi RK, Badakar V. Chemical composition and in vitro antimicrobial activity of the essential oil of the flowers of *Tridax procumbens*. *Nat Prod Commun*. 2012;7(7):941–2. <https://doi.org/10.1177/1934578X1200700736>
90. Kamble VA, Moon AH. Antifungal activity of crude extracts from *Tridax procumbens* L. against potentially pathogenic fungal species. *Int J Curr Res*. 2015;7(6):16930–4.
91. Tiwari S, Antu K, Swati M. Study of the antifungal potential of leaf extracts of *Tridax procumbens*. *Afr J Biol Sci*. 2024;6(15):1295–305.
92. Maldhure AK, Kalambe NA, Bhokare DD. Antifungal activity of extract from the leaves of *Tridax procumbens* Linn. *Int J Res Biosci Agric Technol*. 2017;5(2):962–4.
93. Ashish PV, Annasaheb NS. Antimicrobial and antifungal activity of *Tridax procumbens* Linn whole plant ethanolic extract on different pathogens and its phytochemical screening. *World J Pharm Res*. 2015;4(10):784–813.
94. Miya MF, Islam Z, Shahriyar S, Khan MRH, Reza MS. Antifungal potential of tridhara (*Tridax procumbens*) leaves. *Asian J Med Biol Res*. 2015;1(3):686–9. <https://doi.org/10.3329/ajmbr.v1i3.26497>
95. Acharya S, Srivastava RC. Antifungal property of *Tridax procumbens* L. against three phytopathogenic fungi. *Arch Pharm Sci Res*. 2010;2(1):258–63.
96. Hemalatha R. Anti-hepatotoxic and antioxidant defense potential of *Tridax procumbens*. *Int J Green Pharm*. 2008;2(3):164. <https://doi.org/10.4103/0973-8258.42736>
97. Mundada S, Shivhare R. Pharmacology of *Tridax procumbens*, a weed. *Int J Pharmtech Res*. 2010;2(2):1391–4.
98. Manjamalai A, Varghese SS, Haridas A, Grace VMB. Antifungal, anti-inflammatory and GC-MS analysis for bioactive molecules of *Tridax procumbens* L. leaf. *Asian J Pharm Clin Res*. 2012;5(1):139–45.
99. Greenwell M, Rahman PK. Medicinal plants: their use in anticancer treatment. *Int J Pharm Sci Res*. 2015;6(10):4103–12.
100. Priya VP, Radhika K, Sivakumar R, Sri Ramchandra M, Prameela Devi Y, Srinivas Rao A. Evaluation of anticancer activity of *Tridax procumbens* flower extracts on PC3 cell lines. *Int J Adv Pharm Sci*. 2011;2(1):115–20.
101. Pareek H, Sharma S, Khajja BS, Jain K, Jain GC. Evaluation of hypoglycemic and antihyperglycemic potential of *Tridax procumbens* (Linn.). *BMC Complement Altern Med*. 2009;9:1–7. <https://doi.org/10.1186/1472-6882-9-48>
102. Patel S, Goyal A. Recent developments in mushrooms as anticancer therapeutics: a review. *3 Biotech*. 2012;2:1–15. <https://doi.org/10.1007/s13205-011-0036-2>
103. Oladunmoye MK. Immunomodulatory effects of ethanolic extract of *Tridax procumbens* on Swiss albino rats orogastrically dosed with *Pseudomonas aeruginosa*. *Trends Med Res*. 2006;1(4):152–5.
104. Ghosh P, Biswas S, Biswas M, Dutta A, Sil S, Chatterjee S. Morphological, ethnobiological and phytopharmacological attributes of *Tridax procumbens* Linn. (Asteraceae): a review. *Int J Sci Res Biol Sci*. 2019;6(2):182–91. <https://doi.org/10.26438/ijrbs/v6i2.182191>
105. Nisha MH. Phytochemical and biological investigation of *Tridax procumbens* leaves [doctoral dissertation]. Dhaka: East West University; 2015.
106. Gutiérrez-Rebolledo GA, Drier-Jonas S, Jiménez-Arellanes MA. Natural compounds and extracts from Mexican medicinal plants with antileishmaniasis activity: an update. *Asian Pac J Trop Med*. 2017;10(12):1105–10. <https://doi.org/10.1016/j.apjtm.2017.10.016>
107. Appiah-Opong R, Nyarko AK, Dodoo D, Gyang FN, Koram KA, Ayisi NK. Antiplasmodial activity of extracts of *Tridax procumbens* and *Phyllanthus amarus* in *in vitro Plasmodium falciparum* culture systems. *Ghana Med J*. 2011;45(4):143–50.
108. Choochote W, Chaithong U, Kamsuk K, Jitpakdi A, Tippawangkosol P, Tuetun B, et al. Repellent activity of selected essential oils against *Aedes aegypti*. *Fitoterapia*. 2007;78(5):359–64. <https://doi.org/10.1016/j.fitote.2007.02.006>
109. Fernando SS, Jayasooriya RG, Samarakoon KW, Wijegunawardana NA, Alahakoon SB. Citrus-based bio-insect repellents: a review on historical and emerging trends in utilizing phytochemicals of citrus plants. *J Toxicol*. 2024;2024:6179226. <https://doi.org/10.1155/jt/6179226>
110. Zahir AA, Rahuman AA, Kamaraj C. Adult emergence inhibition and adulticidal activities of medicinal plant extracts against *Anopheles stephensi* Liston. *Asian Pac J Trop Med*. 2010;3:878–83. [https://doi.org/10.1016/S1995-7645\(10\)60211-8](https://doi.org/10.1016/S1995-7645(10)60211-8)
111. Kamaraj C, Bagavan A, Elango G, Zahir AA, Rajakumar G, Marimuthu S, Rahuman AA. Larvicidal activity of medicinal plant extracts against *Anopheles subpictus* and *Culex tritaeniorhynchus*. *Indian J Med Res*. 2011;134(1):101–6. <https://doi.org/10.1007/s00436-010-1816-z>
112. Selvakumar P. Pharmacognostical and phytopharmacological review on coat buttons. *J Prev Diagn Treat Strateg Med*. 2022;202–208. [https://doi.org/10.4103/jpdtm.jpdtm\\_55\\_22](https://doi.org/10.4103/jpdtm.jpdtm_55_22)
113. Shrivastav A, Kumar Mishra A, Abid M, Ahmad A, Fabuzinadah M, Khan NA. Extracts of *Tridax procumbens* Linn leaves cause wound healing in diabetic and non-diabetic laboratory animals. *Wound Med*. 2020;29(1):100185. <https://doi.org/10.1016/j.wndm.2020.100185>
114. Yaduvanshi B, Mathur R, Mathur SR, Velpandian T. Evaluation of wound healing potential of topical formulation of leaf juice of *Tridax procumbens* L. in mice. *Indian J Pharm Sci*. 2011;73(3):303–6.
115. Deshmukh R, Agrawal R, Mishra MU. Formulation and evaluation of ointment containing natural wound healing activity of *Tridax procumbens*. *Res J Pharm Technol*. 2018;11:4543–6. <https://doi.org/10.5958/0974-360X.2018.00831.4>

#### Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonpublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonpublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc  
See [https://horizonpublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

**Publisher information:** Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.