



REVIEW ARTICLE

Sustainable management of saline and sodic soils for growing vegetable crops - A review

J Marimuthu¹, K R Vijayalatha^{1*}, V Jegadeeswari², S Nithila³, A Sabir Ahamed¹, Geethalakshmi Indran⁴, T Uma Maheswari⁴, R Rajeswari¹, N Malini³, S Kumar⁴ & P Yasodha⁵

¹Department of Vegetable Science, Horticultural College and Research Institute for Women, Tamil Nadu Agricultural University, Tiruchirappalli 620 027, Tamil Nadu, India

²Grapes Research Station, Tamil Nadu Agricultural University, Theni 625 526, Tamil Nadu, India

³Department of Basic Science, Horticultural College and Research Institute for Women, Tamil Nadu Agricultural University, Tiruchirappalli 620 027, Tamil Nadu, India

⁴Department of Fruit Science, Horticultural College and Research Institute for Women, Tamil Nadu Agricultural University, Tiruchirappalli 620 027, Tamil Nadu, India

⁵Department of Plant Protection, Horticultural College and Research Institute for Women, Tamil Nadu Agricultural University, Tiruchirappalli 620 027, Tamil Nadu, India

*Correspondence email - vijayalatha.kr@tnau.ac.in

Received: 25 April 2025; Accepted: 19 June 2025; Available online: Version 1.0: 26 July 2025

Cite this article: Marimuthu J, Vijayalatha KR, Jegadeeswari V, Nithila S, Sabir Ahamed A, Geethalakshmi I, Uma Maheswari T, Rajeswari R, Malini N, Kumar S, Yasodha P. Sustainable management of saline and sodic soils for growing vegetable crops - A review. Plant Science Today (Early Access).

<https://doi.org/10.14719/pst.9100>

Abstract

Vegetables are essential that play a crucial role in human nutrition. Salinity is a major constraint affecting agricultural productivity globally. Salt affects soils comprise approximately 20 % of cultivated land and 33 % of irrigated land. Salt stress hampers plant growth, thereby reducing the yield and quality of various crops. Salinity and sodicity adversely affect the biological, physical and chemical properties of soil, leading to reduced productivity and land degradation, particularly in irrigated and rainfed agricultural systems. Salinity reduces protein, fatty acid and total carbohydrate content in crops, while often increasing the accumulation of amino acids. The presence of soluble salts and excess sodium ions (Na^+) in soil adversely affects plant health, emphasizing the need for effective resource management and sustainable practices. High salinity leads to surface crusting, reduced water infiltration, sodium-induced soil dispersion and decreased hydraulic conductivity (HC), all of which negatively impact plant development. Sodicity refers to the presence of excessive exchangeable sodium in soil relative to calcium and magnesium, which disrupts soil structure and fertility. High sodicity inhibits plant growth due to salt toxicity, nutritional imbalances and reduced availability of essential minerals in the soil. This review discusses the impact of saline and sodic soils on various vegetable crops and explores sustainable management practices to mitigate their effects.

Keywords: management practices; saline soil; sodic soil; vegetable crops

Introduction

Vegetables are essential for small-scale farmers because they yield much higher returns per hectare than traditional staple crops (1). Compared to staple crops, vegetables are believed to be more susceptible to adverse climatic conditions such as drought, high or low temperatures, salt, waterlogging, an excess or deficiency of mineral nutrients and variations in soil pH (2). High levels of soluble salts in the soil limit the production of most crops, including vegetables, in many areas of the world particularly in arid and semi-arid regions such as parts of South Asia (e.g., India and Pakistan), the Middle East (e.g., Iran, Iraq and Saudi Arabia), North Africa (e.g., Egypt and Algeria), Central Asia (e.g., Uzbekistan and Turkmenistan) and parts of Australia and southwestern United States. Climate change is expected to exacerbate these environmental issues by increasing evapotranspiration and reducing freshwater availability in

these vulnerable regions (3). Significant crop-to-crop diversity in vegetable crop salt tolerance has been noted, much like with other crops (4). While crops such as broccoli, cauliflower, tomato, eggplant, potato, turnip, radish, lettuce, pumpkin, cucumber and pepper exhibit moderate sensitivity, red beet (*Beta vulgaris*) is relatively salt-tolerant. In contrast, carrots, onions, peas and okra are highly susceptible to salinity (Table 1 and 2).

This term “salinity” is derived from the Latin words “*salinium*,” meaning “position or quality of being,” and “salt cellar.” “Salt” is the term used to describe the dissolved salt found in soil or water. Soil salinity is a significant issue that threatens agricultural plants and restricts worldwide agriculture, especially on irrigated farmlands, in areas where high-salt groundwater is used for irrigation (5). Plants that are salt tolerant, or resistant to their natural strength, may withstand the damaging effects of excessive salt on their roots or leaves without experiencing any severe adverse effects (6).

Table 1. Threshold level of salinity for vegetable crops

Vegetable	Threshold levels (dS m ⁻¹)	Reference
Pea (<i>Pisum sativum</i> L.)	1.5	(55)
Potato (<i>Solanum tuberosum</i>)	1.7	(56)
Cauliflower (<i>Brassica oleracea</i> var. <i>Botrytis</i>)	1.8	(55)
Sweet pepper (<i>Capsicum annuum</i>)	1.5	(56)
Broccoli (<i>Brassica oleracea</i> var. <i>Italic</i>)	1.8	(56, 57)
Carrot (<i>Daucus carota</i>)	1.0	(56, 57)
Bhendi (<i>Abelmoschus esculentus</i> L.)	1.2	(55)
Brinjal (<i>Solanum melongena</i>)	1.1	(56)
Tomato (<i>Solanum lycopersicum</i>)	2.5	(56)

Table 2. Categorization of vegetable cultivars based on salinity tolerance

Moderately sensible (0-4 mS/cm)	Sensible (4-6 mS/cm)	Tolerant (6-8 mS/cm)	Highly tolerant (8-12 mS/cm)	Reference
Carrot	Onion			
Cucumber	Lettuce	Cabbage		
Watermelon	Melon	Tomato	Asparagus	
Beans	Potato	Spinach	Beetroot	
Radish	Pumpkin			(21, 84, 85)

Approximately 25 % of the world's irrigated land equivalent to 45 million hectares is affected by salinity, compromising nearly one-third of global food production (7). Global agriculture, particularly in irrigated places, is hampered by soil salinity, another significant issue in locations where irrigation uses high ground salt water (5). The profitability of many horticultural harvests, especially extremely soft vegetables at the plant's entry, is reduced by excessive soil salt. Symptoms of salt stress in susceptible plants might include decreased growth, turgor loss, leaf shrinkage, folding and epinasty, leaf removal, decreased photosynthesis, respiratory abnormalities, loss of cellular integrity, tissue necrosis and plant death (8, 9).

If the extract saturation (ECE) electrical conductivity in the root zone is greater than 4 dS m⁻¹ (around 40 mM NaCl) at 25 dC and includes 15 % exchangeable sodium, the soil is considered salty (Table 3). The yield of most agricultural plants decreases with ECE, even if the bulk of crops exhibit a dip in productivity at lower ECs (10, 11). Globally, soil erosion and decreased agricultural productivity are caused by soil salinization (12). Most vegetable crops are natively glycophytes, making salt stress one of the most severe environmental conditions that limits their growth.

Approximately one-third of the world's irrigated land is affected by salinity (13), posing a significant threat to agricultural productivity. Alongside other environmental factors such as high winds, elevated temperatures, drought and floods, soil salinity is among the most damaging, especially in uncultivated areas, where it severely reduces crop yield and quality (14, 15). Additionally, poor soil structure and inadequate aeration further limit plant growth and productivity (16).

Soil salinity is a major issue in irrigated agriculture. Saline soils are common in hot, arid parts of the world with

little potential for agriculture. Since, most crops grown in these places are irrigated, 20 % of irrigated land globally suffers from secondary salinization, which is exacerbated by inadequate irrigation (17). Salinity is the result of soil salts building up when rainfall is insufficient to eliminate ions from the soil profile (18).

Excessive gypsum treatment has been demonstrated to reduce electrical conductivity and the sodium adsorption ratio (SAR) while increasing the removal of excess Na⁺ from soil. Gypsum is a common agricultural soil additive used for sodic soil reclamation because of its high solubility, ease of application and low cost, helping to mitigate the negative effects of high salt concentrations in irrigation water.

Salt naturally exists in soil, surface water and groundwater. The most common salt is sodium chloride, though other salts like magnesium, calcium and potassium may also be present. While soluble salts may have less sodic acid, they often contain high levels of sodium. Sodic soils, characterized by high levels of exchangeable sodium, are unsuitable for most crops as they elevate soil pH (typically 8.5 -12), disrupt soil structure and impair water infiltration and aeration (Table 3). This disrupts the soil's chemical balance and structure. As a result, the soil cannot easily absorb air, rain, or irrigation water. It becomes sticky when wet and forms hard crusts when dry. This problem usually doesn't occur in sandy soils, as they lack enough clay to develop such issues (19). This review focuses on the cultivation of various vegetable crops under saline and sodic soil conditions and discusses effective management strategies to enhance productivity in such challenging environments (Table 4).

Saline soil

Saline soils are defined by the presence of excess soluble salts, which can hinder the growth and yield of most agricultural crops, even at electrical conductivity of the

Table 3. Characteristics of salt affected soil

Different Salt Classes Affecting Soils	Electrical conductivity (ECE) at 25 C (dS m ⁻¹)	Exchange Sodium percentage (ESP)	Sodium adsorption ratio (SAR)	Reaction (pH value)	Reference
Saline soil	>4	<15	< 13	<8.5	
Sodic (alkali) soil	<4	>15	>13	>8.5-10	(54)

Table 4. Management of salinity

Techniques of management	Utilizing vegetables	Reference
Lowering of the groundwater	Groundwater withdrawal from rivers can lower coastal areas' groundwater levels, allow seawater intrusion and hinder crop cultivation. Reducing salt leaching is necessary to lessen the consequences of salinity and lowering the water table and ensuring adequate drainage are crucial to avoiding salt buildup	(66)
Building of structures for water harvesting	To mitigate salinity impacts, groundwater use must be substituted by freshwater irrigation. Proper rainwater harvesting structures, such as floodwater harvesting, macro-catchments and micro-catchment, are needed to sustain agricultural livelihoods in drylands, including spat irrigation and runoff farming	(67)
Restoration of salinized soils	Soil reclamation is a method that extracts soluble salts from crop roots, reducing salinity impact through practices like leaching, improved water management, surface and subsurface drainage, organic fertilizers and salt-tolerant cultivars	(6, 68, 69)
Leaching of salt	The leaching process for bell peppers increases with the salt of the irrigation water	(70)
Systems of surface and subsurface drainage	Drainage is the act of eliminating surface or subsurface water using artificial or natural systems, which lowers the water table and lessens the possibility of salt buildup and a rising groundwater table	(71)
Organic or chemical fertilizers	By lowering Na^+ toxicity, boosting water-holding capacity and releasing vital minerals, organic fertilizers enhance soil health. Potash fertilizers improve soil salinity, increase crop production and aid in element uptake. Nitrogen fertilizers overcome boron toxicity and improve soil's Cl^- toxicity effects	(72, 73)
Mulching the soil	Swiss chard-Mulching using rice straw and stones to increase crop yield	(74)
Calcium	Sweet Pepper and tomato improve fruit quality and output while reducing blossom end rot	(75)
Phosphorus	Salinity sensitivity was reduced by up to 3.5 dS m^{-1} by radish	(76)
Sulfur	<i>Brassica</i> spp. and legume crops: enhanced salinity stress and defensive mechanisms	(77)
Relative humidity	The cultivation of melons under salt stress fared better at 70 % relative humidity than at 30 %	(78)
Grafting to tolerant rootstock	Melon and pumpkin rootstock combinations: Melon rootstocks almost eliminate Na, whilst pumpkin rootstocks exclude 74 % of accessible Na	(79)
Seed priming	Melons treated with 18 dS m^{-1} NaCl solution had less adverse effects than those of salty irrigation	(80)
Foliar application of nutrient	Brinjal- K_2HPO_4 increased fruit yield	(81)
Elevated CO_2 concentration	Tomato-increasing aerial CO_2 concentration, alleviate the negative salinity effects	(82)
Irrigation methods	Sprinkler irrigation reduces salt leaching by releasing little quantities of water for infiltration, making it perfect for regions that are often watered	(6)
Crop rotation	Drip irrigation is a highly effective method for saline-irrigated lands, as it maintains low salt levels in the plant root zone	
	Crop rotation, a strategy to combat salinity, commonly uses cultivars that are tolerant of both water and salt	
Crop rotation	Long-fallowing crops, however, may cause groundwater levels to rise, which would impede agricultural development. According to studies, annual and perennial crops should be cultivated alternately	(83)

extract (ECe) levels below standard thresholds (11). This ECe leads to reduced yields for most agricultural plants. Saline soils are well-structured nature, saline soils are not sodic and contain enough soluble salts to negatively impact the development of most agricultural plants (20). Salt types and concentrations are the primary determinants of the chemical properties of soils that are classified as saline. It is the quantity of soluble salts that determines the soil solution's osmotic pressure. Magnesium and calcium concentrations on the exchange complex and in the soil solution might differ significantly. The chemical composition of saline soils varies depending on the dominant salts, with sodium chloride being the most common. However, magnesium, calcium and occasionally potassium both in soluble and exchangeable forms also contribute significantly to the ionic profile (21). Saline soils are widespread in hot, arid regions and generally have limited agricultural potential. Secondary salinization, which affects about 20 % of the world's irrigated land, results primarily from poor irrigation management and is a major constraint to crop production in these areas (17). Human activities, especially irrigated agriculture in arid and semi-arid zones, accelerate secondary salinization by causing salt accumulation in soils and water sources. Additionally, natural

processes such as mineral weathering contribute to the gradual buildup of salts. When minerals weather, they release salts, which are composed of electrically charged atoms or molecules called ions, into the soil. Secondary salinization, which affects 20 % of irrigated land worldwide and is made worse by inadequate irrigation management, is cited as the primary crop harming most crops farmed in these countries. Irrigated agriculture, a common human activity in arid and semi-arid regions, causes secondary salinization of the soil and water sources. Because saline soil is typically found in hot, arid regions of the world, they offer limited promise for agriculture (17). Electrically charged atoms or molecules known as ions make up salts, which are found in soil. Weathering minerals releases ions into the soil. Irrigation and fertilization are possible because they move upstream from shallow groundwater to land. Salts build up when the soil profile isn't sufficiently cleared of ions by rainfall (18).

Impact of salinity stress on the plants

Salinity has a complicated effect on plants and can cause ion toxicity, osmotic stress, imbalances in hormones, nutrient uptake and antioxidant activity. The first physiological reaction to salt stress is a reduction in stomatal closure and leaf

elongation (22, 23). Excessive accumulation of sodium (Na^+) and chloride (Cl^-) ions in the lower leaves disrupts the ionic balance, leading to a decline in nutrient absorption and reduced leaf area index ultimately hindering plant development.

In sodic soils, the high osmotic pressure causes plant cells to lose more water than they can absorb, inducing water stress (24, 25). A scarcity of water results in a series of physical, signaling, gene expression, metabolic and physiological processes and activities that eventually limit photosynthesis, biomass accumulation, the leaf area index and yield (26, 27). The primary cause raising the concentration of Na^+ and Cl^- ions and producing plant toxicity in sodic soils is the fact that 50-80% of the soluble salts are NaCl . These ions significantly impact the photosynthetic signaling system or the chemical process behind enzyme function (28, 30).

Sodium ions in saline soils interfere with potassium uptake due to similarities in ionic radii and transport mechanisms. This competitive inhibition often results in potassium (K^+) deficiency in plant tissues, especially in leaves, where K^+ plays a vital role in stomatal regulation, turgor maintenance and overall photosynthetic efficiency (31-33). Salt tension has a significant impact on the intricate plant-physiological and biochemical process of photosynthesis in vegetables (20, 34). In plants that are susceptible to salt, stomata closure may result in a reduction in the efficiency of carboxylation and photosynthesis. Additionally, the over expression of pheophorbide and oxygenase causes chlorophyll to degrade, which impacts photosynthesis (20). Additionally, salt can cause anomalies in the oxygen-evolving complex, cyclic electron transport and PSII activity (35, 36). In lettuce, onions and tomatoes, it was demonstrated that salinity stress reduced stomatal conductance and transpiration rates, deteriorated pigments and light-harvesting complexes and hence reduced the quantum yield of photosynthetic energy and energy dissipation through non-photochemical means.

The secondary effect of salinity on plant cells is the accumulation of dangerous reactive oxygen species (ROS), which change gene expression and cause DNA methylation, regardless of the primary effects of salt stress (37, 38). Furthermore, ROS can cause lipid peroxidation, which raises the membrane's permeability and fluidity (40). In response, plants activate a complex defense system involving both enzymatic (e.g., superoxide dismutase, catalase) and non-enzymatic (e.g., ascorbate, glutathione) antioxidants to mitigate ROS-induced damage (30, 40) (Table 5).

Changes in soil salinity and sodicity

In arid and semiarid environments, irrigation (irrigation agriculture) is the main cause of soil salinity, an issue that is becoming more common in crop production globally. Salinity

has an impact on almost one-third of all irrigated land on Earth. When evapotranspiration exceeds precipitation and drainage is inadequate, salts accumulate at the soil surface. A high salt content in the soil profile is a characteristic of saline soils. A range of ion species with varying compositions are frequently present in amounts that impact crop growth (Na^+ , Cl^- , HCO_3^- , PO_4^{3-} , Ca^{2+} , Mg^{2+} , SO_4^{2-} and borate) (41).

Soil quality is influenced by land use and soil management practices and it differs regionally from field to greater area size (42). Management practices and land use have an impact on the extent and direction of soil changes. It may be beneficial to use and manage land appropriately to enhance soil properties, reduce soil degradation and eventually attain agricultural sustainability (43). Understanding the spatial variability of soil quality and the factors influencing it is essential for planning sustainable land use and improving long-term agricultural productivity (44).

Primary salinization

This happens naturally and is caused by salts collecting because the parent material or groundwater has a lot of salts (3). Salts are naturally formed in both dry and moist sections of the earth (45). Salt-affected soils (SAS) can develop due to high salt levels in the soil, a shallow groundwater table, or the use of saline water for irrigation (46). A major environmental problem that affects natural farming ecosystems worldwide is salinity during drought (47, 48).

Secondary salinization

Secondary salinization results from human-induced activities such as poor drainage, improper fertilizer application, inefficient irrigation systems and the use of salt-laden water for irrigation. Population increases and socioeconomic conditions are the main sources of these problems (3). As there is insufficient drainage, salts that were once evenly dispersed throughout the soil profile are carried to the upper layers by irrigation water and then left behind as the water evaporates. Secondary salinization caused by soil mineral weathering, fertilizer use, immobilized salts already precipitated in soils, atmospheric salt depositions, such as in coastal areas, water logging and excessive irrigation water salinity are the main causes of excessive salt accumulation in soil surface horizons (49).

Mechanism of salinity tolerance

Salinity tolerance in plants involves a complex interplay of morphological, biochemical, molecular and physiological mechanisms that influence growth and yield (50) (Fig 1). Reducing water loss from stomata and cuticles and increasing water intake by roots promote osmotic adjustment and support morphological and physiological adaptation for

Table 5. Enhancement of possible vegetable crops' resistance to salt through several methods

Vegetable crop	Method used to increase resistance to salt	Characteristics for improved	Reference
Tomato	Application of salicylic acid, thiazuron and amino acids exogenously	Enhanced dry biomass	(60, 66)
Brinjal	Using a foliar spray to apply glycinebetaine (GB) exogenously	Greater yield and growth	(59)
Potato	Applying salts like NaCl and CaCl_2 to potato tubers before they are harvested	The shoots' higher dry weight	(62)
Okra	K and humic acid application in a saline media	Increased dry biomass	(58)
Pea	application of methyl jasmonate (Me-JA) to seeds prior to sowing	Higher concentration of proline	(63)
Cauliflower	Adding nitrogen to the growing medium	Higher yield	(64)
Broccoli	Topically applying urea or methyl-jasmonate (MeJA)	Improved growth	(65)

resistance to salt-induced osmotic stress (51). A number of molecular networks regulate tolerance and adaptation to salt stress. These networks initiate response mechanisms such as stress protein synthesis, antioxidant overexpression and the accumulation of appropriate solutes to stabilize cells, repair damaged membranes and protect proteins (25, 52).

Based on their response to salt, plants are classified into halophytes and glycophytes. Halophytes thrive and complete their life cycles in saline environments, whereas glycophytes are generally salt-sensitive (53). Plants use two primary strategies to combat salt stress: tissue tolerance and salt avoidance. Ion compartmentalization is also carried out by plants in their tissues. To control their osmotic pressure and preserve the integrity of their metabolic processes, plants continuously produce water-soluble, low molecular weight compatible solutes including proline, glycine betaine and sugars. Plants produce a range of enzymatic and non-enzymatic antioxidants to address the detrimental effects of salt. To protect the cytoplasm of the plant cell from ion toxicity and water stress, ion compartmentalization in the vacuole creates a constant salt concentration in the cytosol during plant tissue tolerance (2).

Effect of salinity and crop reaction in vegetable crops

Bhendi (*Abelmoschus esculentus* L.)

Effect of salinity and crop reaction: Bhendi (*Abelmoschus esculentus* L.) is considered a moderately salt-tolerant crop. Salinity stress affects bhendi, a crop that is semi-tolerant. It can prevent seed germination and lower cotyledon activity of Na^+ , sugar and phenol. Fresh pod yield, shoot length and shoot and root weights are all greatly decreased by the salinity of the rooting media. In reaction to salt stress caused by NaCl , ion absorption and ratios are also impacted, with higher concentrations of Na^+ and Cl^- and lower concentrations of K^+ and Ca^{2+} .

Strategies for improving salinity tolerance: Research on enhancing salinity tolerance in okra remains limited (58). Incorporating K and humic acid into a saline medium has been shown to significantly increase okra's tolerance to salt, especially during the seedling stage (Table 5).

Brinjal (*Solanum melongena* L.)

Effect of salinity and crop reaction: In eggplant (*Solanum melongena* L.), salinity stress moderately affects physiological functions, including internal CO_2 concentration, shoot and root biomass and gas exchange parameters, while water use efficiency remains largely unaffected (59). Fruit weight and quantity per plant are greatly reduced by salinity.

Strategies for improving salinity tolerance: There aren't many ways to combat the losses in eggplant production caused by salt in saline environments, however exogenous application of inorganic fertilizers, suitable solutes and bacterial plant growth promotion have been discovered to be effective methods (59). It has been discovered that the detrimental effects of NaCl on plant development, fruit output and total soluble sugar levels can be lessened by applying di-potassium hydrogen orthophosphate (K_2HPO_4) topically.

Potato (*Solanum tuberosum* L.)

Effect of salinity and crop reaction: Salinity stress significantly impairs the growth and productivity of potato (*Solanum tuberosum* L.), particularly in salt-sensitive cultivars. It adversely affects seedling biomass accumulation, reduces shoot and root lengths and alters molecular responses involved in stress signaling and metabolism. The resulting physiological imbalances, such as ion toxicity and osmotic stress, lead to reduced tuber yield and quality.

Strategies for improving salinity tolerance: To enhance salt tolerance in potatoes, several strategies have been employed, including the application of exogenous nutrients, 5-aminolevulinic acid (ALA) and genetic engineering. Pre-treatment with NaCl and CaCl_2 has shown effectiveness in preparing plants for salt stress, while potassium supplementation in the rooting medium helps mitigate salinity effects, particularly under potassium-deficient conditions. However, excessive potassium application beyond the recommended threshold does not confer additional benefits.

Salt-tolerant cultivars

The accumulation of salts in soil is a widespread issue that negatively influences plant physiological processes and reduces crop yields. The development and cultivation of salt-tolerant potato cultivars represent a critical strategy for mitigating these effects.




Fig. 1. Stress signaling, osmotic adjustment, enzyme activity, polyamines and ionic compartmentalization all regulate the physiological and metabolic processes of halophytes.

Sodicity

Sodic soils are those that have so much exchangeable salt in their exchange complex that it hinders the growth of most agricultural plants. A non-saline soil that has enough exchangeable sodium to negatively impact crop yield and soil physical characteristics by altering soil structure is often referred to as a sodic soil (20). Poorly structured soils with colloidal clays scattered over the uppermost layer are known as sodic soils. Some of these soils' characteristics include poor aeration, crusting of the surface soil, low permeability and infiltration rates, difficulty tilling and difficulty permitting plant roots to penetrate.

Sodic soils typically have a high pH (above 8.5) due to the absence of calcium carbonate buffering and the alkaline hydrolysis of sodium salts (86, 87). High sodicity inhibits plant development due to a lack of mineral nutrients in the soil, a plant nutritional imbalance and Na toxicity. Sodicity refers to the ratios of transferable sodium to calcium and magnesium in soil (88). Sodic conditions develop when sodium ions preferentially bind to negatively charged sites on soil colloids, displacing calcium and magnesium and causing clay dispersion (89). Sodic soils are the most common problem in the world's irrigated desert and semi-arid regions. Furthermore, the presence of bicarbonates, soluble sodium carbonates and interchangeable sodium in irrigation water adversely affects the increasing salinity/alkalinity in the agricultural soils in these regions (90). Crop type, variety, developmental stage, soil texture, salt type and quantity, cultural techniques and climate (temperature, relative humidity and rainfall) all affect how severe the negative effects (91, 92). Sodic soils inhibited asparagus and tomato development more than saline soils did, indicating that both plants are vulnerable to sodicity, as shown by the sensitivity of specific varieties listed in Table 6. Similarly, the bean demonstrated a strong sensitivity to sodicity by failing to survive under such conditions (93).

The impacts of sodicity

- Decreased water movement through the soil, which inhibits leaching and might eventually lead to the buildup of salt and the production of salt water.
- Increasing corrosion and spreading throughout the subsurface, which may result in the formation of lanes and tunnels.
- It eliminates aggregation, which manifests as thick, structure-free soil.
- Water infiltration is stopped by dispersion over the soil surface, which causes crusting and sealing.

Measures of sodicity

- There are either too many plants or trees standing, too few heavy plants, or poor plant development that is below average.

Table 6. Sodic soil tolerant crop varieties

Crops	Varieties	Reference
Hot pepper (chilli)	Chaman, Jawala	
Tomato	Azad T2, Angurlata	
Spinach	Chikari, K Hari	(96)
Garlic	Hansa, Gattar Gola	

- Differently sized plants.
- Poor rainfall infiltration overhead.
- The shallowness of plants.
- The blockage of the mud channel makes it challenging to immerse oneself outside the study area (72).
- Green or fluctuating water pools.
- The soil is often darker in the shadow because of the intricate development of Na-humic.

Effect of sodicity on vegetables

Like salinity, alkalinity and tolerance vary greatly among plants and their species. When the ESP and salt content in the soil solution are higher than what is allowed for each crop, crop yields are often impacted. Winter crops are often hardier than summer crops in terms of salt tolerance. Therefore, it is suggested that low ET winter vegetable crops (less than 400 mm) can be grown in areas with low rainfall that are categorized as arable crops during the summer. The best course of action would be to select rainfed crops for Kharif and crops that need less water for *Rabi* (94) (Table 7).

Deficits in nutrients and ion toxicities

Sodic soils, which are created by electron and proton activity, may limit the availability of plant nutrients due to limited water and oxygen flow rates, even though high pH levels can remain throughout the soil profile (95). The sodic soils of IGP have been discovered to exhibit significant deficiencies in OM, accessible N, Ca and Zn. Sodic soils often have low levels of nitrogen and organic carbon (OC). Despite the advantages of applying organic inputs like farmyard manure (FYM), regenerated soils lack organic matter (OC) (96).

Chemical amendments

Sodic conditions need the use of tillage, amendments and Na^+ leaching in crop cultivation. By applying chemicals, industrial wastes, composts, microbial inoculants and polymers, sodic soils' physico-chemical properties are improved. The most often utilized chemical to get beyond structural and nutritional constraints is gypsum. However, there is a growing shortage of high-quality agricultural-grade gypsum, which has raised interest in inexpensive, environmentally friendly soil conditioners for reclamation projects (97, 98).

Restoration and handling of sodic soils

Gypsum

Gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$), a soluble calcium salt, is widely used to reclaim sodic soils, particularly those with an Exchangeable

Table 7. Vegetable crops tolerance for sodic soil

Less tolerant (ESP < 20)	Semi-tolerant (ESP 20-40)	Highly tolerant crops (ESP > 40)	Reference
Ginger, turmeric, cluster bean, pea and cowpea	Onion, potatoes, ash gourd, radishes, carrots, cauliflower, fenugreek, fennel, tomatoes, garlic and okra	Eggplant, spinach and sugar beets	(96)

Sodium Percentage (ESP) below 15. Excess Na^+ may be eliminated by the soil exchange complex because gypsum increases the quantity of interchangeable Ca^{2+} available. In agriculture, gypsum has long been utilized as an ameliorant and as a calcium and sulfur fertilizer. Gypsum has long been used to improve agricultural soil and as a fertilizer for calcium and sulfur. Applying gypsum as a soil conditioner prevents runoff-induced nitrogen depletion and soil erosion (99).

Technology using gypsum beds: In regions where irrigation water contains high levels of sodium carbonate, gypsum-bed technology offers an efficient method to mitigate sodicity. Traditional methods include mixing gypsum into the soil or placing it in irrigation channels wrapped in porous bags. However, using properly designed gypsum dissolving beds yields notably superior outcomes (100). The gypsum-bed technology transports irrigation water via a brick-cement chamber that is filled with a gypsum clod. The size of the chamber is determined by the discharge from irrigation water tube wells and the amount of sodium carbonate that is left over. The water channel is connected to one side, while the water fall box is connected to the other. Ten centimeters above the chamber floor is an iron bar mesh covered with a 2 mm \times 2 mm wire net. If farmers make the necessary modifications, they may also turn their tube wells into gypsum rooms. Dissolving and replenishing the gypsum in the chamber is done by sodic water flowing from the bottom (55). There is no change in the rationale for calculating the gypsum required regardless of the application method. On the other side, the method utilized affects the application time. A single basal dosage of the whole amount of gypsum is administered to the soil. Water-applied gypsum does not accumulate in the soil since it is neutralized before application (100).

Regular applications of agricultural wastes like sawdust, rice hulls, sugar mill waste and so on, as well as thick dressings of organic manures, have been shown to be advantageous. Preserving and enhancing the physical characteristics of soil while mitigating the adverse effects of excessively changing salt levels. Whenever possible, organic fluids that are in danger of becoming alkaline should be supplemented with organic materials. But organic modification is ineffective in reducing the negative effects of alkali water without gypsum. Farmyard Manure (FYM) added gypsum to water to boost the yields of knol-khol, bottle gourds, ridge gourds, bitter gourds, eggplants, broccoli, cluster beans, cauliflower and potatoes (94). Gypsum and FYM were added to counteract the adverse effects of alkaline water, which significantly increased crop growth and yield. They reasoned that high pH and alkalinity altered the rhizosphere's physicochemical environment, delaying tuber formation. The low concentration of potatoes might also be attributed to the detrimental effects of salt in the soil solution. The beneficial effects are attributed to improved rhizosphere conditions, better root development and reduced surface crusting caused by carbonate and bicarbonate precipitation (101).

Sulphur

Elemental sulfur and pyrite can be used to reclaim sodic soils, but their effectiveness depends on their complete oxidation to produce sulfuric acid. Sulfur must be oxidized to generate enough sulfuric acid to replace exchangeable Na^+ for it to function

as effectively as soluble calcium ions. Therefore, as compared to gypsum or sulfuric acid, sulfur does not yield the optimal outcomes, even in chemically equivalent concentrations (85).

Industrial by-products

Phosphogypsum

Phosphogypsum (PG), an acidic by-product of phosphoric acid production from rock phosphate, contains essential nutrients such as sulfur (S) and calcium (Ca), along with trace contaminants and heavy metals. It has been shown to mitigate subsoil constraints including acidity, aluminium (Al) toxicity, limited nutrient availability and sodicity (102). The application of powdered PG (5 Mg ha^{-1}) greatly increased IR in a sodic soil ($\text{ESP} = 21$) by inhibiting crust formation; it increased from 0.9 mm h⁻¹ in the control to 8.3 mm h⁻¹ (105).

According to the study, applying PG and CaCl_2 with canal water successfully decreased the amount of salt in saline-sodic soil. Total Na^+ and soluble salts were eliminated 90 % by CaCl_2 , 79 % by PG and 60 % by PG. In both situations, the soil ESP dropped by around 90 %. Because PG was inexpensive, it was regarded as an effective modification (104).

Fly ash

Large volumes of fly ash (FA), a byproduct of coal and lignite-based thermal power plants, are generated annually. In India, approximately 50 % of this FA is utilized by the cement and concrete industries, while innovative applications for the remainder such as soil restoration are being actively explored (105). In sodic soils, the combined application of gypsum and acidic FA has been shown to reduce the SAR, improve nutrient availability, increase saturated HC and enhance soil water retention. For instance, treatment of sodic soil with this combination (initial soil conditions: $\text{pH} = 9.07$, $\text{EC} = 3.87 \text{ dS m}^{-1}$, Exchangeable Sodium Percentage (ESP) = 26) resulted in significantly higher rice and wheat yields compared to untreated control plots (106). These findings suggest that FA can effectively substitute up to 40 % of the gypsum requirement (GR), offering a sustainable and cost-effective approach for sodic soil reclamation.

Press mud

A by-product of sugar mills, press mud (PM) increases soil surface area (SAR) over untreated soil since it is low in irrigation and high in plant nutrients. Plant growth and biomass output have been reported to increase, which may be explained by improvements in N, P and K contents and decreases in Na^+ in maize roots and shoots (107). When combined with gypsum, PM significantly enhanced zinc (Zn) uptake and rice yields in saline-sodic soils, demonstrating the synergistic effect of organic and chemical amendments (108).

Organic wastes and composts

Organic additives such as mulches and composts enhance the physico-chemical properties of SAS by improving the plants' capacity to exchange cations, hold onto water and access nutrients. Even though these adjustments are less costly than the pricey chemicals (109).

Mulching

Mulching with organic materials and agricultural wastes improves soil organic carbon (OC), promotes the formation of water-stable aggregates and enhances moisture retention. It

protects the root zone by reducing erosion caused by runoff, insulating the soil from heat and minimizing water evaporation (110). Straw mulching in brackish water (3-5 g L⁻¹) When compared to treatments without mulch, irrigated wheat-summer maize rotation soils dramatically reduced salt buildup (111). When compared to severely saline-sodic unmulched soils, a 20-year continuous coverage of saline-sodic soils with tephra mulch that was 10-15 cm thick completely reduced salt danger, as seen by notable decreases in EC se and ESP. Reduced evaporation, enhanced water flow, positive changes in the soil moisture regime and limited upward migration of Na⁺ and other salts were cited as the reasons for these decreases. During the trial, neither soil received any irrigation (112).

Landform and drainage treatments for wet sodic soils

Approximately 2.46 million hectares of land in India are affected by waterlogged areas, which result in crop root zone saturation and limited air and water flow. Soils in areas impacted by salt have both permanent waterlogging, when water tables are within 2 m and seasonal waterlogging, which is brought on by floods, high rains and drainage congestion. Anaerobic conditions, osmotic stress, salinity and nutritional toxicities can result from subsurface waterlogging, which occurs when water tables are within 2 m. The states of Punjab and Uttar Pradesh have significant tracts of such wet sodic soils (113).

Irrigation scheduling and management

Sodic soil frequently has problems with permeability and infiltration, which leads to inadequate use of rainfall and decreased irrigation effectiveness. Prior to irrigation, adequate leveling is necessary to provide a homogeneous soil surface and avoid high water application depth, low efficiency and water inundation (114). For crops like wheat in particular, sodic soils need different irrigation management timing, depth and frequency than regular soils. Despite its moderate salt tolerance and sensitivity to waterlogging, wheat has demonstrated improved grain yields under optimized irrigation in sodic soils (115).

Maintaining reclaimed sodic soils' productivity

The potential to reclaim and manage degraded sodic lands using simple methods is well recognized. However, challenges such as the declining availability and quality of agricultural-grade gypsum hinder widespread adoption, despite its effectiveness in addressing structural and nutrient limitations in sodic soils (116). A study in Etawah, Uttar Pradesh, revealed that 27 % of 3905 acres of reclaimed sodic land had resodified, indicating soil degradation and reduced crop productivity (117). In the Indo-Gangetic Plains (IGP), poor irrigation practices, suboptimal fertilizer use, excessive tillage and climate variability further degrade soil health and crop yields (118). Nevertheless, the implementation of conservation agriculture practices-particularly within the Rice-Wheat Cropping System (RWCS) has shown promise in reducing production costs, mitigating resource degradation and improving overall productivity and profitability.

Conclusion

Sustainable management of saline and sodic soils is essential to maintain vegetable crop production and ensure long-term soil health. Salinity reduces plant growth and yield, but this challenge can be addressed through the development of salt-tolerant varieties, traditional breeding, biotechnology and grafting of tolerant rootstocks. Although many vegetables like okra, tomato, eggplant, potato and carrot show some level of salt tolerance, further research is needed to classify them accurately based on their tolerance levels. Efficient water management practices, such as using agricultural drainage water for initial soil leaching, can significantly reduce freshwater use. Additionally, the application of gypsum improves soil structure and accelerates salt removal by increasing calcium and magnesium levels and reducing harmful sodium and potassium. Integrating these practices with the cultivation of salt-tolerant vegetables offers a practical and sustainable approach to reclaim and manage saline-sodic soils effectively. This holistic approach not only supports agricultural productivity but also contributes to the resilience and sustainability of agroecosystems in salt-affected regions.

Acknowledgements

We are grateful to Tamil Nadu Agricultural University for their ongoing support in the form of regular workshop that helps us with writing review.

Authors' contributions

JM, KRV & VJ performed conceptualization; SN, AS & KRV has done formal analysis; KRV, NS, AS, SK, PY & VJ conducted supervision and validation; JM & KRV corrected language.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None

References

- Genova C, Weinberger K, Chanthasombath T, Inthalungdsee B, Sanatem K, Somsak K. Postharvest loss in the supply chain for vegetables-The case of *tomato*, *yardlong bean*, *cucumber* and *chili* in Lao PDR. Shanhua: AVRDC (The World Vegetable Center); 2006. Working Paper No. 17.
- Chinnusamy V, Jagendorf A, Zhu JK. Understanding and improving salt tolerance in plants. *Crop Sci.* 2005;45:437-48. <https://doi.org/10.2135/cropsci2005.0437>
- FAO (Food and Agricultural Organization). Global network on integrated soil management for sustainable use of salt affected soils; 2000. Available from: www.fao.org/agl
- Maas EV. Crop. In: Tanji KK, editor. Agricultural salinity assessment and management. New York: ASCE Manuals and Reports on Engineering No. 71; 1990. p. 262-304.
- Rausch T, Kirsch M, Low R, Lehr A, Viereck R, Zhigang A. Salt stress responses of higher plants: the role of proton pumps and Na⁺/H⁺ antiporters. *J Plant Physiol.* 1996;148(3&4):425-33. [https://doi.org/10.1016/S0176-1617\(96\)80275-6](https://doi.org/10.1016/S0176-1617(96)80275-6)

6. West DW, Merrigan IF, Taylor JA. Germination, growth and flowering of *cowpea*, *mung bean* and *soybean* in relation to soil salinity; 1978.

7. Shrivastava P, Kumar R. Soil salinity: a serious environmental issue and plant growth promoting bacteria as one of the tools for its alleviation. *Saudi J Biol Sci.* 2015;22(2):123-31. <https://doi.org/10.1016/j.sjbs.2014.12.001>

8. Jones RA. The development of salt-tolerant *tomatoes*: breeding strategies. *Acta Hortic.* 1986;190:101-14. <https://doi.org/10.17660/ActaHortic.1986.190.8>

9. Cheeseman JM. Mechanisms of salinity tolerance in plants. *Plant Physiol.* 1988;87(3):557-60. <https://doi.org/10.1104/pp.87.3.547>

10. Munns R. Genes and salt tolerance: bringing them together. *New Phytol.* 2005;167(3):645-63. <https://doi.org/10.1111/j.1469-8137.2005.01487.x>

11. Jamil A, Riaz S, Ashraf M, Foolad R. Gene expression profiling of plants under salt stress: a review. *J Plant Sci.* 2011;30(5):435-58. <https://doi.org/10.1080/07352689.2011.605739>

12. Acosta JA, Boris J, Karsten K, Martínez SM. Salinity increases mobility of heavy metals in soils. *Chemosphere.* 2011;85(8):1318-24. <https://doi.org/10.1016/j.chemosphere.2011.07.046>

13. Allen JA, Chambers JL, McKinney D. Intraspecific variation in the response of *Taxodium distichum* seedlings to salinity. *For Ecol Manage.* 1994;70(1&3):203-14. [https://doi.org/10.1016/0378-1127\(94\)90087-6](https://doi.org/10.1016/0378-1127(94)90087-6)

14. Yamaguchi T, Blumwald E. Developing salt-tolerant crop plants: challenges and opportunities. *Trends Plant Sci.* 2005;10(12):615-20. <https://doi.org/10.1016/j.tplants.2005.10.002>

15. Ashraf M, Harris PJC. Photosynthesis under stressful environments: an overview. *Photosynthetica.* 2013;51:163-90. <https://doi.org/10.1007/s11099-013-0021-6>

16. Waisel Y. *Biology of halophytes*. Elsevier; 2012. ISBN 0323151582.

17. Glick BR, Cheng Z, Czarny J, Duan J. Promotion of plant growth by ACC deaminase-producing soil bacteria. *Eur J Plant Pathol.* 2007;119:329-39. <https://doi.org/10.1007/s10658-007-9162-4>

18. Blaylock AD. Soil salinity, salt tolerance and growth potential of horticultural and landscape plants. Laramie, Wyoming: Cooperative Extension Service, University of Wyoming, Department of Plant, Soil and Insect Sciences, College of Agriculture; 1994.

19. Devi ND, Arumugam T. Salinity tolerance in vegetable crops: a review. *J Pharmacogn Phytochem.* 2019;8(3):2717-21.

20. FAO (Food and Agricultural Organization). Salt affected soils and their management. FAO Soils Bulletin 39. Rome: FAO; 1988.

21. USSLS (United States Salinity Laboratory Staff). Diagnosis and improvement of saline and alkali soils. Agriculture Handbook 60. Washington (DC): USDA; 1954.

22. Chourasia KN, More SJ, Kumar A, Kumar D, Singh B, Bhardwaj V, et al. Salinity responses and tolerance mechanisms in underground vegetable crops: an integrative review. *Planta.* 2022;255(3):68. <https://doi.org/10.1007/s00425-022-03845-y>

23. Fang S, Hou X, Liang X. Response mechanisms of plants under saline-alkali stress. *Front Plant Sci.* 2021;12:1049.

24. Shams M, Yildirim E, Ekinci M, Turan M, Dursun A, Parlakova F, et al. Exogenously applied glycine betaine regulates some chemical characteristics and antioxidative defence systems in *lettuce* under salt stress. *Hort Environ Biotechnol.* 2016;57(3):225-31. <https://doi.org/10.1007/s13580-016-0021-0>

25. Shah AN, Tanveer M, Abbas A, Fahad S, Baloch MS, Ahmad MI, et al. Targeting salt stress coping mechanisms for stress tolerance in *Brassica*: a research perspective. *Plant Physiol Biochem.* 2021;158:53-64. <https://doi.org/10.1016/j.plaphy.2020.11.044>

26. Machado RMA, Serralheiro RP. Soil salinity: effect on vegetable crop growth. Management practices to prevent and mitigate soil salinization. *Horticulturae.* 2017;3(2):30. <https://doi.org/10.3390/horticulturae3020030>

27. Giordano M, Petropoulos SA, Cirillo C, Roushanel Y. Biochemical, physiological and molecular aspects of ornamental plants adaptation to deficit irrigation. *Horticulturae.* 2021;7(5):107. <https://doi.org/10.3390/horticultae7050107>

28. Dong X, Sun L, Guo J, Liu L, Han G, Wang B. Exogenous boron alleviates growth inhibition by NaCl stress by reducing Cl uptake in *Beta vulgaris*. *Plant Soil.* 2021;464(1):423-39. <https://doi.org/10.1007/s11104-021-04946-5>

29. Azizi F, Farsaraei S, Moghaddam M. Application of exogenous ascorbic acid modifies growth and pigment content of *Calendula officinalis* L. flower heads of plants exposed to NaCl stress. *J Soil Sci Plant Nutr.* 2021;21(4):2803-14. <https://doi.org/10.1007/s42729-021-00567-0>

30. Shams M, Yildirim E. Variations in response of CaPAO and CaATG8c genes, hormone, photosynthesis and antioxidative system in *pepper* genotypes under salinity stress. *Sci Hort.* 2021;282:110041. <https://doi.org/10.1016/j.scienta.2021.110041>

31. Dadashoglu E, Ekinci M, Kul R, Shams M, Turan M, Yildirim E. Nitric oxide enhances salt tolerance through regulating antioxidant enzyme activity and nutrient uptake in *pea*. *Legume Res.* 2021;44(1):41-5. <https://doi.org/10.18805/LR-540>

32. Khalil R, Elsayed N, Khan TA, Yusuf M. Potassium: a potent modulator of plant responses under changing environment. In: *Role of Potassium in Abiotic Stress*. Springer; 2022. p. 221-47.

33. Nieves-Cordones M, Azeem F, Long Y, Boeglin M, Duby G, Mouline K, et al. Non-autonomous stomatal control by pavement cell turgor via the K⁺ channel subunit AtKC1. *Plant Cell.* 2022;34(5):2019-37. <https://doi.org/10.1093/plcell/koac038>

34. Yang X, Han Y, Hao J, Qin X, Liu C, Fan S. Exogenous spermidine enhances the photosynthesis and ultrastructure of *lettuce* seedlings under high-temperature stress. *Sci Hort.* 2022;291:110570. <https://doi.org/10.1016/j.scienta.2021.110570>

35. Ansari HH, Siddiqui A, Wajid D, Tabassum S, Umar M, Siddiqui ZS. Profiling of energy compartmentalization in photosystem II (PSII), light harvesting complexes and specific energy fluxes of primed *maize* cultivar (P1429) under salt stress environment. *Plant Physiol Biochem.* 2022;170:296-306. <https://doi.org/10.1016/j.plaphy.2021.12.015>

36. Dhokane K, Pandey J, Yadav RM, Ramachandran P, Rath JR, Subramanyam R. Change in the photochemical and structural organization of thylakoids from *pea* (*Pisum sativum*) under salt stress. *Plant Physiol Biochem.* 2022;177:46-60. <https://doi.org/10.1016/j.plaphy.2022.02.004>

37. Shams M, Ekinci M, Ors S, Turan M, Agar G, Kul R, et al. Nitric oxide mitigates salt stress effects of *pepper* seedlings by altering nutrient uptake, enzyme activity and osmolyte accumulation. *Physiol Mol Biol Plants.* 2019;25(5):1149-61. <https://doi.org/10.1007/s12298-019-00692-2>

38. Ranjan A, Sinha R, Sharma TR, Pattanayak A, Singh AK. Alleviating aluminum toxicity in plants: implications of reactive oxygen species signaling and crosstalk with other signaling pathways. *Physiol Plant.* 2021;173(4):1765-84. <https://doi.org/10.1111/ppl.13382>

39. Sahin U, Ekinci M, Ors S, Turan M, Yildiz S, Yildirim E. Effects of individual and combined effects of salinity and drought on physiological, nutritional and biochemical properties of *cabbage* (*Brassica oleracea* var. *capitata*). *Sci Hort.* 2018;240:196-204. <https://doi.org/10.1016/j.scienta.2018.06.016>

40. Moradbeigi H, Jamei R, Heidari R, Darvishzadeh R. Investigating the enzymatic and non-enzymatic antioxidant defense by applying iron oxide nanoparticles in *Dracocephalum moldavica* L. plant under salinity stress. *Sci Hort.* 2020;272:109537. <https://doi.org/10.1016/j.scienta.2020.109537>

41. Konrad M. *Principles of plant nutrition*. 5th ed. Giessen, Germany: Justus Liebig University, School of Biology; 2001.

42. Sun B, Zhou S, Zhao Q. Evaluation of spatial and temporal changes of soil quality based on geostatistical analysis in sub-tropical China. *Geoderma*. 2003;115(6):85-99. [https://doi.org/10.1016/S0016-7061\(03\)00078-8](https://doi.org/10.1016/S0016-7061(03)00078-8)

43. Fayed I, Rateb K. Long-term cropping system impacts on some physical and chemical properties and fertility status in alluvial soils of Egypt. *Soil Water Environ*. 2013;58(1):9-18.

44. Qi B, Darilek J, Huang B, Zhao Y, Sun W. Evaluating soil quality indices in an agricultural region of Jiangsu Province of China. *Geoderma*. 2009;149(3-4):325-34. <https://doi.org/10.1016/j.geoderma.2008.12.015>

45. Molatakgosi G. Impact of in-field irrigation management by *Botswana cabbage* farmers on soil salinity [Dissertation]. Cranfield University, UK; 2005.

46. Mali B, Thengal S, Pate N. Physico-chemical characteristics of salt affected soil from Barhanpur, M.S., India. *Ann Biol Res*. 2012;3 (8):4091-3.

47. Ghanbari M, Bahmanyar S, Gilani F, Raiesi M. The effect of different levels of saline irrigation water and some amendments on microbial respiration and acid and alkaline phosphatase activities in the rhizosphere soil during vegetative growth of *soybean*. *J Water Soil Conserv*. 2012;19(3):63-76.

48. Rengasamy P. World salinisation with emphasis on Australia. *J Exp Bot*. 2006;57(5):1017-23. <https://doi.org/10.1093/jxb/erj108>

49. Bhadauria S, Sengar M, Dheeraj M, Singh D, Kushwah B. Sustainable land use planning through utilization of alkaline wasteland by biotechnological intervention. *Am Eurasian J Agric Environ Sci*. 2010;9(3):325-37.

50. Alexieva V, Sergiev I, Mapelli S, Karanov E. The effect of drought and ultraviolet radiation on growth and stress markers in *pea* and *wheat*. *Plant Cell Environ*. 2001;24:1337-44. <https://doi.org/10.1046/j.j.1365-3040.2001.00778.x>

51. Rai AC, Rai A, Shah K, Singh M. Engineered BcZAT12 gene mitigates salt stress in tomato seedlings. *Physiol Mol Biol Plants*. 2021;27:535-41. <https://doi.org/10.1007/s12298-021-00948-w>

52. Nahakpam S, Shah K. Expression of key antioxidant enzymes under combined effect of heat and cadmium toxicity in growing rice seedlings. *Plant Growth Regul*. 2011;63:23-35. <https://doi.org/10.1007/s10725-010-9508-3>

53. Flowers TJ, Flowers SA. Why does salinity pose such a difficult problem for plant breeders? *Agric Water Manage*. 2005;78:15-24. <https://doi.org/10.1016/j.agwat.2005.04.015>

54. Sharma P. Amendments in saline soils for improved growth and yield in plants. 2016.

55. Rhoades JD. Electrical conductivity methods for measuring and mapping soil salinity. In: Sparks DL, editor. *Advances in Agronomy*. Vol. 49. San Diego: Academic Press; 1993. p. 201-51.

56. Maas EV. Salt tolerance of plants. *Appl Agric Res*. 1986;1:12-26.

57. Shannon MC, Grieve CM, Francois LE. Whole-plant response to salinity. In: Wilkinson RE, editor. *Plant-environment interactions*. New York: Marcel Dekker; 1994. p. 199-244.

58. Paksoy M, Turkmen O, Dursun A. Effects of potassium and humic acid on emergence, growth and nutrient contents of *Abelmoschus esculentus* L. seedling under saline soil conditions. *Afr J Biotechnol*. 2010;9:5343-6.

59. Abbas W, Ashraf M, Akram NA. Alleviation of stressed-induced adverse effects in *eggplant* (*Solanum melongena* L.) by glycinebetaine and sugarbeet extracts. *Sci Hort*. 2010;125:188-95. <https://doi.org/10.1016/j.scientia.2010.04.008>

60. Tantawy AS, Abdel-Mawgoud AMR, El-Nemr MA, Chamoun YG. Alleviation of salinity effects on *tomato* plants by application of amino acids and growth regulators. *Eur J Sci Res*. 2009;30:484-94.

61. Shahba Z, Baghizadeh A, Ali VSM, Ali Y, Mehdi Y. The salicylic acid effect on the *tomato* (*Lycopersicon esculentum* Mill.) sugar, protein and proline contents under salinity stress (NaCl). *J Biophys Struct Biol*. 2010;2:35-41.

62. Etehadnia M, Schoenau J, Waterer D, Karen T. The effect of CaCl₂ and NaCl salt acclimation in stress tolerance and its potential role in ABA and scion/rootstock-mediated salt stress responses. *Plant Stress*. 2010;4:72-81.

63. Velitchkova M, Fedina L. Response of photosynthesis of *Pisum sativum* to salt stress as affected by methyl jasmonate. *Photosynthetica*. 1998;35:89-97. <https://doi.org/10.1023/A:1006878016556>

64. Yadav BR, Paliwal KV. Response of *cauliflower* to nitrogen and phosphorus fertilization on irrigation with saline waters. *Veg Sci*. 1990;17:1-6.

65. del Amor FM, Cuadra-Crespo P. Alleviation of salinity stress in *broccoli* using foliar urea or methyl-jasmonate: analysis of growth, gas exchange and isotope composition. *Plant Growth Regul*. 2010;63:55-62. <https://doi.org/10.1007/s10725-010-9511-8>

66. Alam MZ, Carpenter-Boggs L, Mitra S, et al. Effect of salinity intrusion on food crops, livestock and fish species at Kalapara Coastal Belt in Bangladesh. *J Food Qual*. 2017;2017:1-23. <https://doi.org/10.1155/2017/2045157>

67. Gebreyess BF, Abayineh A. Water harvesting technologies in semi-arid and arid areas. *J Degrad Min Land Manag*. 2019;7(1):1921-8. <https://doi.org/10.15243/jdmlm.2019.071.1921>

68. Esenov PE, Redjepbaev KR. The reclamation of saline soils. In: Babaev AG, editor. *Desert problems and desertification in Central Asia*. Berlin: Springer; 1999. https://doi.org/10.1007/978-3-642-60128-6_15

69. Kaledhonkar MJ, Meena BL, Sharma PC. Reclamation and nutrient management for salt-affected soils. *Indian J Fert*. 2019;15(5):566-75.

70. Ben GA, Ityel E, Dudley L, Cohen S, Yermiyahu U, Presnov E, et al. Effect of irrigation water salinity on transpiration and leaching requirements: a case study of *bell peppers*. *Agric Water Manage*. 2008;95:587-97. <https://doi.org/10.1016/j.agwat.2007.12.008>

71. Shahid SA, Zaman M, Heng L. Soil salinity: historical perspectives and a world overview of the problem. In: *Guideline for Salinity Assessment, Mitigation and Adaptation Using Nuclear and Related Techniques*. Cham: Springer; 2018. https://doi.org/10.1007/978-3-319-96190-3_2

72. Bar Y, Apelbaum A, Kafkafi U, Goren R. Relationship between chloride and nitrate and its effect on growth and mineral composition of *avocado* and *citrus* plants. *J Plant Nutr*. 1997;20 (6):715-31. <https://doi.org/10.1080/01904169709365288>

73. Karaivazoglou NA, Papakosta DK, Divanidis S. Effect of chloride in irrigation water and form of nitrogen fertilizer on Virginia (flue-cured) *tobacco*. *Field Crops Res*. 2005;92(1):61-74. <https://doi.org/10.1016/j.fcr.2004.09.006>

74. Zhang QT, Inoue M, Inosako K, Irshad M, Kondo K, Qio GY, Wang SP. Ameliorative effect of mulching on water use efficiency of *Swiss chard* and salt accumulation under saline irrigation. *J Food Agric Environ*. 2008;6:480-5.

75. Rubio JS, Garcia-Sanchez F, Rubio F, Martinez V. Yield, blossom end rot incidence and fruit quality in *pepper* plants under moderate salinity are affected by K⁺ and Ca²⁺ fertilization. *Sci Hort*. 2009;119:79-87. <https://doi.org/10.1016/j.scientia.2008.07.009>

76. De Oliveira Oliveira FRAFAD, Medeiros JFD, Sousa FLD, Freir AG. Phosphorus-salinity interaction in *radish*. *Rev Cienc Agron*. 2010;41:519-26.

77. Rausch T, Wachter A. Sulfur metabolism: a versatile platform for launching defense operations. *Trends Plant Sci*. 2005;10:503-9. <https://doi.org/10.1016/j.tplants.2005.08.006>

78. An P, Inanaga S, Lux A, Li XJ, Enji, Zhu NW. Interactive effects of salinity and air humidity on two *tomato* cultivars differing in salt tolerance. *J Plant Nutr*. 2005;28:459-73.

79. Edelstein M, Plaut Z, Ben-Hur M. Mechanism responsible for restricted boron concentration in plant shoots grafted on *pumpkin* rootstocks. *Isr J Plant Sci*. 2011;59:207-15. <https://doi.org/10.1560/IJPS.59.2-4.207>

80. Sivritepe HO, Sivritepe N, Eris A, Turhan E. The effect of NaCl pre-

treatments on salt tolerance of *melon* grown under long-term salinity. *Sci Hort.* 2005;106:568-81. <https://doi.org/10.1016/j.scienta.2005.05.011>

81. Elwan MWM. Ameliorative effects of di-potassium hydrogen orthophosphate on salt-stressed eggplants. *J Plant Nutr.* 2010;33:1593-604. <https://doi.org/10.1080/01904167.2010.496884>
82. Takagi M, El-Shemih H, Sasaki S, Toyama S, Kanai S, Saneoka H, et al. Elevated CO₂ concentration alleviates salinity stress in *tomato* plants. *Acta Agric Scand B Soil Plant Sci.* 2009;59:87-96. <https://doi.org/10.1080/09064710801932425>
83. Jobbágy EG, Jackson RB. Groundwater and soil chemical changes under phreatophytic tree plantations. *J Geophys Res.* 2007;112 (G2). <https://doi.org/10.1029/2006jg000246>
84. Abrol IP, Yadav JSP, Massoud FI. Salt-affected soils and their management. *FAO Soils Bull.* 1988;39.
85. Singh G, Bundela DS, Sethi M, Lal K, Kamra SK. Remote sensing and geographic information system for appraisal of salt-affected soils in India. *J Environ Qual.* 2010;39:5-15. <https://doi.org/10.2134/jeq2009.0032>
86. Galvani A. The challenge of the food sufficiency through salt-tolerant crops. *Rev Environ Sci Biotechnol.* 2006;6(1-3):3-16. <https://doi.org/10.1007/s11157-006-0010-3>
87. Brady N, Weil R. The nature and properties of soils. 13th ed. USA: Macmillan; 2002.
88. Liluchli A, Epstein E. Plant responses to saline and sodic conditions. In: Tanji KK, editor. *Agricultural salinity assessment and management*. ASCE Manuals Reports Eng Pract No. 71. New York: ASCE; 1996. p. 113-37.
89. Rengasamy P. World salinization with emphasis on Australia. *J Exp Bot.* 2006;57(5):1017-23. <https://doi.org/10.1093/jxb/erj108>
90. Pal DK, Srivastava P, Durge SL, Bhattacharyya T. Role of microtopography in the formation of sodic soils in the semi-arid part of the Indo-Gangetic Plains, India. 2003.
91. Chhabra R. Classification of salt-affected soils. *Arid Land Res Manag.* 2004;19(1):61-79. <https://doi.org/10.1080/15324980590887344>
92. Abrol IP, Bhumbra DR. Crop response to differential gypsum applications in a highly sodic soil and the tolerance of several crops to exchangeable sodium under field conditions. *Soil Sci.* 1979;127(2):79-85.
93. Jumberi A, Oka M, Fujiyama H. Response of vegetable crops to salinity and sodicity in relation to ionic balance and ability to absorb microelements. *Soil Sci Plant Nutr.* 2002;48(2):203-9. <https://doi.org/10.1080/00380768.2002.10409192>
94. Phogat V, Sharma SK, Kumar S, Satyavan, Gupta SK. Vegetable cultivation with poor quality water. *Tech Bull.* 2010;72. Department of Soil Science, CCS Haryana Agricultural University, Hisar.
95. Singh J, Ashok V, Chauhan N. Effect of pyrite on growth and yield of *pearl millet-wheat* cropping system under sodic environment. *J Agric Phys.* 2009;9:24-7.
96. Swarup A. Effect of gypsum, green manure, farmyard manure and zinc fertilization on the zinc, iron and manganese nutrition of *wetland rice* on a sodic soil. *J Indian Soc Soil Sci.* 1982;39:530-6.
97. Sharma DK, Singh A. Salinity research in India: achievements, challenges and future prospects. *Water Energy Int.* 2015;58:35-45.
98. Yunusa IAM, Eamus D, DeSilva DL, Murray BR, Burchett MD, Skilbeck GC, et al. Fly-ash: an exploitable resource for management of Australian agricultural soils. *Fuel.* 2006;85:2337-44. <https://doi.org/10.1016/j.fuel.2006.01.033>
99. Chen L, Dick WA. Gypsum as an agricultural amendment: general use guidelines. The Ohio State University; 2011.
100. Tyagi NK. Managing saline and alkali water for higher productivity. In: Kijne JW, Barker R, Molden D, editors. *Water productivity in agriculture: limits and opportunities for improvement*. Wallingford: CABI Publishing; 2003. p. 69-87.

<https://doi.org/10.1079/9780851996691.0069>

101. Yadav AC, Sharma SK, Kapoor AK, Singh A, Mangal JL. Response of *potato* to sodic water irrigation with and without amendments. *Haryana J Hort Sci.* 2002;31(1&2):129-32.
102. Alcordo IS, Rechcigl JE. Phosphogypsum in agriculture: a review. *Adv Agron.* 1993;49:55-118. [https://doi.org/10.1016/S0065-2113\(08\)60793-2](https://doi.org/10.1016/S0065-2113(08)60793-2)
103. Agassi M, Shainberg I, Morin J. Infiltration and runoff in *wheat* fields in the semi-arid region of Israel. *Geoderma.* 1985;36:263-76. [https://doi.org/10.1016/0016-7061\(85\)90007-2](https://doi.org/10.1016/0016-7061(85)90007-2)
104. Kaushik A, Nisha R, Jagjeet K, Kaushik CP. Impact of long- and short-term irrigation of a sodic soil with distillery effluent in combination with bio amendments. *Bioresour Technol.* 2005;96:1860-6. <https://doi.org/10.1016/j.biortech.2005.01.031>
105. Chatterjee AK. Indian fly ashes: their characteristics and potential for mechanochemical activation for enhanced usability. *J Mater Civ Eng.* 2011;23:783-8. [https://doi.org/10.1061/\(ASCE\)MT.1943-5533.0000279](https://doi.org/10.1061/(ASCE)MT.1943-5533.0000279)
106. Kumar D, Singh B. The use of coal fly ash in sodic soil reclamation. *Land Degrad Dev.* 2003;14:285-99. <https://doi.org/10.1002/ldr.557>
107. Muhammad D, Khattak RA. Growth and nutrient concentrations of *maize* in pressmud-treated saline-sodic soils. *Soil Environ.* 2009;28:145-55.
108. Chand M, Randhawa NS, Sinha MK. Effect of gypsum, pressmud, fulvic acid and zinc sources on yield and zinc uptake by *rice* crop in a saline-sodic soil. *Plant Soil.* 1980;55:17-24. <https://doi.org/10.1007/BF02149704>
109. Lakhdar A, Rabhi M, Ghnaya T, Montemurro F, Jedidi N, Abdelly C. Effectiveness of compost use in salt-affected soil. *J Hazard Mater.* 2009;171:29-37. <https://doi.org/10.1016/j.jhazmat.2009.05.132>
110. Mulumba LN, Lal R. Mulching effects on selected soil physical properties. *Soil Tillage Res.* 2008;98:106-11. <https://doi.org/10.1016/j.still.2007.10.011>
111. Pang HC, Li YY, Yang JS, Liang YS. Effect of brackish water irrigation and straw mulching on soil salinity and crop yields under monsoonal climatic conditions. *Agric Water Manage.* 2010;97:1971-7. <https://doi.org/10.1016/j.agwat.2009.08.020>
112. Tejedor M, Jiménez CC, Díaz F. Use of volcanic mulch to rehabilitate saline-sodic soils. *Soil Sci Soc Am J.* 2003;67:1856-61. <https://doi.org/10.2136/sssaj2003.1856>
113. Sharma DB, Swarup A. Effects of short-term flooding on growth, yield and mineral composition of *wheat* on sodic soil under field conditions. *Plant Soil.* 1988;107:137-43. <https://doi.org/10.1007/BF02371555>
114. Tyagi NK. Effect of land surface uniformity on some economic parameters of irrigation in sodic soils under reclamation. *Irrig Sci.* 1984;5:151-60. <https://doi.org/10.1007/BF00264604>
115. Sharma DK, Kumar A, Singh KN. Effect of irrigation scheduling on growth, yield and evapotranspiration of *wheat* in sodic soils. *Agric Water Manage.* 1990;18:267-76. [https://doi.org/10.1016/0378-3774\(90\)90048-4](https://doi.org/10.1016/0378-3774(90)90048-4)
116. Sharma PC, Thimappa K, Kaledhonkar MJ, Chaudhari SK. Reclamation of alkali soils through gypsum technology. ICAR-CSSRI/Karnal/Technology Folder/2016/01. 2016. p. 4.
117. Yadav MS, Yadav PPS, Yaduvanshi M, Verma D, Singh AN. Sustainability assessment of sodic land reclamation using remote sensing and GIS. *J Indian Soc Remote Sens.* 2010;38:269-78. <https://doi.org/10.1007/s12524-010-0017-7>
118. Gathala MK, Kumar V, Sharma PC, Saharawat YS, Jat HS, Singh M, et al. Optimizing intensive cereal-based cropping systems addressing current and future drivers of agricultural change in the northwestern Indo-Gangetic Plains of India. *Agric Ecosyst Environ.* 2013;177: 85-97. <https://doi.org/10.1016/j.agee.2013.06.002>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc

See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.