



RESEARCH ARTICLE

Analysis of genetic variability and trait association in rice (*Oryza sativa* L.)

S Thirumalraj¹, J Bonipas Antony², R P Gnanamalar³, S Manonmani⁴, C Gopalakrishnan⁴, M Raveendran¹ & J Ramalingam^{1*}

¹Department of Plant Biotechnology, Centre for Plant Molecular Biology and Biotechnology, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

²Department of Genetics and Plant Breeding, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

³Department of Plant Breeding and Genetics, Tamil Nadu Agricultural University, Madurai 625 104, Tamil Nadu, India

⁴Department of Rice, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

*Correspondence email - ramalingam.j@tnau.ac.in

Received: 05 May 2025; Accepted: 24 July 2025; Available online: Version 1.0: 11 September 2025; Version 2.0: 06 October 2025

Cite this article: Thirumalraj S, Bonipas AJ, Gnanamalar RP, Manonmani S, Gopalakrishnan C, Raveendran M, Ramalingam J. Analysis of genetic variability and trait association in rice (*Oryza sativa* L.). Plant Science Today. 2025;12(sp1):01–08. <https://doi.org/10.14719/pst.9309>

Abstract

Rice is a staple food for billions of people worldwide. In recent years unpredictable weather patterns, making rice cultivation more challenging and threatens its productivity. To ensure food security, developing high-yielding and climate-resilient rice varieties is essential. This requires a deep understanding of genetic variability and the relationships between key agronomic traits. Thus, the present study explores the genetic variability, as well as the major yield attributes of a segregating population derived from black rice genotypes, to formulate an effective selection strategy. In this research, 188 F₂ plants derived from crosses between the improved *Chakhao Amubi* and improved *Co 51* varieties were analysed and superior progenies were forwarded to F₃ generation. The variability study in the F₂ population revealed substantial variation for single plant yield and related yield attributes such as productive tillers and panicle length. High heritability estimates across all traits, along with a significant genetic advance over the mean for yield and major yield components, indicate that phenotypic selection would be effective. Principal Component Analysis (PCA) further confirmed the extensive variability present in the F₂ population. Correlation and path analysis highlighted panicle length and flag leaf length as key contributors to single plant yield, emphasizing their importance in selection strategies. The combined analysis of the F₂ and selected F₃ families revealed that single plant yield is predominantly governed by additive gene action, as evidenced by significant inter-generational correlation and high narrow-sense heritability. Furthermore, the observed positive selection response for yield suggests that simple phenotypic selection can lead to substantial genetic gain in rice

Keywords: black rice; intergeneration correlation; principal component analysis; selection gain

Introduction

Black rice (*Oryza sativa* L. indica) is a special cultivar of rice which contains remarkably high anthocyanin pigments in the aleurone layer than white and red rice which account for its violet or dark purple colour (1). Black rice is locally known as 'Chakhao', means delicious rice in Manipuri language is cultivated mainly by Meitei farmers of Manipur. Black rice is packed with nutrients and health's benefits, making it a standout among rice varieties. It has nearly six times more antioxidant activity than other types of rice and boasts a high protein content (8.16 %) while being extremely low in fat (0.07 %) (2). Naturally gluten-free and gentle on the gut, it also acts as a natural detoxifier with various medicinal properties (3). This nutrient-rich rice contains essential amino acids like lysine and tryptophan, along with functional lipids, dietary fibre and vitamin B1, vitamins B2, vitamins E and folic acid. It also has powerful phenolic compounds, including γ-oryzanol, tocopherols and tocotrienols, which contribute to its health benefits. When cooked, black rice has a mildly nutty

flavour and a slightly sticky texture. It's loaded with essential macro and micronutrients like iron, zinc, calcium, phosphorus and selenium, yet remains low in calories (4).

Among the rice varieties grown in Northeast India, black rice stands out for its superior nutritional value and high protein content. Though, its cultivation is becoming increasingly difficult due to climate change. It led to rising temperatures, unpredictable rainfall and salt intrusion in coastal areas (5-7). These environmental challenges situate rice production at risk, emphasizing the need for resilient rice varieties and adaptive farming practices to sustain yields.

Grain yield is influenced by genotype, environment and cultural practices, along with their interactions (8). Still, the most significant variations in yield are primarily driven by the interaction between genotype and environment, even under consistent cultural practices. Additionally, planting time plays a crucial role in determining yield outcomes (9).

Differences in yield are evident across different rice genotypes, environmental conditions and their interactions (10). To analyse these variations, multivariate statistical techniques like PCA and cluster analysis are commonly used (11).

PCA helps identify patterns within large datasets by reducing redundancy and highlighting key relationships among variables, making it useful for classifying genotypes. On the other hand, cluster analysis focuses on grouping previously uncategorized materials based on their similarities. PCA allows researchers to visualize relationships between variables and observations, with 'eigenvalues' indicating how much each principal component contributes to overall variance. Meanwhile, eigenvector coefficients show amount variable influences a given component. The main advantage of PCA is its ability to pinpoint the most critical dimensions responsible for variation within a dataset. It is particularly useful when studying plant traits related to yield (12-14). The global population rising and challenges like climate change, shrinking arable land and water scarcity, improving rice productivity and resilience has become a major focus in agricultural research (15).

To enhance genetic improvement through breeding, researchers must understand variability in rice genotypes, their trait associations and the factors affecting yield and quality (16). Advanced statistical tools have proven effective in identifying key traits that contribute to higher productivity and can help guide breeding strategies for better-performing rice varieties.

Materials and Methods

To study genetic variability crosses were made between the improved *Chakhao Amubi* [Drought-tolerant backcross inbred lines of Chakhao Amubi were successfully developed through marker-assisted breeding by introgressing the drought-responsive QTLs *qDTY1.1*, *qDTY2.1* and *qDTY6.1* from the donor parent Apo] and improved *Co51* rice varieties [Disease resistant line with multiple resistant genes for bacterial blight (*xa5*, *xa13* and *Xa21*) and blast (*Pi54*)]. True hybrids in the F_1 generation were identified using polymorphic markers and then advanced to the F_2 generation. These hybrids were self-pollinated to produce the F_2 population of 188 F_2 plants to assess genetic diversity and variability. The F_2 plants with desirable genes (both drought QTLs and biotic stress resistant genes) were forwarded to F_3 and raised in a family row method.

A total of nine quantitative traits were recorded in individual F_2 plants and all the plants in F_3 family rows. Those are, days to fifty percent flowering (DFF, days), plant height (PH, cm), number of productive tillers (NPT, count), panicle length (PL, cm), FLL (cm), flag leaf width (FW, cm), number of grains per panicle (NGP, count), hundred seed weight (HSW, g) and single plant yield (SPY, g).

The experiment was conducted at the Paddy Breeding Station, TNAU and Coimbatore. It is located at an elevation of 426.72 meters at 11°N latitude and 77°E longitude. The station covers 12.96 hectares of cultivable land with clayey soil and a pH of 7.8. The region receives an average annual rainfall of 670 mm, providing controlled conditions for studying trait variation.

Statistical analysis

The collected data were evaluated for genetic variability parameters using MS Excel software. The variability analysis and parent progeny regression were performed according to Singh and Narayanan (17). The PCA was performed using 'FactoMineR' R package and visualized using 'factoextra' R package. The correlation and path analysis were performed using 'agricolea' R package.

Results and Discussion

Genetic variability

The Genotypic Coefficient of Variation (GCV) and Phenotypic Coefficient of Variation (PCV) values showed that SPY had the highest genetic variability, with GCV and PCV recorded at 41.31 % and 43.61 %, respectively. This close proximity between GCV and PCV also indicated low environmental influence and a high degree of heritable variation. High GCV and PCV were also noted for flower length (FL) and pod length (PL), whereas days to 50 % flowering (DFF) and number of green pods (NGP) exhibited the lowest GCV (1.24 % and 2.37 %, respectively), implying limited genetic variability and strong environmental influence. All the studied traits showed low coefficient of variation (CV) which suggesting more precise measurements and lower environmental fluctuation.

Heritability estimates ranged from moderate to high across traits. Flower length (FL) exhibited the highest heritability (0.95), followed closely by SPY (0.90) and hundred seed weight (HSW) (0.91). These high values indicate that the observed variation is largely due to genetic factors and can be effectively transferred to the next generation. Conversely, Number of Pods per Plant (NPT) recorded the lowest heritability (0.45), reflecting higher environmental influence and reduced selection efficiency.

Genetic Advance (GA) and Genetic Advance as Percent of Mean (GAM) results showed that the SPY had the highest GA (22.09) and GAM (80.60 %), reinforcing its suitability for selection in breeding programs. Similarly, high GAM values were recorded for flower length (63.50 %) and pod length (39.01 %), indicating traits governed primarily by additive gene action. Traits such as number of green pods (NGP) and DFF showed low GAM values (3.74 % and 1.94 %, respectively), suggesting limited potential gain through selection (Table 1).

Correlation

The correlation matrix provides valuable insights into the relationships between various agronomic traits in rice, highlighting key interactions that can guide to formulation of breeding strategies. Strong positive correlations, such as between FLL and PL ($r = 0.72$, $p < 0.001$), suggest that an increase in FLL is associated with longer panicles, which may contribute to higher yield potential. These relationships play a crucial role in selecting desirable traits for breeding programs to improve rice productivity (18). For instance, that flag leaf size is directly linked to grain yield, influencing photosynthetic efficiency and assimilate translocation. Similarly highlighted the role of yield contributing traits in determining overall grain production. Understanding such correlations aids in targeted selection and genetic improvement strategies to enhance the

Table 1. Genetic variability parameters for agronomic traits in selected populations

	DFF	PH	NPT	PL	FL	FW	NGP	HSW	SPY
Ve	1.26	13.90	3.55	4.07	3.61	0.00	7.21	0.01	14.67
Vp	2.94	89.57	6.50	23.78	70.64	0.01	17.50	0.06	142.79
Vg	1.68	75.67	2.95	19.71	67.02	0.01	10.29	0.05	128.12
Mean	104.26	110.56	12.22	21.34	25.87	1.21	135.36	2.44	27.40
GCV	1.24	7.87	14.06	20.80	31.64	6.53	2.37	9.44	41.31
PCV	1.64	8.56	20.87	22.85	32.49	7.60	3.09	9.88	43.61
CV	1.08	3.37	15.42	9.45	7.35	3.90	1.98	2.90	13.97
heritability	0.57	0.84	0.45	0.83	0.95	0.74	0.59	0.91	0.90
GA	2.02	16.47	2.38	8.33	16.43	0.14	5.07	0.45	22.09
GAM	1.94	14.90	19.51	39.01	63.50	11.54	3.74	18.59	80.60
Max	109	141	21	32	44	1.6	147	3.81	67
Min	100	55.7	7	12	2	1.1	121	2.1	7

yield and adaptability of rice varieties (19). The study reveals negative correlations between PL and tillering capacity ($r = -0.17$, $p < 0.05$), suggesting a trade-off between these factors (20). These findings have implications for rice breeding, as traits like FL, PL and FW can be prioritized for yield improvement. The negative correlation between PL and tiller number highlights the complexity of genetic interactions and the need for multi-trait selection strategies (21, 22).

The correlation coefficient measures the strength and direction of the relationship between two traits, making it a valuable tool for selecting and improving correlated traits simultaneously (23). The study indicates a strong positive correlation between traits like plant height, flag leaf size, leaf width and grain yield per plant (24). Additionally, panicle length and other factors like total tillers, effective tillers, PL, grains per panicle, fertile grains per panicle, 1000 grain weight and kernel breadth also contribute positively significantly to grain yield. These findings suggest these traits are valuable targets for improving grain yield in crop breeding (25-31). Grain yield per plant is negatively correlated with days to fifty percent flowering reported by (32). Spikelet fertility has a negative correlation with grain yield (33) and 1000 grain weight has a negative correlation (Fig. 1).

Path analysis

Path coefficient analysis was used to determine the importance of different traits in improving yield (34). Grain yield per plant is negatively correlated with days to flowering, while traits like spikelet fertility, kernel length and L/B ratio show a positive but non-significant correlation. Spikelet fertility has a negative correlation with grain yield, while 1000 grain weight has a negative correlation. Path coefficient analysis was used to determine the importance of different traits in improving yield (35) (Table 2).

Principal Component Analysis (PCA)

PCA was employed to assess the extent to which individual agronomic traits contributed to the total variability within the dataset. The scree plot shows the percentage of explained variance for each principal component (PC), helping determine the optimal number of PCs to retain for analysis (Fig. 2). The first three principal components (PC1, PC2 and PC3) together explained 60 % of the total variation, with PC1 alone accounting for 31 %. Traits like single plant yield, FLL and PL had the strongest influence on PC1, suggesting they play a key role in overall trait variation (36). Plant height and the number of grains per panicle were most important in PC2, while the number of productive tillers and 100-seed weight stood out in



**Fig. 1.** Heatmap of Pearson's correlation coefficients among agronomic traits.

Table 2. Path coefficient analysis indicating direct and indirect effects of agronomic traits

	DFF	PH	NPT	PL	FL	FW	NGP	HSW
DFF	-0.0183	-0.0002	0.0009	0.0094	0.0063	0.0052	0.0017	-0.005
PH	-0.0007	-0.0039	-0.0032	-0.0531	-0.0696	0.0045	0.0091	-0.0032
NPT	-0.0005	0.0004	0.0291	0.0094	0.0759	0.0091	-0.0061	-0.0072
PL	-0.0005	0.0007	0.0009	0.3123	0.3857	0.0181	0.0065	-0.0036
FL	-0.0002	0.0004	0.0035	0.1905	0.6323	0.0227	0.0026	-0.0018
FW	-0.0015	-0.0003	0.0041	0.0874	0.2213	0.0648	0.0074	-0.0032
NGP	-0.0007	-0.0008	-0.0041	0.0468	0.0379	0.011	0.0434	-0.0036
HSW	-0.0026	-0.0003	0.0058	0.0312	0.0316	0.0058	0.0043	-0.0359
Residual	0.205228							

**Fig. 2.** Scree plot showing the percentage of explained variance across principal components.

PC3. Interestingly, days to 50 % flowering had a strong negative impact on PC4. By the time six components are considered, nearly 87 % of the variability is captured, meaning most of the important differences among traits can be understood with just a few key patterns. This makes it easier to focus on the traits that matter most in breeding and selection efforts.

However, Retaining PC1 and PC2 is a reasonable approach for dimensionality reduction and trait selection in rice breeding programs, as they capture the most critical variations among agronomic traits (37). Thus, PCA biplot generated using first two PCs. Principal components with eigenvalues greater than 1.0 indicate high variability among rice genotypes, making them useful for selecting diverse parental lines. Similar findings were reported by (38). Recent genetic analysis highlights key parameters shaping trait inheritance and selection effectiveness. Traits like seed yield per plant and number of productive tillers show high selection gains (46.45 and 75.02, respectively), indicating potential for genetic improvement. Conversely, days to 50 % flowering and flag leaf length show negative selection gains (-4.11 and -20.43), suggesting a generational decline in these traits (39).

The study reveals stable inheritance patterns for FLL (0.74) and hundred-seed weight (HSW) (0.76), as well as strong positive correlations at $p < 0.01$, for SPY (0.61) and NPT (0.67), DFF (72.83) are showed high inheritable, while PL (9.06) and FW (10.66) show lower heritability, suggesting stronger environmental influence (40). These findings suggest that three traits should be prioritized in breeding programs for sustained

genetic improvement. The F_3 families show significant variation in key agronomic traits, such as days to DFF, PH, NPT, PL, FLL and SPY providing valuable insights for breeding programs (41).

PC1 explains the highest variance, accounting for over 30 %, followed by PC2 which contributes around 20 %. A sharp decline in variance is observed after PC2, indicating that the first two PCs capture a substantial proportion of the total variance in the dataset. This pattern aligns with the Kaiser criterion. The gradual decrease in explained variance across subsequent PCs indicates diminishing contributions to overall data variation (42-43). A scree plot shows the percentage of explained variance for each principal component, with PC1 exhibiting the highest variability.

The PCA results reveal that FL, SPY and PL are the most influential in explaining variability (8), while path analysis confirms FL and SPY as key contributors to productivity, emphasizing their role in breeding programs. Additionally, the selection gain for SPY (+12.71) indicates promising yield improvement, further supporting its importance in genetic enhancement strategies (44). High heritability traits (FL, SPY, NGP) indicate strong genetic control, making them ideal for selection (45). Similarly, high GAM values for SPY, FL and PL suggest significant improvement potential under selection pressure. CA simplified the analysis of agronomic traits by identifying key contributors to variation in rice. The first three components explained 60 % of the total variance, with PC1 (31 %) driven by traits like single plant yield, flag leaf length and panicle length-highlighting their importance for yield improvement. PC2 and PC3 emphasized structural and

reproductive traits, while days to 50 % flowering showed an independent trend in PC4. The scree plot supported focusing on the top components, suggesting that a few key traits can effectively guide selection and breeding decisions. The PCA results reveal that FL, SPY and PL are the most influential in explaining variability (46), while path analysis confirms FL and SPY as key contributors to productivity, emphasizing their role in breeding programs. Additionally, the selection gain for SPY (+12.71) indicates promising yield improvement, further supporting its importance in genetic enhancement strategies (47) (Table 3).

Genetic studies in F₃

Family 143, 151 and 141 of the F₃ family showed the highest SPY (45.55), (44.85) and (44.72) respectively indicating superior yield potential. Panicle length varied from 22.20 cm (F₃-169) to 24.99 cm (F₃-143) and flag leaf length showed significant genetic variability, with F₃-58 having the longest FLL (24.58 cm) and F₃-56 having the shortest (16.69 cm). FDFR ranging from 96.57 days (F₃-151) to 103.35 days (F₃-74), indicating genetic differences in growth duration. Plant height also showed considerable variation, with F₃-56 reaching the tallest height (118.84 cm) and F₃-151 the shortest (103.24 cm), pointing to differences in plant architecture (Table 4). These variations in yield-related traits highlight the genetic diversity within the F₃ families, making entries like 143, 151 and 141 strong candidates for further breeding selection (48).

Among the selected, families 143, 151 and 141 exhibited the highest seed yield per plant, making them strong candidates for future breeding efforts. Additionally, differences in panicle length, flag leaf length, flowering time and plant height demonstrate the genetic variability within these families, offering breeders a wide selection pool for trait improvement (49). Interestingly, while traits like seed yield per plant (SPY) and the NPT show high selection gains, others such as days to 50 % flowering (DFF) and FLL exhibit negative selection gains, indicating a potential decline over generations. This suggests that while some traits are strongly inheritable and stable, others may require careful selection to prevent undesirable changes over time (50, 51). Overall, these findings highlight the importance of multi-trait selection strategies in rice breeding (Table 5). By prioritizing high-yielding families, maintaining genetic diversity and balancing trade-offs between correlated traits, breeders can enhance rice productivity while ensuring long-term genetic stability. High heritability traits (FL, SPY, NGP) indicate strong genetic control, making them ideal for selection (52). Similarly, high GAM values for SPY, FL and PL suggest significant improvement potential under selection pressure (53, 54).

Conclusion

Table 3. Principal component contributions to agronomic trait variability

	PC1	PC2	PC3	PC4	PC5	PC6	PC7	PC8	PC9
DFF	0.04	0.28	-0.34	-0.84	0.24	-0.01	0.17	-0.03	-0.01
PH	-0.09	0.60	0.12	0.29	0.28	-0.61	0.24	0.16	0.01
NPT	0.11	-0.15	-0.67	0.38	0.18	0.21	0.53	0.11	0.03
PL	0.50	-0.05	0.11	-0.13	-0.24	-0.05	0.03	0.78	0.22
FL	0.53	-0.10	0.07	0.01	0.03	-0.24	0.10	-0.53	0.59
FW	0.33	0.25	-0.09	0.18	0.59	0.32	-0.58	0.07	0.02
NGP	0.12	0.57	0.28	0.07	-0.27	0.61	0.33	-0.13	0.04
HSW	0.09	0.35	-0.55	0.10	-0.59	-0.18	-0.39	-0.11	-0.04
SPY	0.56	-0.08	0.10	0.00	-0.01	-0.16	0.13	-0.18	-0.77
Standard deviation	1.66	1.17	1.11	0.97	0.91	0.85	0.79	0.61	0.36
proportion of variance	0.31	0.15	0.14	0.10	0.09	0.08	0.07	0.04	0.01
cumulative proportion	0.31	0.46	0.60	0.70	0.79	0.87	0.94	0.99	1.00

Table 4. Selected mean performance F₃ lines

F ₃ family name	DFF	PH	NPT	PL	FLL	FW	NGP	HSW	SPY
2	101.51	116.23	21.74	22.62	22.79	1.17	138.67	2.31	41.03
18	101.91	103.39	20.11	22.41	22.54	1.20	137.10	2.31	37.56
56	98.06	118.84	22.00	22.73	16.69	1.18	137.19	2.31	28.65
58	102.08	108.75	20.84	23.68	24.58	1.22	139.73	2.20	44.53
73	102.60	117.48	21.44	24.26	17.91	1.19	138.72	2.18	37.78
74	103.35	105.84	20.40	22.52	17.01	1.16	137.67	2.19	43.49
141	98.43	106.84	21.51	23.90	18.97	1.18	137.82	2.20	44.72
143	100.15	113.86	21.91	24.99	21.06	1.17	137.61	2.18	45.55
151	96.57	103.24	21.56	22.25	20.69	1.16	136.85	2.17	44.85
169	96.66	109.33	21.66	22.20	19.86	1.17	129.68	2.19	37.37
171	98.57	116.68	22.06	23.98	24.39	1.20	139.51	2.23	35.54

Table 5. Selection response and heritability estimates in F₃ families

	mean F ₂	Mean F ₃	Selection gain	Selection differential	inter generation correlation	parent progeny regression	narrow sense heritability
DFF	104.26	99.97	-4.11	-4.29	-0.02	-0.04	72.83
PH	110.56	110.94	0.35	0.39	0.03	0.02	33.79
NPT	12.22	21.39	75.02	9.17	0.67*	0.13	9.81
PL	21.34	23.23	8.83	1.89	0.19	0.03	9.06
FLL	25.87	20.58	-20.43	-5.29	0.74**	0.24	16.20
FW	1.21	1.18	-2.31	-0.03	-0.07	-0.01	10.66
NGP	135.36	137.28	1.42	1.92	-0.15	-0.10	31.70
HSW	2.44	2.22	-8.95	-0.22	0.76**	0.25	16.12
SPY	27.37	40.08	46.45	12.71	0.61*	0.43	35.08

This study provides meaningful insights into the relationships among key agronomic traits in rice through correlation analysis, path coefficient analysis and PCA. The strong positive association between FLL and PL highlights their potential as reliable traits for selection in breeding programs focused on yield enhancement. In contrast, the negative correlation between PL and tillering capacity points to the need for careful consideration of trade-offs when selecting multiple traits simultaneously. Path analysis helped identify both direct and indirect effects of various traits, revealing the complex roles of spikelet fertility, kernel traits and 1000 grain weight in determining grain yield. The genetic variability and PCA results, indicated that tremendous genetic variation presented in the segregating population, making the selection process more efficient. Genetic variability and heritability observed among F_3 families and genetic correlation of F_2 with F_3 suggested the predominance of additive gene action in single plant yield implying phenotypic selection would be more effective. The mean performance of F_3 families indicated that lines 143, 151 and 141, as potential candidates need to be forwarded further to isolate superior line with improved yield for climate resilience.

Authors' contributions

ST conducted the experiments, collected data, performed statistical analyses and wrote the manuscript. JBA performed statistical analyses, wrote and edited the manuscript. JR conceptualized the experiment, provided materials, reviewed the manuscript and supervised all activities, including field research, data analysis and manuscript preparation. RPG, SM, CG and MR participated in field research and provided technical support for manuscript preparation. All authors have read and agreed to the publication of the final manuscript.

Compliance with ethical standards

Conflict of interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Ethical issues: None

References

- Hou Z, Qin P, Zhang Y, Cui S, Ren G, et al. Identification of anthocyanins isolated from black rice (*Oryza sativa* L.) and their degradation kinetics. *Food Res Int*. 2013;50(2):691–7. <https://doi.org/10.1016/j.foodres.2011.07.037>
- Thomas R, Wan-Nadiah WA, Bhat R, et al. Physiochemical properties, proximate composition and cooking qualities of locally grown and imported rice varieties marketed in Penang, Malaysia. *Int Food Res J*. 2013;20(3). <https://doi.org/10.3136/fstr.20.469>
- Jha P, Das AJ, Deka SC, et al. Optimization of ultrasound and microwave assisted extractions of polyphenols from black rice (*Oryza sativa* cv. Poireton) husk. *J Food Sci Technol*. 2017;54:3847–58. <https://doi.org/10.1007/s13197-017-2832-0>
- Poonia A, Pandey S, et al. Bioactive compounds, nutritional benefits and food applications of black rice: a review. *Nutr Food Sci*. 2021;52(3):466–82.
- Espe MB. Yield gaps and opportunities: exploring the current state of grain yield in US rice production systems. *Univ Calif Davis*. 2017. <http://doi.org/10.1016/j.fcr.2016.07.011>
- Tayade RR, Nguyen TTD. Effective strategies for enhancing tolerance to high-temperature stress in rice during the reproductive and ripening stages. 2018. <https://doi.org/10.9787/PBB.2018.6.1.1>
- Yuan S, Stuart AM, Laborte AG, Rattalino Edreira JI, Dobermann A, Kien LV, et al. Southeast Asia must narrow down the yield gap to continue to be a major rice bowl. *Nat Food*. 2022;3(3):217–26. <https://doi.org/10.1038/s43016-022-00477-z>
- Messina C, Hammer G, Dong Z, Podlich D, Cooper M, et al. Modelling crop improvement in a G×E×M framework via gene-trait -phenotype relationships. In: Sadras VO, Calderini DF, editors. *Crop Physiology: Applications for Genetic Improvement and Agronomy*. Elsevier; 2009:235–65. <http://doi.org/10.1016/B978-0-12-374431-9.00010-4>
- Roy S, Verma BC, Banerjee A, Kumar J, Ray US, Mandal NP, et al. Genetic diversity for drought and low-phosphorus tolerance in rice (*Oryza sativa* L.) varieties and donors adapted to rainfed drought-prone ecologies. *Sci Rep*. 2021;11(1):13671. <https://doi.org/10.1038/s41598-021-93325-2>
- Rana MM. Conservation agriculture for sustainable crop productivity and economic return for the smallholders of Bangladesh: a systematic review. *Turk J Agric Food Sci Technol*. 2023;11(10):2009–15. <https://doi.org/10.24925/turjaf.v11i10.2009-2015.6053>
- Moosavi M, Ranjbar G, Zarrini HN, Gilani A. Correlation between morphological and physiological traits and path analysis of grain yield in rice genotypes under Khuzestan conditions. *Biol Forum*. 2015;7(1):43.
- Mahendran R, Veerabadhiran P, Robin S, Raveendran M. Principal component analysis of rice germplasm accessions under high temperature stress. 2015.
- Maji AT, Shaibu AA. Application of principal component analysis for rice germplasm characterization and evaluation. *J Plant Breed Crop Sci*. 2012;4(6):87–93. <http://doi.org/10.5897/JPBSC11.093>
- Shoba D, Vijayan R, Robin S, Manivannan N, Iyanar K, Arunachalam P, et al. Assessment of genetic diversity in aromatic rice (*Oryza sativa* L.) germplasm using PCA and cluster analysis. *Electron J Plant Breed*. 2019;10(3):1095–104. <http://doi.org/10.5958/0975-928X.2019.00140.6>
- Gewers FL, Ferreira GR, Arruda HF, Silva FN, Comin CH, Amancio DR, et al. Principal component analysis: a natural approach to data exploration. *ACM Comput Surv*. 2021;54(4):1–34. <https://doi.org/10.48550/arXiv.1804.02502>
- Singh V, Zhou S, Ganie Z, Valverde B, Avila L, Marchesan E, et al. Rice production in the Americas. In: Chauhan BS, Jabran K, Mahajan G, editors. *Rice Production Worldwide*. Cham: Springer; 2017:137–68. https://doi.org/10.1007/978-3-319-47516-5_6
- Singh RK, Chaudhary BD. Biometrical methods in quantitative genetic analysis. New Delhi: Kalyani Publishers; 1981.
- Cheabu S, Pansrithong M, Malumpom C, Thongjoo C, Romkaew J. Variability and relationship between agronomic traits and grain yield in rice (*Oryza sativa* L.) under heat stress conditions. *Int J Agric Technol*. 2024;20:1823–42.
- Kumar N, Chhokar RS, Meena RP, Kharub AS, Gill SC, Tripathi SC, et al. Challenges and opportunities in productivity and sustainability of rice cultivation system: a critical review in Indian perspective. *Cereal Res Commun*. 2021;1:29. <https://doi.org/10.1007/s42976-021-00214-5>
- Wang P, Zhou G, Yu H, Yu S, Mahendran R, Veerabadhiran P, Robin

- S, Raveendran M. Principal component analysis of rice germplasm accessions under high temperature stress.
21. Fine mapping a major QTL for flag leaf size and yield-related traits in rice. *Theor Appl Genet.* 2011;123:1319–30. <https://doi.org/10.1007/s00122-011-1669-6>
 22. Liu C, Wu C, Yuan Z, Tian B, Yu P, Xu D, et al. Multi-trait genome-wide association studies reveal novel pleiotropic loci associated with yield and yield-related traits in rice. *J Integr Agric.* 2024. <https://doi.org/10.1016/j.jia.2024.07.026>
 23. Anilkumar C, Sah RP, Muhammed Azharudheen TP, Behera S, Mohanty SP, Anandan A, et al. Integrating multi-trait genomic selection with simulation strategies to improve grain yield and parental line selection in rice. *Ann Appl Biol.* 2025;186(2):216–27. <http://doi.org/10.1111/aab.12964>
 24. Habib MA, Azam MG, Haque MA, Hassan L, Khatun MS, Nayak S, et al. Climate-smart rice (*Oryza sativa* L.) genotypes identification using stability analysis, multi-trait selection index and genotype-environment interaction at different irrigation regimes with adaptation to universal warming. *Sci Rep.* 2024;14(1):13836. <https://doi.org/10.1038/s41598-024-64808-9>
 25. Panigrahi AK, Bharathi M, Kumaravadivel N. Morphological characterization of advanced backcross population for yield and yield attributing character in rice (*Oryza sativa* L.). *Int J Curr Microbiol Appl Sci.* 2018;7(2):2812–9. <https://doi.org/10.20546/ijcmas.2018.702.342>
 26. Sudharani M, Reddy PR, Reddy GH, Raju CS. Correlation and path coefficient analysis for yield and physiological attributes in rice (*Oryza sativa* L.) hybrids under saline soil conditions. 2013.
 27. Zeng L, Kwon TR, Liu X, Wilson C, Grieve CM, Gregorio GB. Genetic diversity analyzed by microsatellite markers among rice (*Oryza sativa* L.) genotypes with different adaptations to saline soils. *Plant Sci.* 2004;166(5):1275–85. <http://doi.org/10.1016/j.plantsci.2004.01.005>
 28. Immanuel SC, Pothiraj N, Thiyagarajan K, Bharathi M, Rabindran R. Genetic parameters of variability, correlation and path coefficient studies for grain yield and other yield attributes among rice blast disease resistant genotypes of rice (*Oryza sativa* L.). *Afr J Biotechnol.* 2011;10(17):3322–34. <http://doi.org/10.5897/AJB10.2575>
 29. Rajamadhan R, Murugan E. Genetic variability, correlation and path coefficient studies for grain yield and other yield attributing traits in rice (*Oryza sativa* L.). *Adv Life Sci.* 2016;5(6):2145–50.
 30. Ramya B, Saravanan K, Kumar PS. Genetic parameters of variability, correlation and path coefficient studies for grain yield and other yield attributes traits in rice (*Oryza sativa* L.). *Plant Arch.* 2012;12(2):711–5.
 31. Seyoum M, Alamerew S, Bantte K. Genetic variability, heritability, correlation coefficient and path analysis for yield and yield related traits in upland rice (*Oryza sativa* L.). *J Plant Sci.* 2012;7(1):13–22. <http://doi.org/10.3923/jps.2012.13.22>
 32. Jambhulkar NN, Bose LK. Genetic variability and association of yield attributing traits with grain yield in upland rice. *Genetika.* 2014;46(3):831–8. <http://doi.org/10.2298/GENSR1403831J>
 33. Sudeepthi K, Srinivas TV, Kumar BR, Jyothula DP, Umar SN. Assessment of genetic variability, character association and path analysis for yield and yield component traits in rice (*Oryza sativa* L.). *Electron J Plant Breed.* 2020;11(1):144–8. <https://doi.org/10.37992/2020.1101.026>
 34. Abhilash R, Thiruvengadam T, Dhatchinamoorthy S. Genetic studies in F2 for biometrical traits in rice (*Oryza sativa* L.). *Electron J Plant Breed.* 2018;9(3):1067–76. <https://doi.org/10.5958/0975-928X.2018.00133.3>
 35. Karim MJ, Jesmin F, Ali MH, Islam MN, Mahmud E. Variability, heritability and genetic advances for quantitative characters in T. Aman rice. *Int J Biol Res.* 2016;1(2):1–5.
 36. Mahalakshmi PJ, Satish Y, Ramana JV, Raju MR. Correlation and path coefficient analysis for yield and its component traits in advanced breeding lines of rice (*Oryza sativa* L.). *J Adv Biol Biotechnol.* 2024;27(9):769–77. <http://doi.org/10.9734/jabb/2024/v27i91350>
 37. Hossain S, Haque M, Rahman J. Genetic variability, correlation and path coefficient analysis of morphological traits in some extinct local Aman rice (*Oryza sativa* L.). *J Rice Res.* 2015;3(158):2. <https://doi.org/10.4172/2375-4338.1000158>
 38. Sharma N, Murty NS, Mall P, Bhardwaj SB. An analysis on the yield and yield contributing characters of rice in tarai region of Uttarakhand. *Int J Chem Stud.* 2018;6(3):42–7.
 39. Sharif S, Javaid RA, Majeed A, Ahmed MS, Sani QU, Siddique F, et al. Genetic diversity estimation of rice genotypes based on morphological and quality parameters through principal component analysis. *Pak J Agric Res.* 2023;36(3):207–16. <https://doi.org/10.17582/journal.pjar/2023/36.3.207.216>
 40. Khatun M, Islam MR, Khan MA, Hossain MK, Islam AK. Principal component analysis and estimated breeding values for selecting suitable parental genotypes in rice (*Oryza sativa* L.). *SABRAO J Breed Genet.* 2023;55(6):1886–96. <http://doi.org/10.54910/sabao2023.55.6.3>
 41. Reddy VR, Rani TS, Kumari A, Soundharya B. Correlation and path analysis for yield and yield attributes in maintainer lines of rice (*Oryza sativa* L.). *Int J Bioresour Stress Manag.* 2023;14(6):900–8. <https://doi.org/10.23910/1.2023.3417>
 42. Sanku G, Hepziba SJ, Sheeba A, Hemalatha G, Senthil K. Genetic variability and relatedness among yield characters in rice landraces and improved varieties. *Electron J Plant Breed.* 2022;13(3):932–9. <http://doi.org/10.37992/2022.1303.122>
 43. SaiVenkat KS, Sekhar MR, Vemireddy LR, Umar SN, Bommisetty R. Multivariate analyses of rice indigenous germplasm for identification of high yield potential genotypes based on yield and its attributes. *Genet Resour Crop Evol.* 2024;71(8):4807–23. <http://doi.org/10.1007/s10722-024-01932-5>
 44. Muthuramu S, Ragavan T. Association analysis for yield traits in rainfed rice (*Oryza sativa* L.). *Int J Curr Microbiol Appl Sci.* 2020;9(4):1512–7. <https://doi.org/10.20546/ijcmas.2020.904.178>
 45. Savitha P, Kumari RU. Genetic variability studies in segregating generation for grain and nutritional quality traits in rice (*Oryza sativa* L.). *J Appl Nat Sci.* 2016;8(1):63. <https://doi.org/10.31018/jans.v8i1.748>
 46. Sabar M, Mustafa SE, Ijaz M, Khan RA, Shahzadi F, Saher H, et al. Rice breeding for yield improvement through traditional and modern genetic tools. *Eur J Ecol Biol Agric.* 2024;1(1):14–19. [https://doi.org/10.59324/ejeba.2024.1\(1\).02](https://doi.org/10.59324/ejeba.2024.1(1).02)
 47. Gour L, Maurya SB, Koutu GK, Singh SK, Shukla SS, Mishra DK. Characterization of rice (*Oryza sativa* L.) genotypes using principal component analysis including scree plot & rotated component matrix. *Int J Chem Stud.* 2017;5(4):975–83.
 48. Abebe T, Alamerew S, Tulu L. Genetic variability, heritability and genetic advance for yield and its related traits in rainfed lowland rice (*Oryza sativa* L.) genotypes at Fogera and Pawe, Ethiopia. *Adv Crop Sci Technol.* 2017;5(2):272. <https://doi.org/10.4172/2329-8863.1000272>
 49. Nadim MK, Rahman M, Sikder MR, Adhikary S, Atiq NB, Alam MM, et al. Performance, adaptability and multi-trait stability of rice varieties (*Oryza sativa* L.) in rainfed conditions. *Agrosyst Geosci Environ.* 2025;8(1):e70038. <http://doi.org/10.1002/agg2.70038>
 50. Bhangar NA, Arain F, Pandrani MI, Khan N, Soomro G, Rao F, et al. Correlation and heritability estimation for agronomic traits in rice (*Oryza sativa* L.) genotypes. *Kashmir J Sci.* 2024;3(3). <https://doi.org/10.63147/kjrs.v3i3.79>
 51. Sandhu N, Yadav S, Kumar Singh V, Kumar A. Effective crop management and modern breeding strategies to ensure higher

- crop productivity under direct seeded rice cultivation system: a review. *Agronomy*. 2021;11(7):1264. <https://doi.org/10.3390/agronomy11071264>
52. Xu Y, Wang X, Ding X, Zheng X, Yang Z, Xu C, et al. Genomic selection of agronomic traits in hybrid rice using an NCII population. *Rice*. 2018;11:1–0. <https://doi.org/10.1186/s12284-018-0223-4>
 53. Christina GR, Thirumurugan T, Jeyaprakash P, Rajanbabu V. Principal component analysis of yield and yield related traits in rice (*Oryza sativa* L.) landraces. *Electron J Plant Breed*. 2021;12(3):907–11. <https://doi.org/10.37992/2021.1203.125>
 54. Bailey-Serres J, Parker JE, Ainsworth EA, Oldroyd GE, Schroeder JI. Genetic strategies for improving crop yields. *Nature*. 2019;575(7781):109–18. <https://doi.org/10.1038/s41586-019-1679-0>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc
See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.