



REVIEW ARTICLE

# Genetic approaches to impart brown leaf spot resistance in rice: A review

A S Mary Navya<sup>1</sup>, C Babu<sup>2\*</sup>, Ramalingam Suresh<sup>3</sup>, Swaminathan Manonmani<sup>3</sup>, Chellapan Gopalakrishnan<sup>4</sup> & G Senthilkumar<sup>5</sup>

<sup>1</sup>Department of Genetics & plant breeding, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

<sup>2</sup>Directorate of Research, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

<sup>3</sup>Department of Rice, Centre for Plant Breeding and Genetics, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

<sup>4</sup>Department of Plant Pathology, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

<sup>5</sup>Department of Agronomy, Tamil Nadu Agricultural University, Coimbatore 641 003, Tamil Nadu, India

\*Correspondence email - [babu.c@tnau.ac.in](mailto:babu.c@tnau.ac.in)

Received: 05 May 2025; Accepted: 19 July 2025; Available online: Version 1.0: 20 September 2025

**Cite this article:** Mary Navya AS, Babu C, Ramalingam S, Swaminathan M, Chellapan G, Senthilkumar G. Genetic approaches to impart Brown leaf spot resistance in rice: A review. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.9310>

## Abstract

Brown leaf spot (BLS) of rice, caused by the fungal pathogen *Bipolaris oryzae*, is a widespread and economically significant disease affecting rice-growing regions worldwide. Historically known for contributing to the Bengal famine during the 1940s, the disease continues to impact both yield and grain quality, especially in rice grown in rainfed and nutrient-deficient ecologies. Despite its long-standing presence, Brown leaf spot remains as an under-researched disease compared to other major rice pathogens, limiting progress in resistance breeding. Climate change and widespread drought events have witnessed an increased incidence of this disease in the recent past. Therefore, developing stable and broad-spectrum resistance to multiple strains of *Bipolaris oryzae* across diverse agro-ecological zones is essential. This review provides insight into recent advances that are transforming BLS resistance breeding from the discovery of novel resistance donors in diverse germplasm to the deployment of high-resolution Quantitative Trait Loci (QTL) mapping, genome-wide association studies (GWAS) and Clustered Regularly Interspaced Short Palindromic Repeats associated protein 9 (CRISPR-Cas9) based functional gene validation. It highlights how combining conventional breeding with molecular tools like marker-assisted selection and gene pyramiding accelerates the development of rice lines with broad and durable BLS resistance. The review emphasizes recent advances that support more efficient and climate-resilient breeding strategies for sustainable rice production.

**Keywords:** *bipolaris oryzae*; breeding approaches; brown leaf spot; host plant resistance; resistance genes

## Introduction

Rice (*Oryza sativa* L.) is a diploid species ( $2n = 24$  chromosomes) belonging to the Poaceae family. It has a panicle-type inflorescence and produces a dry, one-seeded fruit known as a caryopsis, typical of grasses (1). Rice serves as a vital food crop in the developing world and a staple for over half of the global population. More than 3.5 billion people rely on rice for over 20 % of their daily caloric intake (2). Rising population and economic development continue to increase pressure on rice production systems (3). According to FAO's most recent projection, global cereal commerce in 2022-23 is estimated at 468 million tonnes, reflecting the growing global demand for staple crops like rice (4). However, yield-reducing diseases such as brown leaf spot (BLS) remains as a significant threat, particularly in low-input, rain-fed systems. BLS is responsible for average global yield losses of 4-5 % and in severe outbreaks, losses can rise to 45-52 % (5, 6). To maintain steady rice production, it is important to control these disease outbreaks and reduce losses from year to year. This requires knowing extent of damage that diseases cause and keeping track of new

or changing disease problems, especially those linked to changes in farming practices and technology (7). Sustainable agricultural methods and broad-spectrum genetic resistance are complimentary approaches to disease management. Despite significant advancements in recent decades, further integration of these approaches is needed for tangible results in farmer's fields. Preventing losses from biotic and abiotic stresses provides a resource-neutral strategy to increase production in light of growing need for food. The ability to maintain or increase rice yields cost-effectively will depend on developing varieties that remain productive despite these challenges (8).

Although chemical, biological and cultural practices are widely used to manage BLS in rice, they are often costly, environmentally unsafe and provide only limited long-term effectiveness (9). In contrast, host plant resistance provides a sustainable and economically viable solution, especially when employing strategies such as gene pyramiding, multilines and partial resistance accumulation (10, 11). Advances in QTL mapping and rice genomics have equipped breeders with powerful tools to develop brown spot-resistant cultivars suited

to diverse environment. Marker-assisted selection (MAS) now offers a faster, more efficient and reliable alternative to phenotypic selection, enabling early and accurate identification of resistance traits, even for complex characteristics that are challenging to breed through conventional methods. Considerable advances in QTL mapping, cloning of resistance genes and understanding of defence signalling pathways have significantly enhanced rice breeding, making MAS a powerful tool for developing brown spot-resistant cultivars with greater precision and in a shorter timeframe (12, 13). This review provides a comprehensive understanding of molecular defence mechanisms in rice, that are essential for accelerating breeding efforts towards long-lasting resistance against brown leaf spot. It also highlights recent advances and outlines future directions to manage disease outbreaks and enhance rice productivity.

### Brown leaf spot

Rice is susceptible to 55 fungal diseases, of which 43 are seed-borne or seed-transmissible. Among these, three pathogens are particularly significant: *Pyricularia oryzae* (blast), *Bipolaris oryzae* (brown spot) and *Xanthomonas oryzae* (bacterial leaf blight) (14). Together, blast and brown spot diseases contribute to approximately 30 % of yield losses in rice (15). Brown spot, caused by the fungal pathogen *Bipolaris oryzae* Subr. and Jain, has been reported in Japan as early as 1900 (16). In India, brown leaf spot has been reported across all rice-growing states (17), since its first report by Sundararaman in 1919 from Madras. Historically famous for triggering the 1942-43 Bengal famine, which damaged 50-90 % of rice crop in Eastern India and caused two million deaths (18), BLS remains a global threat across Asia, Africa, the Americas and parts of Europe (16). The causal organism of brown leaf spot in rice has undergone multiple taxonomic reclassifications. Initially named *Helminthosporium oryzae* by Breda de Haan, it was later identified as *Ophiobolus miyabeanus* (perfect stage) and reclassified as *Cochliobolus miyabeanus* (19). The anamorph was renamed *Bipolaris oryzae* in 1955 due to its bipolar germination but was also referred to as *Drechslera oryzae* based on conidial features. *Bipolaris oryzae* produces well-branched mycelium and septate conidia of variable size (20). Optimal growth occurs at 27-30 °C, with conidial germination favoured at 25-30 °C and pH 6.8-7 (17). Genetic analysis of 288 isolates from Iran, Japan and the Philippines using 12 SSR

markers revealed 278 haplotypes, indicating high genetic diversity and gene flow likely facilitated by seed exchange (21). This highlights the need for quantitative resistance breeding and diverse pathogen isolates in screening programs to ensure durable disease control.

### Symptoms and economic impact

*Bipolaris oryzae* affects rice at both seedling and mature stages, damaging various plant parts including leaf blades, coleoptiles, sheaths, panicle branches, glumes and spikelets (22-24). Seedlings in nurseries often suffer severe damage (25). The disease is characterized by oval or circular leaf lesions with light grey centers, reddish-brown margins and yellow halos (17). It can also cause seedling blight, necrotic lesions, grain discoloration and dark spots on glumes, which may lead to shrivelled grains (26). Under favorable conditions, conidiophores and conidia form on infected tissues, giving a velvety appearance. In severe cases, the fungus penetrates glumes and endosperm, causing seed rotting, pre-emergence damping-off and reduced germination. Rarely, black lesions may also appear on young roots, though node and internode infections are uncommon (27-29). Fig. 1 shows the characteristic symptoms of brown leaf spot (BLS) on rice leaves.

BLS poses a widespread threat to rice production, resulting in considerable yield losses and economic damage (30-32). The disease causes both quantitative and qualitative losses in grain yield, as highlighted by (33). Currently, brown leaf spot is widespread across rice-growing regions, leading to significant yield losses, second only to blast disease (34). Yield loss averages 10 % across lowland rice systems in South and Southeast Asia, with severely infected fields experiencing reductions of up to 45 % (33, 35). Table 1 summarizes the yield loss caused by brown leaf spot disease in rice in different regions. Besides quantitative yield losses, BLS reduces seed quality and germinability, further compounding its economic impact. The disease's severity is exacerbated by global warming, as the pathogen thrives in high-temperature conditions (36). Climate change and nutritional imbalances, especially deficiencies in nitrogen and potassium, are key factors driving its increased occurrence (37). Considering the substantial yield losses and reductions in grain quality, brown leaf spot remains an economic threat to rice production worldwide, necessitating urgent attention to mitigate its



**Fig. 1.** Shows the characteristic symptoms of brown leaf spot (BLS) on rice leaves.

**Table 1.** Yield loss due to brown leaf spot disease in rice

| Region                 | Yield Loss (%)   | Reference |
|------------------------|--|-----------|
| South & Southeast Asia | 10 % average, up to 45 % in severely infected fields                                     | (33, 40)  |
| Nigeria                | 12 %   | (94)      |
| North Sierra Leone     | 8-23 %   | (95)      |
| Florida, USA           | 16-40 %, with reduced milling quality  | (96)      |
| India (General)        | 19.2 % to 58.8 %   | (97, 98)  |
| India (West Bengal)    | 5.6-11.7 % in susceptible variety Tilakkachery, 2.2 % in moderately resistant Bhasamanik | (98)      |
| Jammu Region, India    | 32.65 % to 74.51 % in 2018, 31.64 % to 75.83 % in 2019                                   | (37)      |
| Punjab, India          | 4.45 % to 34.38 %  | (75)      |

financial burden on farmers and economies.

### Environmental conditions

The development and spread of brown leaf spot (BLS) depend on three key factors: favorable climatic conditions, sufficient pathogen inoculum (from air, water, soil, or seeds) and a susceptible host plant. If any of these is absent, disease development will not occur (38). Optimal environmental conditions for BLS include temperatures between 23.6 °C and 34.2 °C, relative humidity of 80.7-85.8 %, 4.6-6.2 hrs of sunlight and wind speeds of 1.0-1.6 km/h (39). These factors promote pathogen growth, spore production and germination, with peak disease severity typically observed in the third week of September. The disease can affect rice from seedling to milk stage, with yield loss varying based on cultivar and time of infection (28). Deficiencies in potassium, manganese, magnesium, silicon, iron, or calcium intensify disease severity, which is why BLS is often referred to as the “poor farmer’s disease”. Its prevalence in low-input, rain-fed and upland systems underscores the need for better soil nutrient management to reduce its impact on smallholder farmers (40).

Agronomic practices and crop microclimate significantly influence BLS progression. Conventional planting methods result in higher disease incidence, while furrow planting and the System of Rice Intensification (SRI) help reduce severity by lowering humidity and increasing canopy temperature. High plant density and delayed transplanting worsen disease by creating a cooler, more humid microclimate. Meteorology-based regression models have shown high accuracy in predicting outbreaks, enabling timely interventions. Integrated approaches such as optimized planting geometry, reduced plant density and timely transplanting can effectively alter the microclimate, offering an eco-friendly strategy to manage BLS (41). Understanding these environmental and agronomic interactions is essential for developing predictive tools and location-specific management practices.

### Pathogenesis

The pathogenesis of brown leaf spot (BLS) in rice begins with primary infections, which is mainly seed-borne (42, 43). *Bipolaris oryzae* is a necrotrophic pathogen that invades by killing host cells and feeding on the dead tissue. The disease cycle initiates when conidia (asexual spores) land on rice leaves and germinate, with germ tubes penetrating the cuticle directly or through stomata (17). The pathogen releases phytotoxins, such as ophiobolins, which disrupt cell membranes, leading to characteristic brown lesions with grey centres and reddish-brown margins (44). A key gene, BMK1, encoding a MAP kinase, is essential for fungal growth, conidiation and virulence. Its


disruption reduces hyphal growth, impairs aerial hyphae, abolishes conidia production and renders the fungus non-pathogenic-paralleling MAPK roles in other fungal pathogens like *Cochliobolus heterostrophus* and *Magnaporthe grisea* (45). Targeting MAPK signalling may therefore offer novel disease control strategies (46).

Secondary infections arise from wind-dispersed conidia originating from infected residues like straw and stubble (17, 47). Soil and certain weed hosts also act as inoculum sources (48). Conidial production is highly responsive to environmental conditions, thriving at 21-26 °C and relative humidity above 92 %, peaking at 100 % RH (49). High humidity and intermittent rains during cooler months (October-December) favour spore dispersal and infection (50). Disease progression is most severe between flowering and dough stages, with flag leaf infection strongly correlating with kernel contamination. The pathogen can invade all parts of the kernel, especially the rachilla and sterile lemmas. As *B. oryzae* is seed-transmissible, it can cause coleoptile, root and first leaf infections in seedlings. Thus, flag leaf severity can serve as a predictor of seed infection, underscoring the importance of seed health testing and treatment in disease management. Fig. 2 shows the infection cycle of *Bipolaris oryzae*, the causal agent of brown leaf spot in rice.

### Molecular mechanism and host-pathogen interaction

The infection process of *Bipolaris oryzae* in rice involves a series of intricate interactions between the pathogen and host, which are mediated by genetic and biochemical mechanisms. The pathogen employs various molecular strategies to invade host tissues, suppress plant defences and establish infection, ultimately determining the susceptibility or resistance of the host plant. The interaction between *Bipolaris oryzae* and its host is influenced by both the pathogen's virulence strategies and the host's defense mechanisms. In addition to rice, alternative hosts such as *Leersia hexandra*, a common weed in Indian rice fields and wild rice species have been reported to harbour the pathogen in nature (51, 52). Severe infections on *Zizania aquatica*, cultivated wild rice in the USA, demonstrate the broad host range of the pathogen (53). Collateral hosts could play a critical role in the pathogen's epidemiology, serving as reservoirs for inoculum and contributing to disease persistence in rice ecosystems.

The physiological and biochemical effects of *Bipolaris oryzae* infection are severe, especially in susceptible rice cultivars like 'Oochikara.' Infection triggers oxidative stress, increasing malondialdehyde (MDA) levels and electrolyte leakage (EL), which signal cell membrane damage. As the disease progresses, chlorophyll and other photosynthetic



**Fig. 2.** Infection cycle of *Bipolaris oryzae*, the causal agent of brown leaf spot in rice.

pigments decline due to ophiobolin toxins that damage chloroplasts and induce chlorosis. This leads to a 65 % reduction in net carbon assimilation, stomatal conductance and transpiration, resulting in significant loss of healthy leaf area and reduced plant vigour (54). Studies using rice cultivar MTL189 inoculated with two isolates of the pathogen, B5 (highly compatible) and K2 (less compatible), have shown that the severity of infection varies based on the host's defence mechanisms. The B5 isolate caused extensive tissue degradation and sporulation, whereas the K2 isolate led to limited hyphal growth and minimal damage due to the accumulation of hydrogen peroxide ( $H_2O_2$ ) and polyphenolic compounds in the resistant plants. Structural defences such as callose deposition and fluorescent papillae further restricted pathogen development in resistant interactions, showing the role of early defence responses in limiting disease spread (55). The pathogen deploys toxins like ophiobolins that suppress rice's phenolic-based defences by significantly reducing phenolic content, peroxidase and phenylalanine-ammonia lyase activity that are key components of resistance. Although treatments that enhance phenolic metabolism (e.g., catechol, quinic acid, phenylalanine and Ethrel) delayed symptom development, the pathogen's toxins counteracted these responses, facilitating host colonization and disease progression (56).

Rice defence against *B. oryzae* involves physiological and biochemical changes. Silicon (Si) induces resistance by disrupting the pathogen's manipulation of ethylene (ET) signalling, reducing ethylene production, phenolic compound accumulation and fungal spread, independent of salicylic acid (SA) or jasmonic acid (JA) pathways. Instead, it directly targets pathogen virulence factors (57). Abscisic acid (ABA) also enhances resistance by suppressing ET signalling via the OsMPK5 gene, involving hydrogen peroxide ( $H_2O_2$ ) production at infection sites without relying on SA, JA, or callose responses (58). The interplay between ABA and ET signalling highlights the complexity of hormonal crosstalk in regulating host defense. The tryptophan (Trp) pathway is pivotal in rice defense against *Bipolaris oryzae*, with infection triggering increased anthranilate synthase (AS) activity, upregulated Trp biosynthetic genes (OASA2, OASB1, OASB2) and higher levels of

Trp and intermediates like anthranilate and indole. Serotonin, produced via tryptophan decarboxylase (TDC), accumulates in infected tissues and integrates into cell walls through oxidative polymerization, forming a barrier that suppresses fungal growth. In serotonin-deficient mutant rice (sl), higher susceptibility occurs due to tryptamine accumulation, leading to lesion spread (59). Exogenous serotonin application restores cell wall defence and resistance, underscoring its role as a crucial secondary metabolite in rice defence.

Rice defence mechanisms differ against *Magnaporthe grisea* (rice blast) and *Cochliobolus miyabeanus* (brown leaf spot). Resistant varieties like Tetep show hypersensitive responses (HR) and localized cell death against both pathogens, while susceptible varieties (e.g., Nakdong) experience rapid cell death and uncontrolled fungal spread with *B. oryzae*. *B. oryzae* conidial germination fluid (CGF) triggers rapid cell death in both resistant and susceptible cultivars, indicating phytotoxins. Defence inducers like benzo (1,2,3)thiadiazole-7-carbothioic acid S-methyl ester (BTH) enhance resistance to *M. grisea* but not *B. oryzae*, suggesting pathogen-specific responses (60). Understanding *B. oryzae* infection strategies and rice defences highlights targets for breeding varieties with early, robust defence activation.

### Approaches to improve BLS resistance

#### From chemical to biocontrol methods

Management of brown spot disease in rice has traditionally relied on chemical fungicides such as carbendazim and combinations like azoxystrobin + difenoconazole, which significantly reduce disease severity and improve yields. For instance, carbendazim, applied at 0.1 % concentration, significantly decreases disease intensity while improving plant health parameters and yield. Studies on the IRGA 424 rice cultivar in Paraguay revealed that a combination of azoxystrobin (20 %) and difenoconazole (12.5 %) applied at specific growth stages (R2, R3 and R4) reduced disease incidence from 90 % to 15.7 % and increased yields by 23 %. However, concerns over environmental impact, pathogen resistance and high costs limit their long-term use, especially for resource-poor farmers (61).

Botanical extracts have gained attention as sustainable alternatives that combine moderate disease control efficacy with environmental safety. Neem (*Azadirachta indica*) extracts, for example, have demonstrated significant reductions in disease intensity at a 10 % concentration. Besides disease suppression, neem also enhances plant health, contributing to increased root biomass and shoot vigour (62). Biocontrol agents have shown promise in managing brown spot disease. Under laboratory conditions, *Trichoderma harzianum* exhibited the highest inhibition of *B. oryzae* mycelial growth (65.33 %), followed by *Trichoderma viride* (57.76 %) and *Pseudomonas fluorescens* (32.68 %). Field trials with rice varieties Basmati-370, Jaya and PC-19 demonstrated that foliar applications of these biocontrol agents significantly reduced disease severity. *T. harzianum* provided the highest disease reduction (up to 64 %) and led to yield increases of 8-17 % depending on the variety. The efficacy of these biocontrol agents is attributed to the production of antifungal compounds and the induction of systemic resistance in plants (63). Similarly, the integration of *Trichoderma harzianum* with botanical extracts has shown promising results. Combined application of *Trichoderma harzianum* (0.1 %) with garlic bulb extract (10 %) reported a disease intensity of 33.06 %, markedly lower than the untreated control (45.20 %) (64). This treatment not only suppressed the disease and promoted plant growth but also proved economically feasible for farmers, as the *Trichoderma* and neem combination yielded the best cost-benefit ratio of 1:2.67. Utilizing biocontrol agents like *Trichoderma* spp. offers an eco-friendly alternative to chemical fungicides, mitigating issues related to pesticide resistance, environmental pollution and health hazards.


Phenolic antioxidants such as salicylic and benzoic acids have emerged as eco-friendly alternatives for managing brown spot disease. These compounds exhibit strong antifungal activity, with benzoic acid (20 mM) significantly reducing disease severity and improving yield components like grain filling and weight. Treated plants also show enhanced chlorophyll, carbohydrate and protein levels, offering both disease suppression and improved crop performance (65). While chemical fungicides remain effective for rapid control, their environmental and economic drawbacks highlight the

need for integrated, sustainable solutions. Combining botanical extracts, *Trichoderma* formulations and phenolic compounds presents a cost-effective approach. Recent findings show that red light (550-650 nm) induces resistance in rice by activating the tryptophan and phenylpropanoid pathways, reducing lesion formation without directly affecting the pathogen. This light-induced defence, which strengthens with longer exposure, offers a novel, non-chemical strategy for brown spot management (66). Fig. 3 shows the integrated management strategies for brown leaf spot disease in rice.

#### Sources of resistance in breeding lines

Host resistance has been widely recognized as a sustainable method for managing brown leaf spot disease. Several studies have focused on identifying resistant genotypes across different regions. Asia and Africa have been notable sources of resistance, providing valuable genetic material for breeding programs aimed at developing resistant cultivars for farmers. The genetic variability of *Bipolaris oryzae* and the differential susceptibility of rice cultivars advocate the need to use host resistance to combat the disease effectively (67). The earliest recorded investigation of varietal resistance dates back to 1930, when the Korean cultivar 'Kutto-urupe' was reported as highly susceptible based on field observations without artificial inoculation (68). Subsequent studies in the United States identified the cultivar 'Mubo-Aikoku' as resistant through seedling inoculation tests (69). In Japan, Yoshii and Matsumoto employed spray inoculation across different growth stages, demonstrating that cultivars like 'Tetep' and 'Ginnen' were resistant, while others, including 'Tadukan,' showed moderate resistance (70). In Egypt, studies categorized 'Pi1' and 'YNA282' as resistant, contributing further to the pool of resistant germplasm (71).

Recent efforts have screened advanced breeding lines and commercial cultivars for BLS resistance. Table 2 shows the disease rating scale for screening rice varieties against Brown leaf spot. Among 29 advanced breeding lines and four check varieties (BR 11, BR 22, BR 25 and BRRI dhan 38), one accession showed resistance during the *T. aman* season, while the boro season revealed 29 moderately resistant cultivars (72). A field and laboratory assessment of 60 rice genotypes in Nepal identified significant variability in resistance levels, with only



**Fig. 3.** Disease management approaches for brown leaf spot in rice.

**Table 2.** Disease rating scale for screening rice varieties against brown leaf spot

| Rating | Affected leaf area | Host reaction        |
|--------|--------------------|----------------------|
| 1      | No incidence       | Immune               |
| 2      | Less than 1 %      | Highly Resistant     |
| 3      | 1 - 3 %            | Resistant            |
| 4      | 4 - 5 %            | Resistant            |
| 5      | 11 - 15 %          | Moderately Resistant |
| 6      | 16 - 25 %          | Moderately Resistant |
| 7      | 26 - 50 %          | Susceptible          |
| 8      | 51 - 75 %          | Susceptible          |
| 9      | 76 - 100 %         | Highly susceptible   |

one genotype showing moderate resistance. However, most genotypes, including landraces, exhibited high susceptibility, emphasizing the need for broader germplasm exploration (73). In Cameroon, field evaluations of four rice varieties (Nerica 3, Nerica 8, Kamkou and White Rice) during 2015-2016 found Nerica 8 and Nerica 3 to be highly tolerant, with disease severity at 110 DAS recorded at 2.05 % and 2.72 %, respectively. These varieties demonstrated superior resilience compared to Kamkou and White Rice, making them promising candidates for resistance breeding programs (74). Regional surveys in Punjab, India, highlighted its impact on local rice varieties. Commercial cultivars such as PR 116, PR 114 and PR 106 were highly susceptible, with PR 116 reaching up to 28.41 % severity. In contrast, the variety 'Jaya' consistently showed lower susceptibility, with a severity of only 3.23 % (75). These results underscore the importance of identifying and utilizing resistant genotypes, both traditional and modern, for managing brown spot disease. Enhanced screening efforts and the inclusion of diverse land races and wild relatives in breeding programs remain crucial for developing sustainable solutions to this persistent challenge.

Advancements in hybrid rice breeding have identified promising parental lines and hybrids with significant resistance to brown spot (*Bipolaris oryzae*). A combined genetic analysis highlighted *Giza 178*, *Giza 179*, *HR195*, *Giza 181* and *Giza 182* as resistant parental lines suitable for developing hybrids with robust resistance to brown spot disease. Among cytoplasmic male sterile (CMS) lines, *K17A* exhibited strong general combining ability (GCA) for resistance-related traits, making it an effective contributor in hybrid breeding. The hybrid *K17A* × *Giza 182* demonstrated significantly enhanced resistance, with desirable reductions in disease severity and infected leaf percentages. These traits, alongside beneficial heterosis, highlight its potential for commercial deployment (76). The study emphasized the predominant role of additive gene action in conferring resistance, with *K17A*, *HR195* and *Giza 182* emerging as superior general combiners.

#### QTL mapping

Several cultivars that have been categorized as resistant did not show complete resistance (immunity) to Brown spot. Some varieties such as 'Tadukan' and 'Tetep' showed quantitative resistance to BLS (70-78). The first QTL analysis for resistance to BLS was conducted in 2008 using a set of recombinant inbred lines (RILs) derived from a cross between 'Tadukan' (resistant)

and 'Hinohikari' (susceptible). Three QTLs for brown spot resistance were detected on chromosomes 2, 9 and 11, named qBS2, qBS9 and qBS11. The most significant QTL was qBS11, with the highest LOD score of 5.11 in composite interval mapping (CIM) and 2.82 in interval mapping (IM). This QTL accounted for 11.9 % to 15.3 % of the total phenotypic variation and reduced the disease score by 0.34 to 0.58. qBS9 was detected with a LOD score of 3.45 in CIM and 2.72 in IM, explaining 9.7 % to 12.9 % of the phenotypic variation. qBS2, identified only in CIM analysis, explained 11.1 % of the variation. The resistant alleles from the *Tadukan* variety reduced disease scores, confirming its effectiveness against brown spot and setting the stage for future map-based cloning to understand resistance mechanisms further (47). The previously found QTLs (qBS2, qBS9 and qBS11) were associated with brown spot resistance at the seedling stage in controlled conditions. Another research aimed to validate these QTLs under field conditions and introduce them into a regional rice variety, 'Koshihikari'. This study identified three field resistance QTLs; qBSfR1 (chromosome 1), qBSfR4 (chromosome 4) and qBSfR11 (chromosome 11). The major QTL qBSfR11 aligns with the previously identified qBS11, demonstrating consistent effectiveness across conditions. Near-isogenic lines (NILs) containing qBSfR11 displayed significant field resistance, highlighting its potential utility for breeding programs. qBSfR1 showed no significant impact on resistance, suggesting limited practical value. The findings underscore qBSfR11 as a major QTL for developing brown spot-resistant rice varieties through marker-assisted selection (79).

The first practical brown spot (BS)-resistant rice variety named *Mienoyume BSL* was developed by incorporating the QTL qBSfR11 from the local resistant line 'Tadukan' into the background of the high-yielding yet susceptible cultivar 'Mienoyume'. The resistance to BS is controlled by a single recessive gene, designated as *bsr1*, located within a 1.3-Mbp region on chromosome 11. This gene was introduced into *Mienoyume* through successive backcrossing. The resulting near-isogenic line (*bsr1*-NIL) showed significantly improved resistance to multiple strains of *Bipolaris oryzae*, yielding a BS disease score significantly lower than *Mienoyume* and achieving a 28.8 % higher yield in fields with severe BS incidence. The *bsr1*-NIL demonstrated stable agricultural performance, including larger grain width, comparable days to heading and improved grain filling, even under disease stress. The protein content was slightly lower in *bsr1*-NIL compared to *Mienoyume* (80). To verify the resistance conferred by the 'Tadukan' alleles at qBSfR1 and qBSfR11, near-isogenic lines (NILs) containing each of these QTL alleles from 'Tadukan' in the background of susceptible variety 'Koshihikari' were bred and evaluated under field conditions. A NIL containing the 'Tadukan' allele at qBSfR11 acquired significant BLS field resistance, but that with the 'Tadukan' allele at qBSfR1 did not. Besides 'Tadukan', 'Kasalath' was also identified as a resistant cultivar (36).

Quantitative trait loci (QTLs) for brown spot (BS) resistance in rice were discovered on chromosomes 2 (qBSR2-kc), 7(qBSR7-kc), 9(qBSR9-kc) and 11(qBSR11-kc) using backcrossed inbred lines (BILs) from a cross between the resistant indica variety 'CH45' and the susceptible japonica cultivar 'Koshihikari'. qBSR11-kc was the major contributor to

BLS resistance which explained 23.0 %-25.9 % of the phenotypic variation in disease resistance. qBSR9-kc and qBSR11-kc were consistently detected in both 2015 and 2016, while qBSR2-kc and qBSR7-kc were only identified in 2016, potentially due to variations in climate, such as higher temperatures and increased rainfall. The resistance alleles at qBSR2-kc, qBSR7-kc and qBSR11-kc originated from CH45, while the allele at qBSR9-kc came from *Koshihikari* (81). A recombinant inbred line (RIL) population from a cross between *Danteshwari* (moderately tolerant) and *Dagad deshi* (susceptible) was used to identify quantitative trait loci (QTLs) associated with brown spot tolerance. qBS1.1, qBS5.1 and qBS5.2 are the QTLs that were located on chromosomes 1 and 5. qBS1.1 (chromosome 1) exhibited a lower effect, explaining 9.15 % of phenotypic variance. qBS5.1 and qBS5.2 (chromosome 5) had high phenotypic variances of 55.35 % and 55.5 %, respectively, indicating major contributions to brown spot resistance. The negative additive effects suggested that the resistance alleles were inherited from the susceptible parent, *Dagad deshi* (82). Table 3 highlights Quantitative trait loci (QTLs) associated with resistance to Brown Leaf Spot (BLS) in rice.

#### Genome-Wide Association Studies (GWAS)

GWAS has become a powerful approach to identify genomic regions linked to complex traits such as disease resistance. While limited studies have focused specifically on BLS in rice, GWAS applied to related traits and species offers valuable insights. In switchgrass, a relative of rice, GWAS identified 27 SNPs associated with BLS resistance, many of which were linked to defense-related genes. This includes genes encoding nucleotide-binding site leucine-rich repeat (NBS-LRR) proteins, pentatricopeptide repeat (PPR) proteins and ABC transporters, all of which are involved in plant defense mechanisms (83). Although direct GWAS studies on *Bipolaris oryzae* resistance in rice are limited, significant progress has been made for analogous leaf spot diseases. For example, narrow brown leaf spot (caused by *Cercospora janseana*) was associated with a key locus, CRSP2.1, on chromosome 2. Fine mapping and functional characterization of these loci are being carried out to develop resistant rice cultivars. Moreover, GWAS on temperate-adapted japonica and indica rice identified 14 marker-trait associations (MTAs) for blast resistance. Several MTAs co-located with known blast resistance genes (Pi genes), while

three (BRF10, BRF11-2, BRGC11-3) may represent novel resistance loci (84). Given that GWAS was successful in identifying the resistance loci for blast disease, a similar approach for brown leaf spot disease, could identify new resistance genes and quantitative trait loci QTLs that may provide a sound basis for breeding BLS-resistant varieties of rice. GWAS on other rice diseases, such as bacterial leaf blight, identified resistance loci and candidate genes, which increased our understanding of defense mechanisms in rice. Such studies reveal NBS-LRR proteins, crucial for the recognition of pathogens and activation of defense responses (85). In addition to the resistance genes, biocontrol agents like *Trichoderma harzianum* offer a complementary strategy. *Trichoderma harzianum* has demonstrated strong antifungal activity against *Bipolaris oryzae*, both *in vitro* and field studies. *In vitro* assays showed that *T. harzianum* overgrows *B. oryzae* and produces inhibitory metabolites, while foliar sprays (10<sup>8</sup> spores/mL) under field conditions significantly reduced disease severity and incidence and boosted grain yield, carbohydrate and protein content in rice leaves (86). Compelling research shows that genes from *Trichoderma* can be transferred into rice to enhance resistance. For example, the *ech42* gene (encoding an endochitinase) from *T. atroviride* was introduced into rice via transformation, leading to enhanced resistance not only against sheath blight but also against rice blast (87). This strategy could similarly enhance resistance to brown leaf spot by enabling rice to express antifungal genes effective against *Bipolaris oryzae*.

#### Omics approaches

Omics technologies have improved our understanding of complex traits such as disease resistance in rice. These approaches-genomics, transcriptomics, proteomics and metabolomics-offer insights into the host-pathogen interactions underlying Brown Leaf Spot (BLS). Transcriptomics is the most widely used omics tool for BLS studies. Techniques like RNA-Seq and qRT-PCR helps to unravel gene expression changes in rice upon pathogen infection. For instance, resistant cultivars upregulate defense genes such as *OsFBN6*, while susceptible ones show disrupted expression, revealing key candidates for breeding (88). Defense-related pathways, including the tryptophan pathway, phenylpropanoid metabolism and serotonin biosynthesis, are often differentially expressed. These

**Table 3.** QTLs identified for brown leaf spot resistance in rice

| QTL       | Chromosome | Flanking markers                | Mapping population                                    | Donor       | % Phenotypic variance | Additive effect | Reference |
|-----------|------------|---------------------------------|---|-------------|-----------------------|-----------------|-----------|
| qBS2      | 2          | RM3515*-RM263                   | RILs (Tadukan × Hinohikari)                           | Tadukan     | 11.1                  | 0.29            | (47)      |
| qBS9      | 9          | RM3907-RM566*<br>RM566*-RM257   | RILs (Tadukan × Hinohikari)                           | Tadukan     | 9.7<br>12.9           | -0.28<br>-0.62  | (47)      |
| qBS11     | 11         | RM206-RM6449*<br>RM6499*-RM6623 | RILs (Tadukan × Hinohikari)                           | Tadukan     | 15.3<br>11.9          | -0.34<br>-0.58  | (47)      |
| qBSfR 1   | 1          | RM10604                         | RILs (Tadukan × Hinohikari) in Koshihikari background | Tadukan     | 12.3                  | -0.4            | (79)      |
| qBSfR 4   | 4          | RM273                           | RILs (Tadukan × Hinohikari) in Koshihikari background | Tadukan     | 10.8                  | 0.3             | (79)      |
| qBSfR 11  | 11         | RM27096                         | RILs (Tadukan × Hinohikari) in Koshihikari background | Tadukan     | 19.2                  | -0.5            | (79)      |
| qBSR9-kc  | 9          | RM3919*-RM6797                  | BILs (CH45 × Koshihikari)                             | Koshihikari | 6.5                   | 0.2             | (81)      |
| qBSR11-kc | 11         | RM6534*-RM4112                  | BILs (CH45 × Koshihikari)                             | CH 45       | 25.9                  | -0.5            | (81)      |
| qBSR2-kc  | 2          | RM5578-RM5614                   | BILs (CH45 × Koshihikari)                             | CH 45       | 17.1                  | -0.6            | (81)      |
| qBSR7-kc  | 7          | RM1353-1*-RM5672                | BILs (CH45 × Koshihikari)                             | CH 45       | 7.8                   | -0.4            | (81)      |

contribute to early pathogen recognition and physical barrier formation. Notably, serotonin-deficient mutants exhibit higher susceptibility, underscoring its role as a secury metabolite in rice defense (59).

Integration with metabolomics helps to link gene expression to functional outputs such as higher phenolic compound accumulation in resistant lines. Combined omics approaches can detect novel resistance genes and metabolic signatures for use in marker-assisted selection (89). Recent transcriptomic studies in North-East Indian rice cultivars revealed a biphasic fungal infection strategy-early metabolic activation in the fungus followed by late effector expression in susceptible hosts. This suggests the need for early defense activation in breeding programs (90). Going forward, the fusion of omics datasets with machine learning could accelerate the discovery of candidate genes and biomarkers for BLS resistance. Integrative approaches will be essential for designing rice cultivars with durable, multi-layered defense.

#### Genome editing/CRISPR Cas9

Genome editing, particularly CRISPR/Cas9, has emerged as a revolutionary method for precise and efficient crop improvement. In rice, this technology has already been employed to develop resistance against major diseases such as blast and bacterial blight. Its application in managing Brown Leaf Spot (BLS) holds immense promise. CRISPR/Cas9 has demonstrated its potential by knocking out susceptibility genes, thereby enhancing disease resistance. For instance, editing the *OsERF922* gene significantly increased resistance to blast (91), while modifications in *SWEET* gene promoters disrupted effector-triggered susceptibility to bacterial blight (92). This precise technology shows great potential for targeting brown leaf spot (BLS). Possible uses include the knockout of susceptibility genes that allow for pathogen infection or the alteration of resistance-related genes to boost their effectiveness against BLS. Multiplex genome editing can enable the concurrent targeting of multiple genes, allowing for the combination of resistance traits and offering broad-spectrum and long-lasting resistance. CRISPR/Cas9-mediated editing of susceptibility genes (*Bsr-d1*, *Pi21* and *ERF922*) in rice revealed that *ERF922* mutants conferred strong blast resistance, while *Pi21* and *ERF922* knockouts also enhanced bacterial blight resistance via activation of salicylic acid (SA) and jasmonic acid (JA) pathways (93). This approach, which preserved key agronomic traits, suggests that targeting analogous genes in rice could similarly enhance BLS resistance via multiplex genome editing. By merging genome editing with molecular markers and genomic selection, this strategy could hasten the creation of BLS-resistant rice varieties, ensuring sustainable production in the face of evolving pathogen threats.

#### Challenges and Limitations

Breeding rice varieties resistant to brown leaf spot (BLS) presents several challenges due to the disease's complex epidemiology and the multifaceted interactions among the host plant, pathogen and environmental conditions.

- Slow disease initiation and progression, influenced by specific environmental conditions (e.g., optimal temperature and humidity), delay symptom expression and hinder early detection.

- Persistence of the pathogen is aided by continuous rice cultivation, favourable climate and survival on infected plant residues and seeds.
- High initial inoculum levels from contaminated seeds and crop debris lead to more severe disease outbreaks.
- Host susceptibility is affected by physical traits like leaf surface morphology and physiological stresses such as nutrient imbalance and water stress.
- Resistance evaluation is challenging due to environmental variability, difficulty in achieving reliable phenotyping and complex inheritance involving multiple QTLs.
- Pathogen variability and strong genotype × environment interactions complicate breeding for stable resistance.
- Marker-assisted selection (MAS) requires precise identification and mapping of QTLs, which is constrained by the polygenic nature of BLS resistance.
- Pyramiding multiple resistance QTLs is essential for durable resistance but may face compatibility and pleiotropy challenges.

#### Conclusion and Future Perspectives

Brown leaf spot (BLS) continues to pose a significant threat to global rice production, particularly in low-input, rain-fed systems. Its persistence is driven by complex interactions among the host, pathogen and environment, compounded by nutrient deficiencies and the pathogen's broad host range. While chemical and cultural controls offer temporary relief, host plant resistance remains the most sustainable and eco-friendly long-term solution. Considerable progress has been made in identifying QTLs, especially from landraces and wild *Oryza* species and incorporating them into elite lines using molecular tools such as marker-assisted selection (MAS). Advances in genome editing technologies like CRISPR/Cas9 further enable precise and efficient resistance breeding. To reduce the spread of BLS in the field, practical strategies such as the use of disease-free seed, crop rotation, residue management and balanced nutrition should be prioritized alongside genetic solutions.

To overcome the challenges posed by the polygenic nature of resistance and high pathogen variability, future breeding must focus on gene pyramiding, genomic selection and the development of multi-resistant cultivars. Additionally, continuous pathogen surveillance and the strategic deployment of resistance genes can help minimize selection pressure and prolong resistance durability. Future efforts must adopt a multidisciplinary approach, integrating host-pathogen interaction studies, pathogen genomics and systems biology to uncover novel resistance mechanisms. High-throughput phenotyping, environmental modelling and next-generation sequencing, supported by robust bioinformatics, will be crucial for breeding climate-resilient, broad-spectrum BLS-resistant cultivars. Collaboration across disciplines will accelerate the development of durable solutions, safeguarding rice yields in the face of evolving disease pressure and climate change.

## Authors' contributions

ASMN wrote the main manuscript. RS, SM, CG and GS have reviewed the manuscript. BC did the final revision of manuscript. All authors read and approved the manuscript.

## Compliance with ethical standards

**Conflict of interest:** Authors do not have any conflict of interests to declare.

**Ethical issues:** None

## References

1. Hay JO, Spanswick RM. Computational analysis of rice (*Oryza sativa* L.) panicle topology and ripening. *Seed Science Research*. 2006;16(4):243-50. <https://doi.org/10.1017/SSR2006257>
2. Khush GS. IR varieties and their impact. *Int Rice Res Inst*; 2005.
3. Zhang Q. Strategies for developing green super rice. *Proc Natl Acad Sci U S A*. 2007;104(42):16402-9. <https://doi.org/10.1073/pnas.0708013104>
4. FAOStat F. Food and agriculture organization. Rome: FAO; 2005.
5. Savary S, Willocquet L, Pethybridge SJ, Esker P, McRoberts N, Nelson A. The global burden of pathogens and pests on major food crops. *Nat Ecol Evol*. 2019;3(3):430-9. <https://doi.org/10.1038/s41559-018-0793-y>
6. Kaboré KH, Kassankogno AI, Tharreau D. Brown spot of rice: Worldwide disease impact, phenotypic and genetic diversity of the causal pathogen *Bipolaris oryzae* and management of the disease. *Plant Pathol*. 2025;74(4):908-22. <https://doi.org/10.1111/ppa.14075>
7. Mew TW, Leung H, Savary S, Vera Cruz CM, Leach JE. Looking ahead in rice disease research and management. *Crit Rev Plant Sci*. 2004;23(2):103-27. <https://doi.org/10.1146/annurev-arplant-042809-112116>
8. Leung H, Zhu Y, Revilla-Molina I, Fan JX, Chen H, Pangga I, et al. Using genetic diversity to achieve sustainable rice disease management. *Plant Dis*. 2003;87(10):1156-69. <https://doi.org/10.1094/PDIS.2003.87.10.1156>
9. Jeger MJ, Pautasso M, Holdenrieder O, Shaw MW. Modelling disease spread and control in networks: implications for plant sciences. *New Phytol*. 2007;174(2):279-97. <https://doi.org/10.1111/j.1469-8137.2007.02028.x>
10. Manandhar H, Lyngs Jørgensen H, Mathur S, Smedegaard-Petersen V. Suppression of rice blast by preinoculation with avirulent *Pyricularia oryzae* and the nonrice pathogen *Bipolaris sorokiniana*. *Phytopathology*. 1998;88(7):735-9. <https://doi.org/10.1094/PHYTO.1998.88.7.735>
11. Bonman J, Khush G, Nelson R. Breeding rice for resistance to pests. *Annu Rev Phytopathol*. 1992;30(1):507-28. <https://doi.org/10.1146/annurev.py.30.090192.002451>
12. Lopez-Gerena J. Mapping QTL controlling durable resistance to rice blast in the cultivar *Oryzica Llanos 5*. Kansas State University; 2006.
13. Talukder ZI, Tharreau D, Price AH. Quantitative trait loci analysis suggests that partial resistance to rice blast is mostly determined by race-specific interactions. *New Phytol*. 2004;162(1):197-209. <https://doi.org/10.1111/j.1469-8137.2004.01010.x>
14. Richardson M. An annotated list of seed-borne diseases. *Int Seed Test Assoc*; 1979.
15. Bedi K, Gill H. Losses caused by the brown leaf spot disease of rice in the Punjab. 1960.
16. Khalili E, Sadravi M, Naeimi S, Khosravi V. Biological control of rice brown spot with native isolates of three *Trichoderma* species. *Braz J Microbiol*. 2012;43:297-305. <https://doi.org/10.1590/S1517-838220120001000035>
17. Ou S. Rice diseases. Kew: Commonwealth Mycological Institute; 1985.
18. Alam S, Seth R, Singh H, Srivastava J, Shukla D. Screening of disease resistant varieties against brown leaf spot of *Oryza sativa* in Allahabad, India. *Am J Exp Agric*. 2016;14(1):1-11. <https://doi.org/10.9734/AJEA/2016/29059>
19. Sreenivasaprasad S, Johnson R, Chakrabarti N. Epidemiology and disease management of brown spot of rice in India. In: Major fungal diseases of rice: recent advances. 2001:293-306. [https://doi.org/10.1007/978-94-017-2157-8\\_21](https://doi.org/10.1007/978-94-017-2157-8_21)
20. Subramanian C, Jain B. A revision of some graminicolous *Helminthosporia*. *Curr Sci*. 1966;35(14):352-5.
21. Hayward A. Studies on the brown leaf spot disease of rice (*Oryza sativa* L.) and the causal organism *Helminthosporium oryzae* Breda de Haan. *Trans Br Mycol Soc*. 1933;18(4):230-48. [https://doi.org/10.1016/S0007-1536\(33\)80041-9](https://doi.org/10.1016/S0007-1536(33)80041-9)
22. Singh U. Studies on the brown leaf spot disease of rice (*Helminthosporium oryzae*) and its control. *Indian J Agric Sci*. 1947;17:275-82.
23. Misra J. Studies on the genus *Helminthosporium* in India. II. Subgenus *Cochliobolus*. *Trans Br Mycol Soc*. 1971;56(3):395-401. [https://doi.org/10.1016/S0007-1536\(71\)80059-0](https://doi.org/10.1016/S0007-1536(71)80059-0)
24. Rai J, Joshi L. Cultural variability in *Helminthosporium oryzae* Breda de Haan. *Plant Dis Rep*. 1974;58(9):802-4.
25. Sunder S, Satyavathi C, Alice D. Morphological and pathogenic variability of *Bipolaris oryzae* in rice. *Indian Phytopathol*. 2003;56(4):453-6.
26. Rosyara UR, Subedi KD, Baniya BK, Kharel R, Joshi S, Duveiller E. The effect of spot blotch disease on performance of wheat genotypes in warm and humid conditions of Nepal. *Euphytica*. 2007;153(1-2):307-21. <https://doi.org/10.1007/s10681-006-9247-y>
27. Rizvi SMH, Misra AK. Cultural and morphological variability in isolates of *Bipolaris sorokiniana* of wheat. *Ann Pl Protec Sci*. 2005;13(1):119-21.
28. Singh R, Kumar D, Kumar M. Morphological and pathogenic variability among isolates of *Bipolaris sorokiniana* causing leaf blight in wheat. *Indian Phytopathol*. 2008;61(1):55-9.
29. Siddaramaiah A, Somashekar R, Shetty H. Variability among isolates of *Bipolaris sorokiniana* infecting wheat in Karnataka. *Plant Dis Res*. 2000;15(2):168-71.
30. Grover R, Pannu PS, Sharma P. Morphological and pathogenic variability among Indian isolates of *Bipolaris sorokiniana*, the spot blotch pathogen of wheat. *Indian Phytopathol*. 2003;56(4):389-93.
31. Singh A, Maurya S, Kumar S, Kumar S, Srivastava AK. Characterization of *Bipolaris sorokiniana* isolates causing spot blotch in wheat. *Indian Phytopathol*. 2016;69(2):112-6.
32. Ray A, Chowdhury S, Saha S, Khatua DC. Pathogenic and genetic variability among *Bipolaris sorokiniana* isolates infecting wheat in eastern India. *Indian Phytopathol*. 2021;74(1):73-9. <https://doi.org/10.1007/s42360-020-00246-3>
33. Ahiwar RK, Gange D, Patel A. Morphological and cultural variability among isolates of *Bipolaris sorokiniana* (Sacc.) Shoem. incitant of spot blotch of wheat. *Int J Curr Microbiol App Sci*. 2017;6(4):344-52. <https://doi.org/10.20546/ijcmas.2017.604.040>
34. Adhikary SK, Dutta S, Hossain A, Hossain MK, Roy B, Chandra S, et al. Assessment of genetic variability and molecular diversity of *Bipolaris sorokiniana* associated with spot blotch disease of wheat. *Cereal Res Commun*. 2021;49(4):563-71. <https://doi.org/10.1007/s42976-021-00133-z>
35. Smiley RW, Gourlie JA, Easley SA, Patterson LM, Whittaker RG.

Crop damage estimates for crown rot of wheat and barley in the Pacific Northwest. *Plant Dis.* 2005;89(6):595-604. <https://doi.org/10.1094/PD-89-0595>

36. Gupta PK, Chand R, Vasistha NK, Pandey SP, Kumar U, Mishra VK, et al. Spot blotch disease of wheat: the current status of research on genetics and breeding. *Plant Pathol.* 2018;67(3):508-31. <https://doi.org/10.1111/ppa.12773>

37. Kumar U, Mishra VK, Joshi AK, Mishra CN, Bhardwaj SC, Sharma JB, et al. Resistance in wheat to spot blotch caused by *Bipolaris sorokiniana*. *Euphytica.* 2015;207(1):119-29. <https://doi.org/10.1007/s10681-015-1554-z>

38. He X, Singh PK, Dreisigacker S, Singh RP, Duveiller E, Santra D, et al. Identification of resistance to spot blotch in wheat germplasm using genome-wide association mapping. *Plant Genome.* 2020;13(1):e20016. <https://doi.org/10.1002/tpg2.20016>

39. McDonald BA, Linde C. Pathogen population genetics, evolutionary potential and durable resistance. *Annu Rev Phytopathol.* 2002;40:349-79. <https://doi.org/10.1146/annurev.phyto.40.120501.101443>

40. Khaledi N, Taheri P, Falahati-Rastegar M, Tarighi S. Host-pathogen interaction between rice and *Bipolaris oryzae*: a compatible interaction. *Physiol Mol Plant Pathol.* 2017;99:122-32. <https://doi.org/10.1016/j.pmp.2017.05.004>

41. Garg M, Kalra A. Fungal endophytes and their role in biological control and plant health. In: Arora NK, editor. *Plant microbe symbiosis: fundamentals and advances.* Springer; 2013:109-36. [https://doi.org/10.1007/978-81-322-1287-4\\_6](https://doi.org/10.1007/978-81-322-1287-4_6)

42. Rudd JJ, Keon JP, Hammond-Kosack KE. The wheat *Septoria* interaction: a paradigm for the advancement of understanding of plant-pathogen interactions. In: Deising HB, editor. *The Mycota V: plant relationships.* Springer; 2009:271-90. [https://doi.org/10.1007/978-3-540-87407-2\\_16](https://doi.org/10.1007/978-3-540-87407-2_16)

43. Anwar A, Riaz S, Naseem S. Role of plant growth promoting rhizobacteria in improving abiotic stress tolerance in crop plants. In: Egamberdieva D, Ahmad P, editors. *Plant microbiome: stress response.* Springer; 2018:127-52. [https://doi.org/10.1007/978-981-10-5514-0\\_7](https://doi.org/10.1007/978-981-10-5514-0_7)

44. Backer R, Rokem JS, Ilangumaran G, Lamont J, Praslickova D, Ricci E, et al. Plant growth-promoting rhizobacteria: context, mechanisms of action and roadmap to commercialization of biostimulants for sustainable agriculture. *Front Plant Sci.* 2018;871:1473. <https://doi.org/10.3389/fpls.2018.01473>

45. Manzar N, Irfan M, Alam M, Meena RM, Khan SA. Biocontrol potential of *Bacillus* spp. against fungal phytopathogens. In: Meena MK, Upadhyay RS, editors. *Biocontrol mechanisms of endophytic microorganisms.* Springer; 2021:251-73. [https://doi.org/10.1007/978-981-33-6661-9\\_13](https://doi.org/10.1007/978-981-33-6661-9_13)

46. Roy AK, Hossain M, Hossain M, Alam MS, Biswas P, Islam M, et al. Antagonistic potential of rhizospheric bacteria against brown spot pathogen (*Bipolaris oryzae*) and growth promotion in rice. *Arch Phytopathol Plant Prot.* 2022;55(6):746-63. <https://doi.org/10.1080/03235408.2022.2075474>

47. Sato H, Ando I, Hirabayashi H, Takeuchi Y, Arase S, Kihara J, et al. QTL analysis of brown spot resistance in rice (*Oryza sativa* L.). *Breeding Science.* 2008;58(1):93-6. <https://doi.org/10.1270/jsbbs.58.93>

48. Biswas S, Das RH, Sharma GL, Das HR. Isolation and characterization of a novel cross-infective rhizobia from *Sesbania aculeata* (Dhaincha). *Current Microbiology.* 2008;56:48-54. <https://doi.org/10.1007/s00284-007-9037-0>

49. Chattopadhyay S, Chakraborti N. Effect of agronomic practices on susceptibility of rice plants to infection of *Helminthosporium oryzae*. *Rice Newsletter.* 1965;13:39-44.

50. Chakraborti N. Studies on certain aspects of *Helminthosporium* disease of paddy with particular reference to conditions in West Bengal: Doctoral thesis, University of Calcutta, India; 1955.

51. Chattopadhyay S, Chakraborti N. Occurrence in nature of an alternative host (*Leersia hexandra* Sw.) of *Helminthosporium oryzae* Breda de Haan. *Nature.* 1953;172(4377):550.

52. Subramanian C, Ramakrishnan G. The occurrence of brown spot *Helminthosporium oryzae* Breda de Haan on some wild species of *Oryza*. 1973.

53. Kernkamp M, Kroll R, Woodruff W. Diseases of cultivated wild rice in Minnesota. 1976.

54. Dallagnol LJ, Rodrigues FA, Martins SC, Cavatte PC, DaMatta FM. Alterations on rice leaf physiology during infection by *Bipolaris oryzae*. *Australasian Plant Pathology.* 2011;40(4):360-5. <https://doi.org/10.1007/S13313-011-0048-8>

55. Thuy TTT, Lübeck M, Smedegaard-Petersen V, de Neergaard E, Jørgensen HJ. Infection biology of *Bipolaris oryzae* in rice and defence responses in compatible and less compatible interactions. *Agronomy.* 2023;13(1):231. <https://doi.org/10.3390/agronomy13010231>

56. Vidhyasakaran P, Borromeo E, Mew T. *Helminthosporium oryzae* toxin suppresses phenol metabolism in rice plants and aids pathogen colonization. *Physiological and Molecular Plant Pathology.* 1992;41(5):307-15. [https://doi.org/10.1016/0885-5765\(92\)90018-q](https://doi.org/10.1016/0885-5765(92)90018-q)

57. Van Bockhaven J, Spíchal L, Novák O, Strnad M, Asano T, Kikuchi S, et al. Silicon induces resistance to the brown spot fungus *Cochliobolus miyabeanus* by preventing the pathogen from hijacking the rice ethylene pathway. *New Phytologist.* 2015;206(2):761-73. <https://doi.org/10.1111/nph.13270>

58. De Vleesschauwer D, Yang Y, Vera Cruz C, Höfte M. Abscisic acid-induced resistance against the brown spot pathogen *Cochliobolus miyabeanus* in rice involves MAP kinase-mediated repression of ethylene signaling. *Plant Physiology.* 2010;152(4):2036-52. <https://doi.org/10.1104/pp.109.152702>

59. Ishihara A, Hashimoto Y, Tanaka C, Dubouzet JG, Nakao T, Matsuda F, et al. The tryptophan pathway is involved in the defense responses of rice against pathogenic infection via serotonin production. *The Plant Journal.* 2008;54(3):481-95. <https://doi.org/10.1111/j.1365-313X.2008.03441.x>

60. Ahn I-P, Kim S, Kang S, Suh S-C, Lee Y-H. Rice defense mechanisms against *Cochliobolus miyabeanus* and *Magnaporthe grisea* are distinct. *Phytopathology.* 2005;95(11):1248-55. <https://doi.org/10.1094/PHYTO-95-1248>

61. Barúa M, Quintana LA, Ortíz AA. Chemical control of rice brown spot (*Bipolaris oryzae*) in Paraguay. *The Journal of the Society for Tropical Plant Research.* 2019;6(1):148-51. <https://doi.org/10.22271/tpr.2019.v6.i1.021>

62. Kumar M, Simon S. Efficacy of certain botanical extracts in the management of brown leaf spot of rice caused by *Helminthosporium oryzae*. *Biosciences Biotechnology Research Asia.* 2016;13(4):2015. <http://doi.org/10.13005/bbra/2358>

63. Gupta V, Shamas N, Razdan V, Mahajan S, Fatima K, Sharm S, et al. Management of brown spot of rice (*Oryza sativa* L.) caused by *Bipolaris oryzae* by bio-control agents. *Int J Curr Microbiol Appl Sci.* 2018;7(04):3472-7. <https://doi.org/10.20546/ijcmas.2018.704.393>

64. Lal T. Effect of *Trichoderma harzianum* and selected botanicals on brown leaf spot of paddy (*Oryza sativa* L.) caused by *Helminthosporium oryzae* (Breda de Haan). 2024. <https://doi.org/10.9734/arrb/2024/v39i92131>

65. Shabana Y, Abdel-Fattah G, Ismail A, Rashad Y. Control of brown spot pathogen of rice (*Bipolaris oryzae*) using some phenolic antioxidants. *Brazilian Journal of Microbiology.* 2008;39:438-44. <https://doi.org/10.1590/S1517-83822008000300010.9734/arrb/2024/v39i9213106>

66. Parada RY, Mon-nai W, Ueno M, Kihara J, Arase S. Red-light-induced resistance to brown spot disease caused by *Bipolaris oryzae* in rice. *Journal of Phytopathology.* 2015;163(2):116-23. <https://doi.org/10.1111/jph.12288>

67. Nneke N. Screening lowland rice varieties for resistance to brown

spot disease in Enyong creek rice field in Akwa Ibom state of Nigeria. Global Journal of Pure and Applied Sciences. 2012;18(1):5-10.

68. Nagai L, Hara S. On the inheritance; of variegation disease in a strain of rice plant. 1930. <https://doi.org/10.1266/JJG.5.151>
69. Adair CR. Inheritance in rice of reaction to *Helminthosporium oryzae* and *Cercospora oryzae*. 1941.
70. Yoshii H. Studies on the resistance to Helminthosporiose of the rice varieties introduced to Japan (1). Bull of Matsuyama Agricultural College. 1951;6:25-60.
71. Balal M, Omar R, El-Khadem M, Aidy I. Inheritance of resistance to the brown spot disease of rice, *Cochliobolus miyabeanus*. 1979. <https://doi.org/10.5897/JPBCS2016.0613>
72. Hossain M. Reaction of breeding lines/cultivars of rice against brown spot and blast under field condition. Asian Journal of Plant Sciences (Pakistan). 2004;3(5). <https://doi.org/10.3923/ajps.2004.614.617>
73. Bhandari R, Khatri S, Acharya B, Gauchan U, Adhikari G, Thapa S. Screening of rice genotypes against brown leaf spot (*Bipolaris oryzae*) (Breda de Haan) Shoemaker, under field and laboratory conditions at Banke, Nepal. Discover Plants. 2024;1(1):10. <https://doi.org/10.1007/s44372-024-00008-z>
74. Tueguem WNK, Dooh JPN, Songwe TA, Heu A, Sale CE, Nanfack AD, et al. Reaction of some rice (*Oryza sativa* L.) varieties to brown spot disease caused by *Bipolaris oryzae* (Breda de Haan) Shoemaker. 2021. <https://doi.org/10.30574/msarr.2021.2.2.0008>
75. Pannu P, Chahal S, Sharma VK, Kaur M, Bagga P. Occurrence of brown leaf spot of rice in Punjab, its effect on grain yield and its control. Indian Phytopathology. 2006;59(2):190-3.
76. Elshenawy MM, Elgamal WH, Anis GB, Awad F. Combined genetic analysis of brown spot (*Helminthosporium oryzae*) disease for developed hybrid combinations and their parental lines in hybrid rice. Sustainable Food Production. 2018;1:37-48. <https://doi.org/10.18052/www.scipress.com/SFP.1.37>
77. Ohata K, Kubo C. Studies on the mechanism of disease resistance of rice varieties to *Cochliobolus miyabeanus*. 1974.
78. Eruotor P. Varietal reaction of rice to isolates of *Cochliobolus miyabeanus*. 1986.
79. Sato H, Matsumoto K, Ota C, Yamakawa T, Kihara J, Mizobuchi R. Confirming a major QTL and finding additional loci responsible for field resistance to brown spot (*Bipolaris oryzae*) in rice. Breeding Science. 2015;65(2):170-5. <https://doi.org/10.1270/jsbbs.65.170>
80. Matsumoto K, Ota Y, Yamakawa T, Ono T, Seta S, Honda Y, et al. Breeding and characterization of the world's first practical rice variety with resistance to brown spot (*Bipolaris oryzae*). 2021. <https://doi.org/10.1270/jsbbs.21023>
81. Matsumoto K, Ota Y, Seta S, Nakayama Y, Ohno T, Mizobuchi R, et al. Identification of QTLs for rice brown spot resistance in backcross inbred lines derived from a cross between Koshihikari and CH45. Breeding Science. 2017;67(5):540-3. <https://doi.org/10.1270/jsbbs.17057>
82. Mandal L, Verma S, Kotasthane A, Verulkar S. QTL mapping for brown spot tolerance using RIL population of rice (*Oryza sativa* L.). European Journal of Biotechnology and Bioscience. 2017;5(3):17-20.
83. Songsomboon K, Crawford R, Crawford J, Hansen J, Cummings J, Mattson N, et al. Genome-wide associations with resistance to bipolaris leaf spot (*Bipolaris oryzae* (Breda de Haan) Shoemaker) in a northern switchgrass population (*Panicum virgatum* L.). Plants. 2022;11(10):1362. <https://doi.org/10.3390/plants11101362>
84. Volante A, Tondelli A, Desiderio F, Abbruscato P, Menin B, Biselli C, et al. Genome wide association studies for japonica rice resistance to blast in field and controlled conditions. Rice. 2020;13:1-17. <https://doi.org/10.1186/s12284-020-00431-2>
85. Shu X, Wang A, Jiang B, Jiang Y, Xiang X, Yi X, et al. Genome-wide association study and transcriptome analysis discover new genes for bacterial leaf blight resistance in rice (*Oryza sativa* L.). BMC Plant Biology. 2021;21(1):255. <https://doi.org/10.1186/s12870-021-03041-2>
86. Abdel-Fattah GM, Shabana YM, Ismail AE, Rashad YM. *Trichoderma harzianum*: a biocontrol agent against *Bipolaris oryzae*. Mycopathologia. 2007;164:81-9. <https://doi.org/10.1007/s11046-007-9032-9>
87. Liu M, Sun Z-x, Zhu J, Xu T, Harman GE, Lorito M. Enhancing rice resistance to fungal pathogens by transformation with cell wall degrading enzyme genes from *Trichoderma atroviride*. Journal-Zhejiang University Science. 2004;5(2):133-6. <https://doi.org/10.1631/BF02840913>
88. Cao F-Y, Zeng Y, Lee A-R, Kim B, Lee D, Kim S-T, et al. OsFBN6 enhances brown spot disease resistance in rice. Plants. 2024;13(23):3302. <https://doi.org/10.3390/plants13233302>
89. Ontoy JC, Ham JH. Mapping and omics integration: towards precise rice disease resistance breeding. Plants. 2024;13(9):1205. <https://doi.org/10.3390/plants13091205>
90. Singh S, Sarki YN, Marwein R, Singha DL, Velmurugan N, Chikkaputtaiah C. Unraveling the role of effector proteins in *Bipolaris oryzae* infecting North East Indian rice cultivars through time-course transcriptomics analysis. Fungal Biology. 2023;127(7-8):1098-110. <https://doi.org/10.1016/j.funbio.2023.05.003>
91. Wang F, Wang C, Liu P, Lei C, Hao W, Gao Y, et al. Enhanced rice blast resistance by CRISPR/Cas9-targeted mutagenesis of the ERF transcription factor gene OsERF922. PLoS One. 2016;11(4):e0154027. <https://doi.org/10.1371/journal.pone.0154027>
92. Oliva R, Ji C, Atienza-Grande G, Huguet-Tapia JC, Perez-Quintero A, Li T, et al. Broad-spectrum resistance to bacterial blight in rice using genome editing. Nature Biotechnology. 2019;37(11):1344-50. <https://doi.org/10.1038/s41587-019-0267-z>
93. Zhou Y, Xu S, Jiang N, Zhao X, Bai Z, Liu J, et al. Engineering of rice varieties with enhanced resistances to both blast and bacterial blight diseases via CRISPR/Cas9. Plant Biotechnology Journal. 2022;20(5):876-85. <https://doi.org/10.1111/pbi.13766>
94. Aluko M. Crop losses caused by the brown leaf spot disease of rice in Nigeria. Plant Disease Reporter. 1975;59(7):609-13.
95. Fomba S, Singh N. Crop losses caused by rice brown spot disease in mangrove swamps of northwestern Sierra Leone. International Journal of Pest Management. 1990;36(4):387-93.
96. Datnoff LE, Lentini RS. Brown spot in Florida rice. 1994.
97. Vidhyasekaran P, Ranganathan K, Palaniappan S, Ramasamy S. Effect of slow-release nitrogen fertilizers on rice brown spot disease. 1983.
98. Chattopadhyay S, Chakrabarti N, Ghosh A. Estimation of loss in yield of rice due to infection of brown spot incited by *Helminthosporium oryzae*. 1975.

#### Additional information

**Peer review:** Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

**Reprints & permissions information** is available at [https://horizonpublishing.com/journals/index.php/PST/open\\_access\\_policy](https://horizonpublishing.com/journals/index.php/PST/open_access_policy)

**Publisher's Note:** Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Indexing:** Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc  
See [https://horizonpublishing.com/journals/index.php/PST/indexing\\_abstracting](https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting)

**Copyright:** © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

**Publisher information:** Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.