



RESEARCH COMMUNICATION

Three species of lichenized fungi, new to Assam from Chirang district

Madhusmita Deka¹, Pungbili Islary² & Rebecca Daimari^{1*}

¹Department of Botany, Bodoland University, Kokrajhar 783 370, Assam, India

²Department of Botany, Kokrajhar University, Kokrajhar 783 370, Assam, India

*Correspondence email - rebeccadmr@gmail.com

Received: 28 May 2025; Accepted: 14 August 2025; Available online: Version 1.0: 06 March 2026

Cite this article: Madhusmita D, Pungbili I, Rebecca D. Three species of lichenized fungi, new to Assam from Chirang district. Plant Science Today (Early Access). <https://doi.org/10.14719/pst.9465>

Abstract

This paper presents three species of lichen viz. *Herpothallon minutum*, *Lecanora andina* and *Phaeophyscia kairamoi*, from the Chirang District, Assam. The specimens were identified using morphological, anatomical and chemical characters. The three species are reported for the first time from Assam. Detailed descriptions of the species with photographs are provided.

Keywords: Assam; lichenized fungi; new records; taxonomy

Introduction

Assam is a region primarily composed of fertile river valleys, hills and forests and is characterised by a highly humid tropical climate. It falls within the Eastern Himalayas, a biodiversity hotspot known for its rich and diverse flora and fauna, which encompasses over half of India's documented phanerogams. The botanical significance of this area is highlighted by its remarkable diversity and abundance of plant species, such as *Alstonia scholaris* (L.) R. Br., *Bombax ceiba* L., *Cassia fistula* L and *Lannea coromandelica* (Houtt.) Merr. (1). Owing to its favourable geographical conditions, the region harbours a rich diversity of lichens.

In India, a total of 3236 lichen taxa have been reported, comprising 3200 species, 29 varieties, 6 subspecies and 1 forma, distributed across 487 genera and 88 families. Of these, North-East India harbours the highest diversity, comprising 1842 taxa and the state Assam exhibits a remarkable increase in recorded lichen diversity, rising from 300 species in 2018 to 657 species in 2021 and currently recording 749 species of lichens (2). However, studies on lichens in Assam are still insufficient. The pioneering research on lichens in this state was conducted in 1881 by Stirton, a Scottish lichenologist (3). Subsequent contributions to the floristic study of lichen diversity in Assam have been made by various researchers (4–18). However, lichens have become an important cryptogamic group to study and document for their ecological significance. Thus, the present work aims to explore lichens growing in various localities within the Chirang District of Assam.

Materials and Methods

The collection of corticolous lichen specimens were conducted

during 2022–2023 in the Chirang district of Assam, specifically in areas near Bongaigaon Refinery and Petrochemicals Limited (BRPL) (Fig. 1). The collected specimens were air dried and identified under the microscope.

Morphological characterisation of the lichen thallus was performed using a stereo zoom microscope (Leica EZ4). Anatomical observation was conducted under a compound microscope (Leica DM750). Chemical spot tests on the thallus, medulla and apothecium were carried out using standard laboratory reagents—aqueous potassium hydroxide solution (K), aqueous calcium hypochlorite solution (C), 0.5 g of para-phenylenediamine dissolved in 5 mL of ethanol (P) and iodine solution (I).

Thin-layer chromatography (TLC) was performed using solvent system A (toluene-dioxane-acetic acid, 180:45:5) to identify lichen compounds (19). The taxa were identified in the Department of Botany, Bodoland University, Kokrajhar, Assam using morphological, anatomical and chemical information, supported by relevant published literature and were deposited in the herbarium of the Department of Botany, Bodoland University, Kokrajhar, Assam (abbreviated here as BUBH) (20–22). The nomenclature and families of the identified lichen species were updated according to IndexFungorum.org and outline of fungi and fungus-like taxa (23).

Results

From the ongoing study, only the species representing new records for the state are considered here. Detailed descriptions of the three newly recorded species are provided below:




Fig. 1. Map showing the study site of the Chirang district (Image created using QGIS software (version 3.38)).

Taxonomic description

1. *Herpothallon minutum* Jagad. Ram, Lichenologist 46(1): 45 (2014)(Fig. 2A-C)

Family: Arthoniaceae

Thallus crustose, corticolous, white to grey, round patches, thallus up to 3–11 cm in diameter, hyphae in the upper part, hypothallus whitish grey, loosely to firmly attached. The prothallus is white up to 4.5 mm wide, whitish and interwoven, with radiating hyphae. Pseudoisidia minute, numerous, granular. Chemistry: thallus K –, C + red, KC + red, KI + blue, P –, UV –; TLC: gyrophoric acid.

Distribution: India (Andaman & Nicobar Islands, Arunachal Pradesh, Uttarakhand and West Bengal).

Species examined: India– Assam, Chirang district, 26.33.04" N, 90.33.09" E, elevation 84 m, on the bark of *Citrus maxima* (Burm.) Merr., 18/10/2022, 2022-1655 (BUBH); collector: Madhusmita Deka.

2. *Lecanora andina* Räsänen, Anal. Soc. cient. argent. 128(3): 139 (1939) (Fig. 2D-F)

Family: Lecanoraceae

Thallus crustose, rimose to areolate, brownish grey, epruinose, esorediate, shiny. Ascomata apothecia, many, immersed when young, sessile in mature, 0.5–0.7 mm in diameter, with well-developed thalline margin, lecanorine, disc pale brown, margin entire and thin. Hymenium hyaline, 60–73 µm high, hypothecium light yellow, amphithecium with large crystals (pulcaris type), epihymenial pigment not dissolving in K.

Paraphyses simple, apical cell swollen, brown pigmented. Asci 8-spored, 51–57 µm, ascospore colourless, simple, ellipsoid, 10–11 × 5–6 µm. Chemistry: thallus K+ yellow, C –, P –, UV –; TLC: no substances detected.

Distribution: India (Goa, Himachal Pradesh, Maharashtra and Punjab), Argentina and Australia.

Species examined: India– Assam, Chirang district, 26.32.49" N, 90.29.12" E, elevation 85 m, on the bark of *Litchi chinensis* Sonn., 06/08/2022, 2022-1652 (BUBH); collector: Madhusmita Deka.

3. *Phaeophyscia kairamoi* (Vain.) Moberg, Symb. bot. upsala. 22(1): 40 (1977); *Physcia kairamoi* Vain., in Räsänen, Meddn Soc. Fauna Flora fenn. 46: 3 (1921) (Fig. 2G, H)

Family: Physciaceae

Thallus foliose, corticolous, irregularly branched, loosely adnate, corticated on both sides, lobes narrow, 1–1.5 mm wide, upper side grey, marginally isidiate, soredia absent, lacking both cyphellae and pseudocyphellae, lower side brown, rhizines black and simple, lower cortex paraplectenchymatous. Medulla is white and solid. Apothecia not seen. Chemistry: thallus K –, KC –, C –, P –, UV –; TLC: no substances detected.

Distribution: India (Jammu & Kashmir, Maharashtra and Uttarakhand), Canada, Finland, Italy, Kazakhstan, Kyrgyzstan, Mexico, Mongolia, Norway, Poland, Russian Federation, Sweden, Switzerland and USA.

Species examined: INDIA– Assam, Chirang district, 26.29.18.57" N, 90.36.46.41" E, elevation 74 m, on the bark of *Bombax ceiba* L.,




Fig. 2. A–C - *Herpothallon minutum*: A - thallus, B and C - thallus with chemical reagents K, KC and KI; D–F - *Lecanora andina*: D - thallus, E - hymenium with crystals, F - ascospores; G–H - *Phaeophyscia kairamoii*: G - thallus with isidia, H - medulla white and solid. 17/06/2023, 2023–1689 (BUBH); collector: Madhusmita Deka.

Discussion

The present work continues the ongoing research on lichens of Assam, which resulted in the discovery of three lichen species, *H. minutum*, *L. andina* and *P. kairamoii*, in the Chirang District, Assam, which represents a vital addition to the lichenological catalogue of Assam, as well as to Northeast India, a globally recognised biodiversity hotspot. It also emphasises the need for ongoing

taxonomic exploration of lichen diversity in less-documented areas like Chirang. Employing morphological, anatomical and chemical approaches for species identification adheres to stringent taxonomic protocols, ensuring precise classification. Such methods are essential in lichenology, where phenotypic variability often complicates accurate identification. Comprehensive descriptions and photographic documentation of the recorded species serve as key resources for reliable identification and comparative studies by future researchers working in similar ecological contexts.

Conclusion

The identified lichen species are reported for the first time from Assam, extending their known distribution and adding to the lichen biota of the state. The state stands out for its remarkable year-to-year increase in the number of recorded lichen species. The finding highlights the richness and yet-to-be-explored lichen biota of Assam, underlining the need for further taxonomic and ecological research. These additions enhance our understanding of lichen diversity in India. Future research should focus on investigating their ecological roles, biogeographical significance and potential applications.

Acknowledgements

Authors are thankful to the Department of Botany, Bodoland University, for providing laboratory facilities. Additionally, the authors express gratitude to DST-SERB, Government of India, for the instrumental facilities under the project, EMEQ (EEQ/2019/000547).

Authors' contributions

The specimens were collected and identified by MD and PI. MD wrote the initial draft of the manuscript, while RD reviewed and edited it, supervised the entire project.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interest to declare.

Ethical issues: None

References

- Assam State Biodiversity Board. Biodiversity of Assam [Internet]. Assam: Government of Assam. 2023. <https://asbb.assam.gov.in/information-services/biodiversity-of-assam>
- Sinha GP, Nayaka S, Mishra GK. A comprehensive checklist of lichens from India – 2024. *Cryptogam Biodivers Assess*. 2024;8(2):1-134.
- Stirton J. On the vegetable parasite of the tea plant, more especially of Assam. *Proc Roy Phil Soc Glasgow*. 1881;13:181-93.
- Awasthi DD, Singh KP. A note on the revision of some Graphidaceous lichen taxa. *Curr Sci*. 1973;42:656-7.
- Pant G, Upreti DK. The lichen genus *Diploschistes* in India and Nepal. *Lichenologist*. 1993;25:33-50. <https://doi.org/10.1006/lich.1993.1011>
- Rout J, Rongmei R, Das P. Epiphytic lichen flora of a pristine habitat (NIT Campus) in southern Assam, India. *Phytotaxonomy*. 2005;5:117-9.
- Rout J, Das P, Upreti DK. Epiphytic lichen diversity in a reserve forest in South Assam, northeast India. *Trop Ecol*. 2010;51:281-8.
- Gupta P, Sinha GP. A new record of lichen in the genus *Graphis* for India from Assam. *Ind J Forest*. 2011;35:133-4. <https://doi.org/10.54207/bsmps1000-2012-33J2LQ>
- Gupta P, Sinha GP. A first note on foliicolous lichens of Assam, India. *J Threat Taxa*. 2016;8:9014-23. <https://doi.org/10.11609/jott.1882.8.7.9014-9023>
- Sinha GP, Gupta P, Jagadeesh Ram TAM, Solanki CM. A contribution

to the lichen flora of Assam, India. *Ind J Forest*. 2013;36:393-400. <https://doi.org/10.54207/bsmps1000-2013-18746D>

- Daimari R, Hazarika N, Houque RR, Upreti DK, Nayaka S. New records of epiphytic lichens from three districts of Assam. *Indian Forest*. 2014;140:807-11.
- Daimari R, Islary P, Brahma S, Basumatary S, Goswami JP, Nayaka S, et al. Additions to the lichen flora of Assam with special reference to *Astrothelium alboverrucoides*, a new record to India. *Stud Fungi*. 2021;6:342-51. <https://doi.org/10.5943/sif/6/1/25>
- Gupta P, Sinha GP. Lichen flora of Assam. *Journal of Forestry Add Series V*. Dehradun: Bishen Singh Mahendra Pal Singh. 2018.
- Biswas S, Daimari R, Islary P, Nayaka S, Joseph S, Upreti DK, et al. New additions to the lichen biota of Assam from Dhubri district, northeastern India. *J Threat Taxa*. 2022;14:21084-90. <https://doi.org/10.11609/jott.7606.14.5.21084-21090>
- Gogoi R, Yasmin F, Nayaka S. Addition to the lichen biota of undivided Nagaon district with special reference to *Tylophoron protrudens* Nyl., a new record to Assam state of India. *Cryptogam Biodivers Assess*. 2022;6:34-8.
- Islary P, Thakur M, Kumar V, Biswas S, Mishra GK, Nayaka S, et al. New additional records of lichenized fungi to Indian lichen biota. *Vegetos*. 2024;37:2035-40. <https://doi.org/10.1007/s42535-024-01036-6>
- Islary P, Daimari R, Nayaka S, Joseph S, Upreti DK, Biswas S. Occurrence of 26 new additional records to the lichen biota of Assam. *Asian J Conserv Biol*. 2021;10:211-21. <https://doi.org/10.53562/ajcb.68956>
- Behera PK, Joseph S, Nayaka S, Chauhan RS. New additions of lichens from Assam, India. *Plant Sci Today*. 2024;11:333-40. <https://doi.org/10.14719/pst.3310>
- Orange A, James PW, White FJ. *Microchemical methods for the identification of lichens*. 2nd ed. London: British Lichen Society; 2010.
- Awasthi DD. *A compendium of the Macrolichens from India, Nepal, Sri Lanka*. 1st ed. Dehradun: Bishen Singh Mahendra Pal Singh; 2007.
- Awasthi DD. *A key to the Microlichens of India, Nepal, Sri Lanka*. *Biblioth Lichenol*. 1991;40:1-340. <https://doi.org/10.2307/1222799>
- Nayaka S. *Revisionary studies on lichen genus Lecanora sensu lato in India* [PhD thesis]. Faizabad (UP): Dr RML Avadh University; 2005.
- Wijayawardene NN, Hyde KD, Al-Ani LKT, Tedersoo L, Haelewaters D, Rajeshkumar KC, et al. Outline of fungi and fungus-like taxa. *Mycosphere*. 2020;11(1):1060-456. <https://doi.org/10.5943/mycosphere/11/1/8>

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.