



REVIEW ARTICLE

Advances in analytical techniques for phytochemical identification and quality control: A comprehensive review

Dwaipayan Sinha^{1*}, Zoofishan Kazi², Aqsa Majgaonkar², Moumita Datta³, Swastika Banerjee⁴, Arun Kumar Maurya^{5*}, Sharmistha Ganguly⁶, Nasrin Parvin¹, Subhamita Sen Niyogi⁷, Shahana Chowdhury⁸, Supti Das⁹, Arpan Dey¹⁰ & Ranjan Shaw¹¹

¹Department of Botany, Government General Degree College, Mohanpur 721 436, India

²Department of Botany, St. Xavier's College (Autonomous), 5, Mahapalika Marg, Mumbai 400 001, India

³Department of Botany, Rammohan College, Kolkata 700 009, India

⁴Department of Science, Alliance University, Bengaluru 562 106, India

⁵Department of Botany, Multanimal Modi College, Ghaziabad 201 204, India

⁶University Department of Botany, Ranchi University, Ranchi 834 008, India

⁷Central water testing laboratory, Office of the Executive Engineer, Rajarhat, Kolkata 700 163, India

⁸German University Bangladesh, T & T Road, Gazipur 1702, Bangladesh

⁹Department of Biochemistry, University of Calcutta, Kolkata 700 019, India

¹⁰Department of Biosciences and Bioengineering, Indian Institute of Technology, Jodhpur 342 030, India

¹¹Central National Herbarium, Botanical Survey of India, Howrah 711 103, India

*Correspondence email - dwaipayansinha@hotmail.com; akmauryahrc@gmail.com

Received: 11 June 2025; Accepted: 03 October 2025; Available online: Version 1.0: 26 November 2025; Version 2.0: 11 December 2025

Cite this article: Dwaipayan S, Zoofishan K, Aqsa M, Moumita D, Swastika B, Arun KM, Sharmistha G, Nasrin P, Subhamita SN, Sahana C, Supti D, Arpan D, Ranjan S. Advances in analytical techniques for phytochemical identification and quality control: A comprehensive review. *Plant Science Today*. 2025; 12(4): 1-11. <https://doi.org/10.14719/pst.9998>

Abstract

Phytochemistry, the study of plant chemicals, has been a longstanding tradition since ancient times, when plants were used as medicines. Early scientists, such as Dioscorides and Theophrastus, made significant contributions to its development and in the 20th century, the introduction of advanced technologies led to substantial progress in this field. Herbal medicine has gained popularity following the advent of technologies such as mass spectrometry (MS) and nuclear magnetic resonance (NMR) spectroscopy, which enable the identification and characterization of bioactive ingredients or phytochemicals. Scientific study is crucial for the authenticity, purity and efficacy of herbal medicines, particularly in light of the growing global demand for them. Herbal medication development follows standardized processes and principles, from plant identification to pharmacological testing. Quality control entails a thorough evaluation that includes identity, validity, physical and chemical characteristics, detection of adulterants and detection of contamination. The World Health Organisation (WHO) and Pharmacopoeias contribute to global standards for herbal drugs, where novel and emerging tools, such as: High-Performance Liquid Chromatography (HPLC), Gas Chromatography-Mass Spectrometry (GC-MS), Deoxyribonucleic Acid (DNA) barcoding and several other advanced techniques, aid in the process. This paper focuses on highlighting some of these contemporary analytical procedures that enable researchers to meet the professional requirements for developing herbal medications. Modern quality control in phytochemistry employs advanced methods, including thermal and chromatographic analysis. These techniques help identify plant chemicals more accurately, making it easier to develop and deliver effective medicines. This paper also provides practical insights for professionals in industries related to herbal medicines, natural products and manufacturers of phytochemical-based products, helping them stay informed about the latest technologies. Moreover, it will help develop guidelines and standards for the quality control and safety assessment of phytochemicals, supporting policymakers and regulatory bodies in making informed and updated decisions regarding the use and marketing of phytochemical products.

Keywords: contaminants; drugs; HPLC; phytochemicals; phytochemistry.

Introduction

Phytochemistry, the exploration of plant chemistry and bioactive compounds, boasts a long and diverse history spanning centuries, including the Chinese and Egyptians, who harnessed the medicinal potential of plants, acknowledging the therapeutic attributes inherent in specific herbs and botanicals. An essential aspect of phytochemical examination, involving the recognition

and characterization of the chemical constituents present in plant extracts or natural products, is the chemical delineation of phytochemicals. The therapeutic qualities and health benefits associated with medicinal plants are primarily attributed to phytochemicals, which are bioactive substances. The chemical characterization of phytochemicals serves several crucial functions. It aids in establishing the presence of specific bioactive components, understanding the chemical complexity and

composition of plant extracts and evaluating the efficacy, consistency and purity of phytochemical-based products (1). In medicinally important plants, the application of thermal analysis techniques has become indispensable for researchers and scientists seeking a comprehensive understanding of the thermodynamic properties and transformations of natural products. The thermal analysis of these plants involves studying the heat-induced changes in their structure and composition (2). This branch of analysis encompasses various techniques, such as differential scanning calorimetry (DSC), thermogravimetric analysis (TGA) and thermal gravimetric-differential thermal analysis (TG-DTA) (3, 4). These methods enable researchers to analyze the thermal transitions, breakdown patterns and phase shifts of medicinal plants and their associated phytochemicals at various temperatures and meteorological conditions. Pharmaceutical and nutraceutical enterprises must understand the heat stability of medicinal plants and phytochemicals, which aids in improved extraction, protection of bioactive ingredients and product purity and efficacy. The thermal research also demonstrates how temperature affects the therapeutic properties of these plants, helping to retain their efficacy throughout storage (5).

The 20th century witnessed a revolution in phytochemistry, driven by the development of chemical methods and equipment, which facilitated faster extraction, isolation and purification techniques. These advancements were crucial in identifying a diverse range of bioactive substances (6). The structural elucidation of phytochemicals has seen advancements with the introduction of spectroscopic methods, such as Infrared Spectroscopy (IR), Ultraviolet-Visible Spectroscopy (UV-Vis) and Nuclear Magnetic Resonance Spectroscopy (NMR) (7). Chromatographic techniques, including high-performance thin-layer chromatography (HPTLC) and high-performance liquid chromatography (HPLC), facilitated the isolation and quantification of plant elements (8). Electrospray ionization (ESI) and air pressure chemical ionization (APCI), two recent developments in mass spectrometry (MS), have made it easier to identify and characterize phytochemicals at the molecular level (9).

Techniques such as HP-TLC have also enhanced resolution and separation, facilitating the identification and quantification of individual phytochemical compounds. Using various stationary phases, mobile phases and detection techniques, HP-TLC can proficiently separate and visualize different phytochemical classes, including alkaloids, polyphenols, terpenoids and aromatic compounds (10). On the other hand, HPLC can handle complex plant extracts, raw materials and finished goods. It possesses exceptional separation qualities, enabling the resolution of complex mixtures and the identification of specific phytochemical components (11). Supercritical fluid chromatography (SFC) has garnered considerable interest and recognition in phytochemical quality control due to its unique advantages and capabilities. One of the primary benefits of SFC in phytochemical analysis is its capability to effectively separate a range of compounds, encompassing both polar and non-polar analytes. Variable selectivity and solvating power can be achieved by utilizing a supercritical fluid, such as carbon dioxide (CO₂), by varying temperature, pressure and co-solvent content. Supercritical fluid chromatography is a state-of-the-art option for separating complex mixtures of phytochemicals due to its versatility, which provides high-resolution separations while minimizing matrix interferences (12).

Particularly for volatile and semi-volatile substances, GC-MS is an excellent choice. Gas chromatography (GC) and MS work together to separate complicated mixtures and identify specific substances based on their mass spectra (13). The detection and quantification of trace components in phytochemical samples are made possible by the highly sensitive and specific analysis provided by GC-MS (14). It is frequently used to examine terpenes, essential oils and other volatile substances found in plant materials (15-17). Liquid chromatography-mass spectrometry is more adaptable and versatile in handling various compound classes, including both polar and non-polar compounds (18). While MS provides precise mass determination and structural elucidation of substances, liquid chromatography (LC) delivers effective separation (19). Liquid chromatography-mass spectrometry is particularly useful for detecting low concentrations of non-volatile substances, such as flavonoids, alkaloids and phenolics, which are commonly found in many plant extracts. Its high sensitivity enables the identification and measurement of phytochemicals in low quantities (20). An essential instrument for phytochemical quality control, inductively coupled plasma-mass spectroscopy (ICP-MS) enables the precise and effective measurement of trace elements and heavy metals (HMs) (21). These techniques provide detailed information on compound compositions, concentrations and the presence of contaminants or impurities. Confirming the identity of the compounds under investigation is made easier by comparing experimental data with recognized reference standards, such as authentic compounds or spectral libraries (22). Advancements in scientific instrumentation, computing capabilities and data analysis methodologies have led to the development of modern analytical techniques for quality control and the identification of phytochemicals (23).

The motivation behind writing this paper is to share the latest advancements in analytical techniques for quality control and to disseminate information on modern methods to a broader audience, including researchers, academics and professionals in the field of quality control and chemical identification of phytochemicals. This will contribute to the scientific community by fostering discussions and advancements, providing insights into how these techniques can be applied to verify the authenticity, purity and safety of phytochemical products and encouraging interdisciplinary research by showcasing the integration of modern analytical methods in the study of phytochemicals.

Significance of quality control of medicinal plants

Herbal medications require strict quality control to ensure their safety, potency, purity and consistency. It also entails inspecting the compound or medicine for the presence of adulterants and determining its hazardous/damaging effects on living systems (24). With rising global interest in traditional medicine, there is an increasing demand for medicinal plants, necessitating robust quality control measures. Furthermore, quality control extends to the identification and screening of contaminants, including HMs, pesticides and microbial agents, reinforcing the commitment to maintaining the highest standards in herbal medicine production (24). Quality control addresses this by evaluating plant identity, purity and potency (25). Accurate species identification is crucial, as misidentification can lead to contamination or toxicity (26). DNA barcoding enables precise species authentication by analyzing specific DNA regions (Fig. 1) (27). A study using 1,452 *rbcl* and *matK* barcodes from 521 herbs achieved species




Fig. 1. Workflow of DNA barcoding.

resolution rates of 74.4 % %, 90.2 % % and 93.0 % %, respectively, revealing adulteration in 20 % % of samples, 6 % % of which had toxic properties (27). The resulting barcode library supports routine drug authentication for stakeholders (27).

The variability in growing, storage and processing conditions frequently increases the presence of toxic chemicals, microbial load, contaminants and other undesirable elements in medicinal plant raw materials, making purity essential (28). It is thus imperative to detect and monitor these levels meticulously, to ensure the safety and quality of the medicinal herbs Fig. 2 illustrates the phytochemical quality control process in human health. Overall, DNA barcoding, chromatography and spectroscopy are crucial for ensuring the authenticity, purity and potency of herbal medicines, key to their safe global use.

Development of herbal medicine

Stages in the development of herbal medicines

The first step in developing an herbal medicine is collecting the desired plant material, followed by a proper evaluation of its

identity and authenticity (29). The collected plant is then processed systematically, adhering to standardized parameters. Accurate identification involves taxonomy, nomenclature and family classification, typically verified by experts or recognized authorities, such as herbaria (30). Chemical evaluation helps determine the purity and presence of key plant therapeutic constituents. It includes measuring extractive yield, selecting suitable extraction methods and conducting preliminary phytochemical screening (31). Recent advances utilize marker compounds in chemical fingerprinting to assess the potency of crude materials. This technique identifies characteristic patterns of chemical markers-categorized as bioactive, therapeutic, principal, or toxic components. Chemical fingerprinting enhances quality control by validating the authenticity and detecting adulterants, thereby ensuring the efficacy and safety of herbal products (32). Fig. 3 provides a general outline of the steps involved in standardizing and assuring the quality of medicinal plants, while Fig. 4 depicts the step-by-step process for authenticating plant materials when developing herbal medicines.




Fig. 2. Blueprint of the routine of quality control.




Fig. 3. Outline the methods involved in the standardization and quality assurance of medicinal plants.




Fig. 4. Schematic diagram showing different aspects of identification and authentication of a plant sample.

Tools and techniques used in standardization, quality assurance and control of herbal medicines

Thermal analysis of medicinal plants and phytochemicals: Medicinal plants have been harnessed for their therapeutic potential. In this context, understanding the thermal behavior of their bioactive constituents is crucial for pharmaceutical applications (33). In this section, the various thermal analysis techniques will be discussed.

Differential scanning calorimetry

Differential scanning calorimetry is a thermal analysis tool employed to quantify the heat flow linked with physical and chemical transformations in a substance relative to temperature changes, crystalline structure and phase transitions (34, 35). The main components of a DSC instrument include a sample pan, a reference pan (often an empty pan), a heating element and a temperature controller (Fig. 5) (36). In a standard DSC experiment, the sample and reference pans undergo identical temperature changes and any heat taken up or released by the sample is compared with that of the reference. The temperature variance between the sample and reference is documented as a time- or temperature-dependent variable (37). A study aimed to enhance the gastrointestinal absorption of *Ginkgo biloba* extract (GBE) by creating phospholipid complexes (GBP) and solid dispersions (GBS). The physicochemical properties of the compounds mentioned above were assessed using DSC. Subsequent pharmacokinetic assessments in rats demonstrated a significant enhancement in the oral bioavailability of various flavonoids, including quercetin, when administered in GBP and GBS compared to GBE. Notably, the bioavailability of GBP exhibited a substantial increase compared to GBS, resulting in improved bioavailability (38).

Thermogravimetric analysis

Thermogravimetric analysis is another thermal analysis method used to examine how a material's weight changes with temperature or time under controlled conditions. It is particularly valuable for assessing the thermal stability, composition and decomposition kinetics of diverse substances (39). In a TGA

experiment, the sample and a reference material, often an empty pan, are heated simultaneously and the alteration in the sample's weight is continuously monitored. The temperature is systematically increased and the TGA instrument records the sample's weight loss or gain as it undergoes thermal processes (40). This technique is widely used in research and industry to gain insights into the thermal characteristics of materials and their responses to varying temperatures. A study focused on enhancing the aqueous solubility and bioavailability of *Boswellia serrata* Roxb. extract (BSE), renowned for its anti-inflammatory properties, by incorporating it into a lamellar solid hydrotalcite-like anionic clay (41). The ionic lamellar compound, chosen for its layered crystal structure, was expected to encapsulate boswellic acids, the active components of BSE. The structure was calcined to enhance the loading of organic acids. The resulting hybrid composites were extensively characterized using TGA to understand their characteristics, interplay and estimation of the active molecules within the materials loaded with plant extract (41).

Thermal gravimetric-differential thermal analysis

Thermal Gravimetric-Differential Thermal Analysis is a combined thermal analysis technique that integrates the principles of TGA and DTA, used to simultaneously measure a sample's weight changes (TGA) and temperature (DTA) as it undergoes thermal processes (42). The main components of a TG-DTA instrument include a sample pan, a reference pan, a balance for weight measurements (TGA) and a thermocouple for temperature measurements (DTA). The sample and reference pans are heated simultaneously and the balance records any weight loss or gain of the sample, while the temperature probe notes the temperature variance between the pans (43). A study analyzed the constituents and lipid profile of *Caryocar villosum* oil and assessed its heart health benefits, thermogravimetric-differential, calorimetric and spectroscopic behavior (44).

Dynamic mechanical analysis (DMA)

Dynamic mechanical analysis is a versatile and sophisticated method used to assess the mechanical characteristics of materials in relation to temperature, time, frequency, or a combination of




Fig. 5. Workflow of differential scanning calorimetry.

these factors (45). DMA is particularly valuable in characterizing the viscoelastic behavior of materials, providing insights into their dynamic mechanical response under various conditions (46). A study focused on enhancing the overall stability of natural fiber composites (NFCs), which are recognized for their environmentally beneficial properties, including biodegradability. The research introduces three novel fibers and fillers-*Morinda tinctoria* wood powder and myrobalan-aiming to enhance the performance of biodegradable hybrid composites (47).

Chromatographic analysis

The chromatographic profiles mostly mirror the sample's chemical composition, as observed through the selected detection techniques. The content of this substance relies heavily on determiners such as the specific species, plant component, ambient circumstances and extraction process (31). In chromatographic analysis, the quality and degree of authenticity of herbal preparations are assessed by utilizing one or two marker molecules or pharmacologically active constituents found in the mixtures (48). Since most recent state-of-the-art facilities generate massive amounts of data, which can be challenging to process, the relevance of data-processing techniques that utilize chemometrics is instrumental. Chemometrics utilizes statistics and mathematical concepts to analyze data, thereby optimizing the collection and application of valuable knowledge (49). The efficacy of chemometric techniques in facilitating multidimensional calibration of specific spectroscopic, electrochemical and chromatographic procedures is exemplified. The primary application of this approach is to interpret UV-Vis and near-IR (NIR) spectra, as well as data obtained from other instrumental methods, which enables the characterization and quantitative analysis of active substances in complex mixtures, particularly in the study of medicinal preparations available on the market (49).

High-performance thin-layer chromatography analysis

Currently, one of the most effective analytical methods for analysing phytochemicals is HPTLC, which offers superior

separation efficiency when identifying and detecting complex biomolecules in plant extracts used in botanical research and herbal medicines (50). The benefits of HP-TLC are scanning, optimization, automation, computerization, selective detection and minimal sample preparation (51). Table 1 presents selected studies that utilize HPTLC for detecting marker molecules and evaluating the quality of herbal formulations.

High-performance liquid chromatography analysis

High-performance liquid chromatography is a user-friendly, highly automated and widely used technique for the quality assessment and standardization of herbal preparations, due to its capability to identify the presence of markers in both qualitative and quantitative manners (57). An important advantage of HPLC is the ability to perform hyphenation with various detecting molecules like diode array detection (DAD), fluorescence detection (FD), evaporative light scattering detection (ELSD), MS and computer-aided design (CAD) (58-62). The working procedure of HPLC is depicted in Fig. 6, while its applications in assessing polyherbal formulations and herbal preparations are illustrated in Table 2.

Liquid chromatography mass spectrometry (LC-MS) analysis


High performance liquid chromatography alone has limitations in analyzing complex raw extracts, which are overcome by integrating LC-MS, enabling enhanced accuracy through combined separation and mass detection. Liquid chromatography mass spectrometry offers a discriminating method that enables the measurement of both quantity and structure and can attain very low sensitivities of picograms per milliliter (68). The LC-MS-ion trap mass spectrometry method can be simplified by coupling it with quadrupole time-of-flight high-resolution mass spectrometry (Q-TOF HRMS) and triple-quadrupole mass spectrometry (TQ-MS), alternative techniques that can be used in conjunction with the HPLC analytical approach (69-71). The LC-MS approach possesses unique capabilities, such as structure characterization, determination of molecular mass, fragmentation information,

Table 1. Evaluation of selected polyherbal formulations and plant-based drugs by HPTLC

Name of the plant material/formulation/drug	Major constituents	Pharmacological properties	Important compounds detected	Major outcome of the investigation	Reference
Sanwujiao pills	Polygoni Multiflori Radix, Aconiti Radix, Aconiti Lateralis Radix Praeparata, Aconiti Vilmoriniani Radix, Olibanum, Typhonii Rhizoma	Treatment of rheumatism and blood stasis	Emodin, Physcion, 2,3,5,4'-tetrahydroxy stilbene-2-O- β -D-glucoside, Benzoylaconine, Benzolymesaconine, β -Boswellic acid and 3-acetyl- β -boswellic acid	Qualitative analysis of intricate herbal preparations and detection of phytochemicals.	(52)
Divya-swasari-vati	<i>Glycyrrhiza glabra</i> , <i>Syzygium aromaticum</i> , <i>Cinnamomum zeylanicum</i> , <i>Pistacia integerrima</i> , <i>Cressa cretica</i> , <i>Zingiber officinale</i> , <i>Piper nigrum</i> , <i>Piper longum</i> , <i>Anacyclus pyrethrum</i>	Treatment of pulmonary ailments	Gallic acid, Glycyrrhizin and Eugenol	The swift standardization of Divya-Swasari-Vati.	(53)
Jatyadi taila	<i>Curcuma longa</i> , <i>Terminalia chebula</i> and <i>Jasminum officinale</i>	Antiseptic, treatment of wounds	Curcumin, Gallic acid and Ursolic acid	Standardized method for quality control and determination of stability of the marker compounds.	(54)
Mahasudarshan Churna	<i>Swertia chirata</i>	Treatment of fever, cold and malaria	Oleanolic acid, Mangiferin, Gallic acid, Ursolic acid, Curcumin, Quercetin	An economical, rapid, precise and accurate method to measure the marker chemicals in the formulations.	(55)
<i>Centella asiatica</i> and its commercial products	<i>Centella asiatica</i>	Treatment of the nervous system	Asiaticoside	Quantification of Asiaticoside in the formulation.	(56)

Table 2. Evaluation of selected polyherbal formulations and plant-based drugs by HPLC

Name of the plant material/formulation/ drug	Major constituents	Pharmacological properties	Important compounds detected	Major outcome of the investigation	Reference
Brahmi vati	<i>Bacopa monnieri</i> L. and <i>Piper longum</i> L.	Treatment of epilepsy	Bacoside A ₃ and Piperine	Substantial increase in bacoside A ₃ and piperine in the in-house sample compared to all three marketed samples.	(63)
Trichup tablet	<i>Eclipta alba</i> , <i>Glycyrrhiza glabra</i> , <i>Emblica officinalis</i> , <i>Centella asiatica</i> , <i>Hibiscus rosa-sinensis</i> , <i>Tinospora cordifolia</i> , <i>Tribulus terrestris</i> , Powder of <i>Triphala Churna</i> and <i>Shukti Bhasma</i>	Treatment of hair and scalp	Gallic acid	Quick and accurate estimation of gallic acid in the sample.	(64)
Qurs-e-Gul and crude drug	<i>Glycyrrhiza glabra</i> , <i>Pistacia lentiseus</i> , <i>Bamboosa bamboo</i> , <i>Nardostachys jatamansi</i> , <i>Rosa damascene</i>	Treating jaundice and cardiac ailments	Glabridin	Accurate estimation of glabridin in the sample.	(65)
Ashwagandhadi lehyam	<i>Withania somnifera</i> , <i>Hemidesmus indicus</i> , <i>Cuminum cyminum</i> , <i>Smilax china</i> , <i>Vitis vinifera</i> , <i>Elettaria cardamomum</i>	Used as a tonic, hypnotic, sedative and diuretic	Withaferin-A	The Ashwagandha lehyam formulation has a concentration of 0.092 % % of withaferin-A.	(66)
Pathyashadangam kwath	<i>Terminalia chebula</i> , <i>Terminalia bellirica</i> , <i>Phyllanthus emblica</i> , <i>Andrographis paniculata</i> , <i>Tinospora cordifolia</i> , <i>Curcuma longa</i> , <i>Azadirachta indica</i>	Treatment of diseases of the eye, ear and tooth, migraine and cluster headache	Gallic acid	Quantification of the gallic acid in three batches of the sample (16.14 mg/mL, 23.22 mg/mL and 19.29 mg/mL) of kwath, respectively.	(67)

**Fig. 6.** Workflow of high-performance liquid chromatography.

retention time, a broad detection range and excellent separation of analytical substances that are not present in HPLC alone (72-75). The working principle of LC-MS is represented in Fig. 7, while Table 3 illustrates the identification and quality evaluation of raw plant products and polyherbal formulations using the technique.

Gas chromatography-mass spectrometry analysis

Gas chromatography is used to measure the concentration of volatile compounds in plant matrices (81). The benefits of GC include high sensitivity, low sample requirements, robust separation capabilities, excellent selectivity, broad applicability and rapid analysis (82). Gas chromatography separates mixture components based on their interactions with a stationary phase and a mobile carrier gas. Variations in the compounds' structures and affinities cause them to elute at different times under the same driving force (82). The motivation behind coupling a GC (83) with an MS was to combine the complementary features of these two analytical tools. The GC separated components of mixtures into distinct pure compounds, which eluted progressively. On the contrary, the MS offered a strong approach to detecting previously undiscovered pure compounds (84). Gas chromatography-mass spectrometry is a method that combines two analytical techniques to detect and characterize chemicals found in a plant sample. Gas chromatography-mass spectrometry is a valuable tool for analyzing phytochemicals and studying chemotaxonomy in medicinal plants, as it enables the generation of high-quality chemical fingerprints using capillary columns (85). Additionally, GC-MS provides qualitative and semi-quantitative data through mass spectral databases, aiding in research into the pharmacological relevance of

herbal constituents (86). The working mechanism of GC-MS is illustrated in Fig. 8, while its application in the quality evaluation and standardization of herbal medicine is exemplified in Table 4.

Supercritical fluid chromatography analysis

Supercritical fluid chromatography (SFC) utilizes a low-viscosity supercritical fluid, typically CO₂, as the mobile phase to dissolve primarily nonpolar compounds (92). By adding cosolvents like methanol or ethanol, SFC can also analyze polar analytes, offering superior separation efficiency and faster analysis than LC and effectively separating heat-sensitive molecules, unlike GC (93). Its lower use of organic solvents makes it more environmentally friendly and its low viscosity results in reduced pressure drops compared to traditional LC (94). SFC is a versatile method for separating, quantifying and assessing the quality of both polar and nonpolar natural compounds in herbal products. Table 5 illustrates the application of SFC in the quality assessment of the herbal tenets.

Technique for the analysis of elements

Inductively coupled plasma mass spectrometry (ICP-MS) analysis

Inductively Coupled Plasma Mass Spectrometry (ICP-MS) is employed for elemental analysis (28). It is a reliable analytical method for qualitative multi-elemental determination, quantitative concentration measurement and isotopic abundance assessment across diverse matrices. An ICP-MS system typically includes a sample introduction system, an ion source (ICP), electrostatic lenses, an interface, a mass spectrometer and a detector (98). In a standard setup, samples are introduced via an autosampler, then

Table 3. Evaluation of selected polyherbal formulations and plant-based drugs by LC-MS

Name of the plant material/formulation/drug	Major constituents	Pharmacological properties	Important compounds detected	Major outcome of the investigation	Reference
Majoon-e-Nisyan	<i>Boswellia serrate</i> Roxb., <i>Acorus calamus</i> L., <i>Cyperus rotundus</i> L., <i>Piper nigrum</i> L., <i>Zingiber officinale</i> L.	Treatment of amnesia	($\alpha + \beta$) boswellic acid, β -asarone, luteolin, 6-gingerol and piperine	Boswellic acid ($\alpha + \beta$), β -asarone, luteolin, 6-gingerol and piperine were quantified as follows: 0.973 ± 0.009 , 0.093 ± 0.004 , 0.003 ± 0.001 , 0.0256 ± 0.002 and 0.0266 ± 0.008 mg/g of the formulation.	(76)
Peedantak Vati	<i>Balsamodendron mukul</i> , <i>Colchicum luteum</i> , <i>Withania somnifera</i> , <i>Strychnos nux-vomica</i> , <i>Cyperus scariosus</i> , <i>Pluchea lanceolata</i> , <i>Vitex negundo</i> , <i>Boerhaavia diffusa</i>	Treatment of inflammation	Rutin, Caffeic acid, Colchicine, Withaferin A and Curcumin	Confirmation of the presence of marker phytochemicals.	(77)
YANG XIN	<i>Angelica sinensis</i> , <i>Cuscuta chinensis</i> , <i>Nelumbo nucifera</i> , <i>Dioscorea batatas</i> , <i>Cistanche salsa</i>	Improve cognitive function	4,5 Dicafeoylquinic acid, Forsythoside A, Ginsenoside R, Sibiricoses A5 and Schisandrol A,	18 analytical markers were identified for YANG XIN formulations.	(78)
Andrographis paniculata Capsules	<i>Andrographis paniculata</i>	Treatment of fever and diarrhoea	Andrographolide, Dehydroandrographolide and Neoandrographolide	In <i>A. paniculata</i> capsules, the amounts of andrographolide, dehydroandrographolide and neoandrographolide were found to be 3.90-4.08, 4.77-5.04 and 4.32-4.48 mg/g, respectively.	(79)
Hyeonggaeyeongyo-tang	<i>Schizonepetae spica</i> , <i>Forsythiae fructus</i> , <i>Saposhnikoviae radix</i> , <i>Angelicae gigantis radix</i> , <i>Cnidii rhizoma</i> , <i>Bupleuri radix</i> , <i>Aurantii fructus</i> <i>Immaturus scutellariae radix</i> , <i>Angelicae dahuricae radix</i> , <i>Platycodonis radix</i> , <i>Paeoniae radix</i> , <i>Gardeniae fructus</i> and <i>Glycyrrhizae radix et rhizome</i>	Antioxidant and antimicrobial properties	Gallic acid, Narirutin, Baicalin, Neohesperidin, Paeoniflorin, Naringin, Pulegone, Glycyrrhizin and Wogonoside	The marker compounds were effectively estimated.	(80)




Fig. 7. Workflow of liquid chromatography-mass spectrometry.




Fig. 8. Workflow of gas chromatography-mass spectrometry.

Table 4. Evaluation of some polyherbal formulations and polyherbal formulations by GC-MS


Name of the plant material/formulation/ drug	Major constituents	Pharmacological properties	Important compounds detected	Major outcome of the investigation	Reference
Succus Bambusae	<i>Succus Bambusae</i>	Acts as a natural beverage		Twenty-nine volatile components were initially discovered from a sample of forty batches and six quality markers were identified.	(87)
<i>Clinacanthus nutans</i>	<i>Clinacanthus nutans</i>	Treatment of diabetes, inflammatory disease, skin disease, etc.	Squalene, stigmasterol, betulin,	Identification of chemical markers of the plant for future quality evaluation.	(88)
Aswagandharishtam	<i>Withania somnifera</i> , <i>Chlorophytum tuberosum</i> , <i>Rubia cordifolia</i> , <i>Terminalia chebula</i> , <i>Curcuma longa</i> , <i>Berberis aristata</i> , <i>Glycyrrhiza glabra</i> etc.	Used as a nerve tonic	Prostaglandin A2, cholesterol, piperine, gentamicin a, d-mannose, eugenol and pipradrol	A preliminary report was of indication of the different types of biomolecules contained in Aswagandharishtam.	(89)
Arq-e-Nana (AeN), Arq-e-Gazar (AeG) and Arq-e-Brinjasif (AeB)	<i>Mentha Arvensis</i> , <i>Daucus carota</i> , <i>Achillea mellifolium</i>	Treatment of vomiting, dyspepsia, palpitation, inflammation of the liver and intestine	Limonene, 1,8 cineole, fenchone, camphor, citronellol, terpineol, thymol and carvone	Quality control and stability analysis of Arq formulations from the Unani Pharmacopoeia of India.	(90)
Safoof-e-Pathar Phori	<i>Didymocarpus pedicellata</i> , <i>Dolichos biflorus</i> , <i>Rheum emodi</i> , <i>Raphanus sativus</i>	Antiurolithiatic activity	3-furano carboxylic acid, cyclooctene, linolenic acid, n-hexadecanoic acid and alpha humulene	An authentic tool for quality analysis of the polyherbal formulation.	(91)

Table 5. Application of SFC in the quality assessment of herbal principles

Name of the herbal drug	Pharmacological properties	Important elements detected	Major outcome of the investigation	Reference
<i>Rhodiola rosea</i>	Treatment for stress, fatigue and weakness	p-tyrosol, rosin, rosidin, salidroside, rosin, rosin and tricin-5-O- β -d-glucopyranoside	The established methodology enabled a comprehensive extraction of all seven chemicals using a mixture of 60 % CO_2 and 40 % methanol.	(95)
Isatidis Radix (root of <i>Isatis indigotica</i>)	Treatment of cold, fever, sore throat, mumps and tonsillitis	(R, S)-goitrin	This endeavor investigated the utilization of SFC in the field of chiral research about traditional Chinese medicine. The results suggest that SC- CO_2 and SC- CO_2 + ethanol extraction methods show promise as alternatives for extracting artemisinin.	(96)
<i>Artemisia annua</i>	Antimalarial activity	Artemisin	These methods avoid the need for organic solvents like hexane and yield extracts suitable for making antimalarial pharmaceuticals.	(97)

directed by a peristaltic pump to the nebulizer, where they are combined with argon gas to form an aerosol. The spray chamber eliminates large droplets and controls the liquid flow (99). The fine aerosol is then ionized in high-temperature argon plasma through vaporization, atomization and ionization. The extent of ionization depends on plasma temperature and the ionization potential of elements (100). ICP-MS, utilizing an argon plasma, converts the sample into ions and analyzes them using a mass spectrometer

(21). Unlike ICP-OES, which uses optical detection, ICP-MS directly quantifies ions and offers lower detection limits (101). It detects trace elements in ppb or ppt levels and provides advantages such as a wide linear range and isotopic analysis (101). ICP-MS is extensively used to determine in medicinal plants, addressing global health risks when levels exceed threshold limits (102). Fig. 9 illustrates the working principle of ICP-MS, while Table 6 shows the application of ICP-MS in quantifying HMs and assessing the

**Fig. 9.** Workflow of inductively coupled plasma-mass spectrometry.**Table 6.** Evaluation of selected herbal formulations or medicinal plants by ICP-MS

Name of the plant material/ formulation/drug	Pharmacological properties	Important elements detected	Investigation outcomes	Reference
Niu Huang Qingwei Pills	Treatment of gastrointestinal problems	Excess of lead (Pb), arsenic (As) and Antimony (Sb)	A set of fundamental procedures for evaluating the potential risks associated with Chinese patent medicine has been developed for the first time.	(103)
<i>Astragalus membranaceus</i>	Improvement function of the immune system	The computed recommended maximum residue limits (MRLs) for Pb, cadmium (Cd), mercury (Hg) and copper (Cu) in the aforementioned medical items were found to be higher than the MRLs established by the International Organization for Standardization (ISO) and the Chinese Pharmacopoeia.	This study employed a pragmatic research methodology to assess the safety of medicinal herbs contaminated with various HMs.	(104)
Ayurvedic dietary supplements	Multifarious health benefits largely boost the immune system	The samples of dietary supplements exhibited Hg levels ranging from 0.002 to 56 $\mu\text{g/g}$.	An effective way of estimating metallic content in ayurvedic formulations.	(105)
Qishiwei Zhenzhu Pills (Tibetan medicine)	Treatment of stroke, paralysis, hemiplegia, cerebral hemorrhage	The elements in Qishiwei Zhenzhu tablets were ranked in the order of $\text{Cu} > \text{Hg} > \text{Pb}$, with mass fractions above 6000 $\mu\text{g/kg}$. The mass fractions of silver, arsenic, manganese, gold, strontium, barium, chromium and nickel ranged from 33 to 1034 $\mu\text{g/kg}$.	Analysis of trace elements in Qishiwei Zhenzhu Pills.	(106)
Vaiśvānaracūrṇa Tablet	Treatment of flatulence, duodenal ulcers, rheumatism and cardiac ailments	Presence of Cd, Pb, Cr, As, Cu and Zn within the permissible limits	Effective quality evaluation of the polyherbal formulation.	(107)

quality of selected herbal samples.

Conclusion

The development of modern analytical techniques for quality control and chemical identification of phytochemicals represents a pivotal stride in the convergence and revolution of traditional herbal medicine. It has deciphered the intricate compositions of diverse plant extracts with precision and reliability, assessing the quality and authenticity of phytochemical-based products through enhanced phytochemical analysis, quality control, quantification and evaluation. This ensures industries have powerful tools to ensure product integrity and safety. These modern analytical techniques have helped in isolating new bioactive compounds, formulating herbal medicines and advancing natural product research, particularly through the use of thermal analysis and chromatographic methods. The synergy between chromatography and thermal analysis has facilitated the comprehensive profiling of phytochemical mixtures, aiding in the detection and separation of specific compounds crucial for the expansion of pharmaceuticals, nutraceuticals and other applications. Such steps are essential to quality control because they ensure that herbal goods adhere to legal requirements and are free from harmful substances. Rigorous quality assurance protocols, facilitated by cutting-edge analytical tools, safeguard consumer health and bolster the credibility of phytochemical-based interventions in diverse sectors. Lastly, chemometrics and sophisticated data analysis techniques are essential to quality control. The interpretation of complicated data sets, classification of samples and detection of outliers or adulterants are all made conceivable by using statistical tools, multivariate analysis and pattern recognition techniques. These techniques improve the consistency and precision of quality control procedures. In recent years, artificial intelligence and bioinformatics have enhanced the quality of herbal drug development through in-silico studies, utilizing software for screening phytochemicals and molecular docking to determine their efficacies.

Like in any scientific discipline, there are still issues to be resolved. Among the active areas of study and development in this subject are the complexity of plant matrices, sample preparation, the need for standardization and the use of reference materials. Overall, the multidisciplinary subject of chemical identification of phytochemicals uses analytical methods, spectral analysis and reference standards. It is essential to comprehend the chemical complexity of plants, discover novel bioactive substances, ensure product quality, advance the study of natural products and identify new drugs. As technology advances, these methods will remain essential in releasing the full potential of phytochemicals for various uses, from medicine to agriculture and beyond. In essence, the integration of modern analytical techniques with the rich heritage of traditional herbal wisdom strengthens the foundations of phytochemical research. It propels the field toward new horizons of discovery and application. These findings serve as a testament to the transformative power of analytical innovation in unraveling the secrets of nature's pharmacopoeia, offering a roadmap for future endeavors in harnessing the full extent of phytochemicals beneficial to mankind.

Acknowledgements

The authors are grateful to the libraries of their respective institutes and the information centres for providing the requisite information and assisting in the manuscript preparation.

Authors' contributions

DS conceptualized and supervised the manuscript and participated in the composition of the original draft, internal reviewing and overall coordination. ZK, AM, MD, SG, SD, SSN, AD and RS contributed to the sectional composition, editing and revision of the manuscript. AKM edited the manuscript, provided overall supervision and contributed to the artwork. SB was responsible for quality checking, formatting and revision. NP and SC contributed to the revision and reference management of the manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflicts of interest to declare.

Ethical issues: None

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used Grammarly Premium, Quillbot Premium and Chat GPT to remove plagiarism, grammatical, sentence construction and typographical errors. After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the publication.

References

1. Sasidharan S, Chen Y, Saravanan D, Sundram KM, Yoga Latha L. Extraction, isolation and characterization of bioactive compounds from plants' extracts. *Afr J Tradit Complement Altern Med*. 2011;8 (1):1-10.
2. Xue Y, Hu W, Hu X. Thermal analysis of natural fibers. In: Jaffe M, Menczel JD, editors. *Thermal analysis of textiles and fibers*. Cambridge: Woodhead Publishing; 2020. p. 105-32. <https://doi.org/10.1016/B978-0-08-100572-9.00007-0>
3. Peltonen L. Practical guidelines for the characterization and quality control of pure drug nanoparticles and nano-cocrystals in the pharmaceutical industry. *Adv Drug Deliv Rev*. 2018;131:101-15. <https://doi.org/10.1016/j.addr.2018.06.009>
4. Alvarenga LM, Xavier TP, Barrozo MAS, Bacelos MS, Lira TS. Dynamic analysis of reaction kinetics of carton packaging pyrolysis. In: Karimi IA, Srinivasan R, editors. *Computer aided chemical engineering*. Vol. 31. Amsterdam: Elsevier; 2012. p. 180-4. <https://doi.org/10.1016/B978-0-444-59507-2.50028-7>
5. Costa RS, Pinheiro WBdS, Arruda MSP, Costa CEF, Converti A, Costa RMR, et al. Thermoanalytical and phytochemical study of the cupuassu (*Theobroma grandiflorum* Schum.) seed by-product in different processing stages. *J Therm Anal Calorim*. 2022;147:275-84. <https://doi.org/10.1007/s10973-020-10347-0>
6. Sinha D, Odoh UE, Ganguly S, Muhammad M, Chatterjee M, Chikeokwu I, et al. Phytochemistry, history and progress in drug discovery. In: Egbuna C, Rudrapal M, Tijjani H, editors. *Phytochemistry, computational tools and databases in drug discovery*. 1st ed. Amsterdam: Elsevier; 2023. p. 1-26. <https://doi.org/10.1016/B978-0-323-90593-0.00001-0>

7. Altemimi A, Lakhssassi N, Baharlouei A, Watson DG, Lightfoot DA. Phytochemicals: Extraction, isolation and identification of bioactive compounds from plant extracts. *Plants* (Basel). 2017;6(4):42. <https://doi.org/10.3390/plants6040042>
8. Loescher CM, Morton DW, Razic S, Agatonovic-Kustrin S. High performance thin layer chromatography (HPTLC) and high performance liquid chromatography (HPLC) for the qualitative and quantitative analysis of *Calendula officinalis*-advantages and limitations. *J Pharm Biomed Anal*. 2014;98:52-9. <https://doi.org/10.1016/j.jpba.2014.04.023>
9. Gallagher RT, Balogh MP, Davey P, Jackson MR, Sinclair I, Southern LJ. Combined electrospray ionization-atmospheric pressure chemical ionization source for use in high-throughput LC-MS applications. *Anal Chem*. 2003;75(4):973-7. <https://doi.org/10.1021/ac0205457>
10. Attimarad M, Ahmed KK, Aldhubaib BE, Harsha S. High-performance thin layer chromatography: A powerful analytical technique in pharmaceutical drug discovery. *Pharm Methods*. 2011;2(2):71-5. <https://doi.org/10.4103/2229-4708.84436>
11. Malviya R, Bansal V, Pal O, Sharma P. High performance liquid chromatography: A short review. *J Glob Pharma Technol*. 2010;2:22-6.
12. Kupnik K, Knez Ž, Primožič M, Leitgeb M. Separation of amino acids and peptides with supercritical fluids chromatography. *Sep Purif Rev*. 2022;52(1):58-74. <https://doi.org/10.1080/15422119.2022.2038625>
13. Profumo A, Gorrioni A, Guarnieri SA, Mellerio GG, Cucca L, Merli D. GC-MS qualitative analysis of the volatile, semivolatile and volatilizable fractions of soil evidence for forensic application: A chemical fingerprinting. *Talanta*. 2020;219:121304. <https://doi.org/10.1016/j.talanta.2020.121304>
14. Konappa N, Udayashankar AC, Krishnamurthy S, Pradeep CK, Chowdappa S, Jogaiah S. GC-MS analysis of phytoconstituents from *Amomum nilgircum* and molecular docking interactions of bioactive serverogenin acetate with target proteins. *Sci Rep*. 2020;10:16438. <https://doi.org/10.1038/s41598-020-73442-0>
15. Joy N, Jackson D, Coolong T. A validated GC-MS method for major terpenes quantification in hydrodistilled *Cannabis sativa* essential oil. *Phytochem Anal*. 2025. <https://doi.org/10.1002/pca.3526>
16. Sedikelo GK, Lenetha GG, Malebo NJ. Chromatography-mass spectrometry and chemical characteristics of *Thymus zygis* and *Cymbopogon winterianus* essential oils: Possible insect repellents. *Sci Afr*. 2022;15:e01095. <https://doi.org/10.1016/j.sciaf.2022.e01095>
17. Sharma A, Rai PK, Prasad S. GC-MS detection and determination of major volatile compounds in *Brassica juncea* L. leaves and seeds. *Microchem J*. 2018;138:488-93. <https://doi.org/10.1016/j.microc.2018.01.015>
18. Hird SJ, Lau BPY, Schuhmacher R, Krska R. Liquid chromatography-mass spectrometry for the determination of chemical contaminants in food. *TrAC Trends Anal Chem*. 2014;59:59-72. <https://doi.org/10.1016/j.trac.2014.04.005>
19. Kostianinen R, Kaupilla TJ. Effect of eluent on the ionization process in liquid chromatography-mass spectrometry. *J Chromatogr A*. 2009;1216(4):685-99. <https://doi.org/10.1016/j.chroma.2008.08.095>
20. Mukherjee PK. LC-MS: a rapid technique for understanding the plant metabolite analysis. In: Mukherjee PK, editor. *Quality control and evaluation of herbal drugs*. Elsevier; 2019. p. 459-79. <https://doi.org/10.1016/B978-0-12-813374-3.00011-9>
21. Wilschefske SC, Baxter MR. Inductively coupled plasma mass spectrometry: introduction to analytical aspects. *Clin Biochem Rev*. 2019;40(3):115-33. <https://doi.org/10.33176/AACB-19-00024>
22. Stein S. Mass spectral reference libraries: an ever-expanding resource for chemical identification. *Anal Chem*. 2012;84(17):7274-82. <https://doi.org/10.1021/ac301205z>
23. Verma N. Advances and trends in analytical techniques in natural product research: challenges and future perspective. *Indian J Nat Prod Resour*. 2021;12(4):506-26.
24. van Wyk AS, Prinsloo G. Health, safety and quality concerns of plant-based traditional medicines and herbal remedies. *S Afr J Bot*. 2020;133:54-62. <https://doi.org/10.1016/j.sajb.2020.06.031>
25. Govindaraghavan S, Sucher NJ. Quality assessment of medicinal herbs and their extracts: criteria and prerequisites for consistent safety and efficacy of herbal medicines. *Epilepsy Behav*. 2015;52:363-71. <https://doi.org/10.1016/j.yebeh.2015.03.004>
26. Unnikrishnan R, Dev SA, Jayaraj R. Pitfalls and promises of raw drug identification techniques in the ayurvedic industry: an overview. *3 Biotech*. 2020;10:497. <https://doi.org/10.1007/s13205-020-02482-0>
27. Nithaniyal S, Vassou SL, Poovitha S, Raju B, Parani M. Identification of species adulteration in traded medicinal plant raw drugs using DNA barcoding. *Genome*. 2017;60(2):139-46. <https://doi.org/10.1139/gen-2015-0225>
28. Kumar N, Kulsoom M, Shukla V, Kumar D, Priyanka, Kumar S, Tiwari J, Dwivedi N. Profiling of heavy metal and pesticide residues in medicinal plants. *Environ Sci Pollut Res*. 2018;25:29505-10. <https://doi.org/10.1007/s11356-018-2993-z>
29. Wang H, Chen Y, Wang L, Liu Q, Yang S, Wang C. Advancing herbal medicine: enhancing product quality and safety through robust quality control practices. *Front Pharmacol*. 2023;14:1265178. <https://doi.org/10.3389/fphar.2023.1265178>
30. Culley TM. Why vouchers matter in botanical research. *Appl Plant Sci*. 2013;1(11):1300076. <https://doi.org/10.3732/apps.1300076>
31. Klein-Junior LC, de Souza MR, Viaene J, Bresolin TMB, de Gaspar AL, Henriques AT, et al. Quality control of herbal medicines: from traditional techniques to state-of-the-art approaches. *Planta Med*. 2021;87(12/13):964-88. <https://doi.org/10.1055/a-1529-8339>
32. Fan XH, Cheng YY, Ye ZL, Lin RC, Qian ZZ. Multiple chromatographic fingerprinting and its application to the quality control of herbal medicines. *Anal Chim Acta*. 2006;555(2):217-24. <https://doi.org/10.1016/j.aca.2005.09.037>
33. Pirsaeheb M, Seifi H, Dawi EA, Gholami T, Badraddin SQ, Ryadh A, Salavati-Niasari M. Thermal analysis techniques in herbal medicine: a comprehensive review on unveiling integrity and quality for future perspectives. *J Anal Appl Pyrolysis*. 2023;175:106192. <https://doi.org/10.1016/j.jaap.2023.106192>
34. Gill P, Moghadam TT, Ranjbar B. Differential scanning calorimetry techniques: applications in biology and nanoscience. *J Biomol Tech*. 2010;21(4):167-93
35. Lukas K, LeMaire PK. Differential scanning calorimetry: fundamental overview. *Reson*. 2009;14:807-17. <https://doi.org/10.1007/s12045-009-0076-7>
36. Odarchenko Y, Kaźmierczak-Bałata A, Bodzenta J, Ferrari E, Soloviev M. AC/DC thermal nano-analyzer compatible with bulk liquid measurements. *Nanomaterials* (Basel). 2022;12(21):3799. <https://doi.org/10.3390/nano12213799>
37. Haines PJ, Reading M, Wilburn FW. Differential thermal analysis and differential scanning calorimetry. In: Brown ME, editor. *Handbook of thermal analysis and calorimetry*. Vol. 1. Amsterdam: Elsevier Science B.V.; 1998. p. 279-361. [https://doi.org/10.1016/S1573-4374\(98\)80008-3](https://doi.org/10.1016/S1573-4374(98)80008-3)
38. Chen ZP, Sun J, Chen HX, Xiao YY, Liu D, Chen J, Cai H, Cai BC. Comparative pharmacokinetics and bioavailability studies of quercetin, kaempferol and isorhamnetin after oral administration of *Ginkgo biloba* extracts, *Ginkgo biloba* extract phospholipid complexes and *Ginkgo biloba* extract solid dispersions in rats. *Fitoterapia*. 2010;81(8):1045-52. <https://doi.org/10.1016/j.fitote.2010.06.028>
39. Ng HM, Saidi NM, Omar FS, Ramesh K, Ramesh S, Bashir S.

- Thermogravimetric analysis of polymers. In: Encyclopedia of Polymer Science and Technology. Hoboken: John Wiley & Sons, Inc.; 2002. <https://doi.org/10.1002/0471440264.pst667>
40. Lothenbach B, De Weerd K. Thermogravimetric analysis. In: Scrivener K, Snellings R, Lothenbach B, editors. A practical guide to microstructural analysis of cementitious materials. Boca Raton: CRC Press; 2016
 41. Cometa S, Busto F, Castellana A, Cochis A, Najmi Z, Rizzi R, et al. Development, analytical characterization and bioactivity evaluation of *Boswellia serrata* extract-layered double hydroxide hybrid composites. *Molecules*. 2023;28(18):6449. <https://doi.org/10.3390/molecules28186449>
 42. Chiu J. Technique for simultaneous thermogravimetric, derivative thermogravimetric, differential thermal and electrothermal analyses. *Anal Chem*. 1967;39(8):861-7. <https://doi.org/10.1021/ac60252a042>
 43. Hoffman R, Pan WP. Measuring ΔH using TG-DTA and incorporating mass change into the result. *Thermochim Acta*. 1991;192:135-46. [https://doi.org/10.1016/0040-6031\(91\)87155-P](https://doi.org/10.1016/0040-6031(91)87155-P)
 44. Lorenzo ND, Santos OVD, Lannes SCDS. Fatty acid composition, cardiovascular functionality, thermogravimetric-differential, calorimetric and spectroscopic behavior of pequi oil (*Caryocar villosum* (Alb.) Pers.). *Food Sci Technol*. 2021;41(2):524-9. <https://doi.org/10.1590/fst.16420>
 45. Saba N, Jawaid M, Allothman OY, Paridah MT. A review on dynamic mechanical properties of natural fibre reinforced polymer composites. *Constr Build Mater*. 2016;106:149-59. <https://doi.org/10.1016/j.conbuildmat.2015.12.075>
 46. Bashir MA. Use of dynamic mechanical analysis (DMA) for characterizing interfacial interactions in filled polymers. *Solids*. 2021;2(1):108-20. <https://doi.org/10.3390/solids2010006>
 47. Sathees Kumar S, Mugesh Raja V, Sudhagar S, Kanagaraj G, Vignesh V, Manimaran P. Static, dynamic mechanical and thermal characteristics of Luffa, Morinda tinctoria and Myrobalan reinforced epoxy hybrid biocomposites. *Fibers Polym*. 2023;24:2093-105. <https://doi.org/10.1007/s12221-023-00196-7>
 48. Bansal A, Chhabra V, Rawal RK, Sharma S. Chemometrics: a new scenario in herbal drug standardization. *J Pharm Anal*. 2014;4(4):223-33. <https://doi.org/10.1016/j.jpha.2013.12.001>
 49. El-Gindy A, Hadad GM. Chemometrics in pharmaceutical analysis: an introduction, review and future perspectives. *J AOAC Int*. 2012;95(3):609-23. https://doi.org/10.5740/jaoacint.SGE_El-Gindy
 50. Vyas A, Jain V, Sahu U, Kumar N, Joshi N. HPTLC method development of herbal drugs and its validation: an overview. *Res J Pharm Technol*. 2023;16(8):3964-6. <https://doi.org/10.52711/0974-360X.2023.00652>
 51. Dharmamoorthy G, Saiteja K, Keerthana J, Dinesh L, Ganesh A, Balaji A. An overview on high-performance thin-layer chromatography (HPTLC). *Asian J Pharm*. 2023;17(1):5-11. <https://doi.org/10.22377/ajp.v17i1>
 52. Pei W, Huang Y, Qu Y, Cui X, Zhou L, Yang H, et al. A strategy for quality evaluation of complex herbal preparations based on multi-color scale and efficacy-oriented high-performance thin-layer chromatography characteristic fingerprint combined with chemometric method: Sanwujiu Pills as an example. *Heliyon*. 2023;9(11):e22098. <https://doi.org/10.1016/j.heliyon.2023.e22098>
 53. Balkrishna A, Sharma P, Joshi M, Srivastava J, Varshney A. Development and validation of a rapid high-performance thin-layer chromatographic method for quantification of gallic acid, cinnamic acid, piperine, eugenol and glycyrrhizin in Divya-Swasari-Vati, an ayurvedic medicine for respiratory ailments. *J Sep Sci*. 2021;44(16):3146-57. <https://doi.org/10.1002/jssc.202100096>
 54. Patel MN, Nandpal MN, Patel AJ, Raval MA, Patel SG. Development and validation of a green stability-indicating HPTLC method for estimation of curcumin, gallic acid and ursolic acid from polyherbal formulation Jatyadi Taila. *J AOAC Int*. 2023;106(4):979-91. <https://doi.org/10.1093/jaoacint/qsac147>
 55. Kaur P, Gupta RC, Dey A, Malik T, Pandey DK. Validation and quantification of major biomarkers in 'Mahasudarshan Churna'-an Ayurvedic polyherbal formulation through high-performance thin-layer chromatography. *BMC Complement Med Ther*. 2020;20:184. <https://doi.org/10.1186/s12906-020-02970-z>
 56. Patel LJ, Raval MA, Patel SG, Patel AJ. Development and validation of stability indicating high-performance thin-layer chromatographic (HPTLC) method for quantification of asiaticoside from *Centella asiatica* L. and its marketed formulation. *J AOAC Int*. 2019;102(4):1014-20. <https://doi.org/10.5740/jaoacint.18-0381>
 57. Govindarajan R, Vyas T, Pushpangadan P. High-performance liquid chromatography (HPLC) as a tool for standardization of complex herbal drugs. *J AOAC Int*. 2019;102(4):986-92. <https://doi.org/10.5740/jaoacint.18-0378>
 58. Xu Y, Wang Y, Yue X, Yang H, Lv P, Bao H. Application of a new validated HPLC-DAD method for simultaneous determination of ten active components in Xiedu San. *Pak J Pharm Sci*. 2021;34(6):2235-45
 59. Lopes AJF, Tsochatzis ED, Emons H, Hoekstra E. Development and validation of an HPLC method with fluorescence detection for the determination of fluorescent whitening agents migrating from plastic beverage cups. *Food Addit Contam Part A*. 2018;35(7):1438-46. <https://doi.org/10.1080/19440049.2018.1459053>
 60. Zheng X, Chen S, Zheng M, Peng J, He X, Han Y, et al. Development of the HPLC-ELSD method for the determination of phytochelators and glutathione in *Perilla frutescens* under cadmium stress conditions. *R Soc Open Sci*. 2018;5(171659). <https://doi.org/10.1098/rsos.171659>
 61. Rodell R, Tsao N, Ganguly A, Mosammamaparast N. Use of High-Performance Liquid Chromatography-Mass Spectrometry (HPLC-MS) to Quantify Modified Nucleosides. In: Mosammamaparast N, editor. DNA Damage Responses. *Methods Mol Biol*. 2022;2444:125-40. https://doi.org/10.1007/978-1-0716-2063-2_8
 62. Langer C, Süß R. HPLC-DAD-CAD-based approach for the simultaneous analysis of hydrophobic drugs and lipid compounds in liposomes and for cyclodextrin/drug inclusion complexes. *J Pharm Biomed Anal*. 2021;201:114120. <https://doi.org/10.1016/j.jpba.2021.114120>
 63. Mishra A, Mishra AK, Tiwari OP, Jha S. HPLC analysis and standardization of Brahmi vati - An Ayurvedic poly-herbal formulation. *J Young Pharm*. 2013;5(3):77-82. <https://doi.org/10.1016/j.jyp.2013.09.001>
 64. Kardani K, Gurav N, Solanki B, Patel P, Patel B. RP-HPLC method development and validation of gallic acid in polyherbal tablet formulation. *J Appl Pharm Sci*. 2013;3(5):37-42
 65. Kamal YT, Singh M, Tamboli ET, Parveen R, Zaidi SMA, Ahmad S. Rapid RP-HPLC method for the quantification of glabridin in crude drug and in polyherbal formulation. *J Chromatogr Sci*. 2012;50(9):779-84. <https://doi.org/10.1093/chromsci/bms063>
 66. Meena AK, Rekha P, Perumal A, Gokul M, Swathi KN, Ilavarasan R. Estimation of Withaferin-A by HPLC and standardization of the Ashwagandhadhi lehyam formulation. *Heliyon*. 2021;7(2):e06116. <https://doi.org/10.1016/j.heliyon.2021.e06116>
 67. Abraham A, Mathew L, Samuel S. HPLC analysis of Pathyashadagam kwath, a classical Ayurvedic polyherbal formulation. *Mater Today Proc*. 2020;25(2):115-21. <https://doi.org/10.1016/j.matpr.2019.12.165>
 68. Want EJ, Metz TO. MS Based Metabonomics. In: Lindon JC, Tranter GE, Koppenaal DW, editors. Encyclopedia of Spectroscopy and Spectrometry. 3rd ed. Academic Press; 2017. p. 926-35. <https://doi.org/10.1016/B978-0-12-803224-4.00030-3>
 69. Martínez Bueno MJ, Agüera A, Gómez MJ, Hernando MD, García-Reyes JF, Fernández-Alba AR. Application of liquid

- chromatography/quadrupole-linear ion trap mass spectrometry and time-of-flight mass spectrometry to the determination of pharmaceuticals and related contaminants in wastewater. *Anal Chem.* 2007;79(24):9372-84. <https://doi.org/10.1021/ac0715672>
70. Zhou JL, Qi LW, Li P. Herbal medicine analysis by liquid chromatography/time-of-flight mass spectrometry. *J Chromatogr A.* 2009;1216(44):7582-94. <https://doi.org/10.1016/j.chroma.2009.05.054>
 71. Chen XF, Wu HT, Tan GG, Zhu ZY, Chai YF. Liquid chromatography coupled with time-of-flight and ion trap mass spectrometry for qualitative analysis of herbal medicines. *J Pharm Anal.* 2011;1(4):235-45. <https://doi.org/10.1016/j.jpha.2011.09.008>
 72. Aslam L, Kaur R, Hussain S, Kapoor N, Mahajan R. LC-MS/MS identification and structural characterization of isolated cyclotides from precursor sequences of *Viola odorata* L. petiole tissue using computational approach. *J Biosci.* 2022;47:50
 73. Lippens JL, Egea PF, Spahr C, Vaish A, Keener JE, Marty MT, et al. Rapid LC-MS method for accurate molecular weight determination of membrane and hydrophobic proteins. *Anal Chem.* 2018;90(22):13616-23. <https://doi.org/10.1021/acs.analchem.8b03843>
 74. Plumb RS, Johnson KA, Rainville P, Smith BW, Wilson ID, Castro-Perez JM, et al. UPLC/MS E; a new approach for generating molecular fragment information for biomarker structure elucidation. *Rapid Commun Mass Spectrom.* 2006;20(13):1989-94. <https://doi.org/10.1002/rcm.2550>
 75. Ammann AA, Macikova P, Groh KJ, Schirmer K, Suter MJF. LC-MS/MS determination of potential endocrine disruptors of cortico signalling in rivers and wastewaters. *Anal Bioanal Chem.* 2014;406:7653-65. <https://doi.org/10.1007/s00216-014-8206-9>
 76. Thakkar AP, Vora A, Kaur G, Akhtar J. An LC-MS/MS method for the quantification of major biomarkers in "Majoon-e-Nisyan"-an Unani polyherbal formulation. *J Anal Chem.* 2023;78:1549-56. <https://doi.org/10.1134/S1061934823110102>
 77. Balkrishna A, Ranjan R, Sakat SS, Sharma VK, Shukla R, Joshi K, et al. Evaluation of polyherbal ayurvedic formulation 'Peedantak Vati' for anti-inflammatory and analgesic properties. *J Ethnopharmacol.* 2019;235:361-74. <https://doi.org/10.1016/j.jep.2019.01.028>
 78. Almalki A, Zaher A, Simithy J, Keller W, Tripp M, Calderón A. LC-MS-based quality assessment of a traditional Chinese medicine YANG XIN formulation. *Planta Med.* 2016;82(13):1208-16. <https://doi.org/10.1055/s-0042-106170>
 79. Fu S, Du L, Yuan Y, He X. Quality evaluation of *Andrographis paniculata* capsules based on rapid and accurate LC-ESI-MS/MS assay of three diterpenoids. *J Chromatogr Sci.* 2019;57(8):708-14. <https://doi.org/10.1093/chromsci/bmz044>
 80. Seo CS, Shin HK. Quality assessment of traditional herbal formula, Hyeonggaeyeongyo-tang through simultaneous determination of twenty marker components by HPLC-PDA and LC-MS/MS. *Saudi Pharm J.* 2020;28(4):427-39. <https://doi.org/10.1016/j.jsps.2020.02.003>
 81. Kännaste A, Copolovici L, Niinemets Ü. Gas chromatography-mass spectrometry method for determination of biogenic volatile organic compounds emitted by plants. In: Rodríguez-Concepción M, editor. *Plant Isoprenoids. Methods Mol Biol.* 2014;1153:161-9. https://doi.org/10.1007/978-1-4939-0606-2_11
 82. Creative Proteomics [Internet]. Gas Chromatography VS Liquid Chromatography [cited 2024 Dec 11]. Available from: <https://www.creative-proteomics.com/resource/gas-chromatography-vs-liquid-chromatography.htm>
 83. Coskun O. Separation techniques: Chromatography. *North Clin Istanbul.* 2016;3(2):156-60. <https://doi.org/10.14744/nci.2016.32757>
 84. Grayson MA. A history of gas chromatography mass spectrometry (GC/MS). In: Gross ML, Caprioli RM, editors. *The encyclopedia of mass spectrometry.* Amsterdam: Elsevier; 2016. p. 152-8. <https://doi.org/10.1016/B978-0-08-043848-1.00020-1>
 85. Olivia NU, Goodness UC, Obinna OM. Phytochemical profiling and GC-MS analysis of aqueous methanol fraction of *Hibiscus asper* leaves. *Futur J Pharm Sci.* 2021;7:59. <https://doi.org/10.1186/s43094-021-00208-4>
 86. George UU, Mbong EO, Bolarinwa KA, Abiaobo NO. Ethnobotanical verification and phytochemical profile of ethanolic leaves extract of two medicinal plants (*Phragmenthera capitata* and *Lantana camara*) used in Nigeria using GC-MS technique. *Acta Biol Forum.* 2023;2(3):1-7
 87. Feng H, Li S, Hu Y, Zeng X, Qiu P, Li Y, et al. Quality assessment of *Succus Bambusae* oral liquids based on gas chromatography/mass spectrometry fingerprints and chemometrics. *Rapid Commun Mass Spectrom.* 2021;35(24):e9200. <https://doi.org/10.1002/rcm.9200>
 88. Devasvaran K, Baharom NH, Chong HW, Ramli RN, Chiu HI, Lee CK, et al. Quality assessment of *Clinacanthus nutans* leaf extracts by GC-MS-based metabolomics. *Curr Sci.* 2020;119(4):641-8. <http://doi.org/10.18520/cs/v119/i4/641-648>
 89. Kotteswari M, Rao MRK, Kumar S, Prabhu K, Sundaram RL, Dinakar S. GC-MS analysis of one Ayurvedic preparation 'Aśwagandharishtam.' *Biomed Pharmacol J.* 2018;11(2):1061-72. <https://doi.org/10.13005/bpj/1467>
 90. Zahiruddin S, Parveen A, Khan W, Ibrahim M, Akhtar J, Khan AA, et al. Quality control and stability testing of Arq formulations of Unani Pharmacopeia of India using HPTLC and GC-MS. *J AOAC Int.* 2020;103(3):699-704. <https://doi.org/10.5740/jaoacint.19-0230>
 91. Ahmad W, Parveen R, Mujeeb M, Zaidi SMA. Comparative fingerprint profiling of Unani polyherbomineral (Safoof-e-Pathar Phori) formulation by HPTLC, HPLC and GC-MS. *J AOAC Int.* 2020;103(3):659-68. <https://doi.org/10.5740/jaoacint.19-0286>
 92. Chen M, Wen SS, Wang R, Ren QX, Guo CW, Li P, et al. Advanced development of supercritical fluid chromatography in herbal medicine analysis. *Molecules.* 2022;27(13):4159. <https://doi.org/10.3390/molecules27134159>
 93. Yuan H, Olesik SV, West C. Supercritical fluid chromatography. In: *Encyclopedia of Analytical Chemistry.* Chichester: John Wiley & Sons, Ltd; 2019. p. 1-19. <https://doi.org/10.1002/9780470027318.a5917.pub2>
 94. Liu TT, Cheong LZ, Man QQ, Zheng X, Zhang J, Song S. Simultaneous profiling of vitamin D metabolites in serum by supercritical fluid chromatography-tandem mass spectrometry (SFC-MS/MS). *J Chromatogr B.* 2019;1120:16-23. <https://doi.org/10.1016/j.jchromb.2019.04.050>
 95. Langeder J, Grienke U. A supercritical fluid workflow for the quality assessment of herbal drugs and commercial preparations from *Rhodiola rosea*. *Phytochem Anal.* 2021;32(6):982-91. <https://doi.org/10.1002/pca.3040>
 96. Nie L, Dai Z, Ma S. Stereospecific assay of (R)- and (S)-Goitrin in commercial formulation of *Radix Isatidis* by reversed phase high-performance liquid chromatography. *J Anal Methods Chem.* 2017;2017:2810565. <https://doi.org/10.1155/2017/2810565>
 97. Ciftci ON, Cahyadi J, Guigard SE, Saldaña MDA. Optimization of artemisinin extraction from *Artemisia annua* L. with supercritical carbon dioxide + ethanol using response surface methodology. *Electrophoresis.* 2018;39(15):1926-33. <https://doi.org/10.1002/elps.201800084>
 98. Mazarakioti EC, Zotos A, Thomatou AA, Kontogeorgos A, Patakas A, Ladavos A. Inductively coupled plasma-mass spectrometry (ICP-MS), a useful tool in authenticity of agricultural products' and foods' origin. *Foods.* 2022;11(22):3705. <https://doi.org/10.3390/foods11223705>
 99. Sharp BL. Pneumatic nebulisers and spray chambers for inductively coupled plasma spectrometry. A review. Part 1. Nebulisers. *J Anal At Spectrom.* 1988;3(5):613-52. <https://doi.org/10.1039/JA9880300613>

100. Pupyshev AA, Semenova EV. Formation of doubly charged atomic ions in the inductively coupled plasma. *Spectrochim Acta B At Spectrosc.* 2001;56(12):2397-418. [https://doi.org/10.1016/S0584-8547\(01\)00301-9](https://doi.org/10.1016/S0584-8547(01)00301-9)
101. Agilent. An introduction to the principles of Inductively Coupled Plasma - Mass Spectrometry (ICP-MS) [Internet]. Santa Clara (CA): Agilent Technologies, Inc.; 2024 [cited 2024 Dec 11].
102. Luo L, Wang B, Jiang J, Fitzgerald M, Huang Q, Yu Z, et al. Heavy metal contaminations in herbal medicines: determination, comprehensive risk assessments and solutions. *Front Pharmacol.* 2021;11:595335. <https://doi.org/10.3389/fphar.2020.595335>
103. Nie LX, Zha YF, Zuo TT, Jin HY, Yu JD, Dai Z, et al. [Determination and risk assessment of heavy metals and harmful elements residues in Niu Huang Qingwei Pills based on ICP-MS]. *Zhongguo Zhong Yao Za Zhi.* 2019;44(1):82-7. <https://doi.org/10.19540/j.cnki.cjcmm.20181108.002>
104. Yao JJ, Kong DD, Luo JY, Qin WJ, Qin XM, Fan ZW, et al. [Safety evaluation of heavy metals contaminated Astragalus membranaceus using health risk assessment model]. *Zhongguo Zhong Yao Za Zhi.* 2019;44(14):3094-9. <https://doi.org/10.19540/j.cnki.cjcmm.20190517.201>
105. Abdalla AA, Smith RE. Determination of mercury in Ayurvedic dietary supplements that are not Rasa Shastra using the Hydra-C direct mercury analyzer. *Int J Anal Chem.* 2013;2013:1-4. <https://doi.org/10.1155/2013/628397>
106. Fu K, Song Y, Zhang D, Xu M, Wu R, Xiong X, et al. Determination of 18 trace elements in 10 batches of the Tibetan medicine Qishiwei Zhenzhu Pills by direct inductively coupled plasma-mass spectrometry. *Evid Based Complement Alternat Med.* 2022;2022:8548378. <https://doi.org/10.1155/2022/8548378>
107. Kumar R, Samajdar S, Mazumder R, Chandra A. Formulation, standardization and evaluation of Vaiśvānaracūrṇa tablet. *Ancient Sci Life.* 2018;38(2):59-67. https://doi.org/10.4103/asl.asl_97_22

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work.

Reprints & permissions information is available at https://horizonpublishing.com/journals/index.php/PST/open_access_policy

Publisher's Note: Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Indexing: Plant Science Today, published by Horizon e-Publishing Group, is covered by Scopus, Web of Science, BIOSIS Previews, Clarivate Analytics, NAAS, UGC Care, etc. See https://horizonpublishing.com/journals/index.php/PST/indexing_abstracting

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

Publisher information: Plant Science Today is published by HORIZON e-Publishing Group with support from Empirion Publishers Private Limited, Thiruvananthapuram, India.