



REVIEW ARTICLE

Insect as food: search for new protein dietary sources – A review

Ismat Parvin¹, Arpita Gope¹, Jayeeta Khanra¹, Anjali Rawani^{1*}

¹Laboratory of Parasitology, Vector Biology, Nanotechnology, Department of Zoology, The University of Gour Banga, Malda, West Bengal, India, PIN-732103

*Email: micanjali@gmail.com

OPEN ACCESS

ARTICLE HISTORY

Received: 07 November 2023
Accepted: 01 January 2023

Available online
Version 1.0 : 20 January 2024
Version 2.0 : 22 February 2024

Additional information

Peer review: Publisher thanks Sectional Editor and the other anonymous reviewers for their contribution to the peer review of this work

Reprints & permissions information is available at https://horizonepublishing.com/journals/index.php/TCB/open_access_policy

Publisher's Note: The article processing was done by atom e-Publishing, Thiruvananthapuram, India. Horizon e-Publishing Group remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Copyright: © The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited (<https://creativecommons.org/licenses/by/4.0/>)

CITE THIS ARTICLE

Parvin I, Gope A, Khanra J, Rawani A. Insect as food: search for new protein dietary sources – A review. Trends in Current Biology. 2024;2(1): 15-25.
<https://doi.org/10.14719/tcb.3073>

Abstract

The increasing global need for food has sparked a great deal of interest in the consumption of novel and diversified sources of protein. Dietary protein not only plays a critical function in the prevention of Protein Energy Malnutrition (PEM) but also in how the human diet affects the environment. Eating insects has been practiced for a very long time. Edible insects have clear economic and environmental benefits, and recently, their huge potential in numerous industries has brought them to the forefront. Due to cultural prejudices towards hazardous insects, edible insects in Western nations are more likely to be used as animal feed than as human food, even though their abundance should be advantageous to human health. However, insects are a major source of protein for many countries in Asia, Africa, and Latin America. Insects may be used to address issues with the conservative food supply chain, such as the scarcity of land, water, and energy resources. Our analysis shows that, despite the fact that consumers affected by Western eating habits are unfamiliar with it, entomophagy is steadily growing throughout the world. Numerous studies have been done to increase their value, however questions about the security of the food supply have been raised. In this review of literature study, comparison have been made with ingestible insects with those of other conventional protein sources. The development of contemporary entomophagy was discussed along with their historical perspective. The practical methods and tactics have been also suggested to encourage the consumption of edible insects.

Keywords

edible insects, entomophagy, nutritional value, food processing, food security

Introduction

Humans have historically consumed the eggs, larvae, pupae, and adults of insects, and this practice continues today in some orthodox countries. In the beginning, man was an omnivore who ate a lot of insects. Prior to the invention of tools for farming or hunting, insects were a major component of human diets. The majority of humanity also resided in warm climates with a year-round diversity of insects. When there was no meat available from vertebrates, insects were often a welcome source of protein (1). One hundred and thirteen nations throughout the world still practice eating human insects. There are around 2000 bug species that can be consumed. Beetles, caterpillars, bees, wasps, and ants are the species that are eaten most frequently on a global scale. They are followed by termites, dragonflies, cicadas, crickets, flies, leafhoppers, grasshoppers, locusts, and other species. In Africa, Asia, and Latin America, people eat insects the most (2). Human ingestion of

insects are exceedingly uncommon and frequently socially unacceptable or even taboo in the majority of European nations. Insects have nutritional value that is comparable to that of regular meats. Consideration should be given to using edible insects as an animal protein source given the expanding human population and the rising demand for traditional meats like beef, hog, and chicken(3).The following insect species could be raised and consumed in Europe under the right farming conditions: the house cricket (*Acheta domestica*), the Jamaican field cricket (*Gryllus assimilis*), the yellow meal-worm beetle (*Tenebrio molitor*), the superworm (*Zophobas morio*), the lesser mealworm (*Alphito biusdiaperinus*), the wax moth (*Galleria mellonella*) and the African migratory locust (*Locust amigratoria*) (4). In comparison to mammalian cattle, insects exhibit better feed efficiency index, it quantifies how well the animal converts feed bulk into body mass. Van Huis et al. (5) even claimed that the feed conversion of the house cricket (*A. domestica*) was more than 12 times higher than that of cattle, four times higher than that of pigs, and double that of chickens. A thorough overview of the usage of insects as food for humans may be found in Figure 1 below.

Nutritional value of insects

Palatable insects have a wide range of nutritional value because of their abundance and variety of species. Even within a same species, an insect's nutritional value might differ greatly depending on its diet, source, and stage of metamorphosis (9). Payne et al. (10) found that the nutritional content of insects differed significantly between species. On the nutrient value scale, mealworms, crickets, and palm weevil larvae all scored higher than beef and poultry, and none of the six insects under study was statistically less healthy than meat. They may also include vitamins such as folic acid, pantothenic acid, biotin, and riboflavin under specific conditions (11).

Energy value

Energy content is typically higher in larvae or pupae than in adults. In contrast, insect species with high protein content have lower energy levels (12). In 78 different insect species, Ramos- Elorduy et al. (13) assessed the calorific value, which ranged from 293 to 762 kcal per 100 g of dry mass. The energy content of a few species of edible insects, represented in kcal per 100 g of fresh weight, is shown in Table 1(5).

Table 1: Energy value of edible insect (5)

Order	Common name	Scientific name	Phase	Area	En. value (kcal/100g)
Orthoptera					
	Australian plague locust	<i>Chortoicetes foraminifera</i>	Adult	Australia	499
	wo-spotted cricket	<i>Gryllus bimaculatus</i>	Adult	Thailand	120
	Japanese grasshopper	<i>Oxyajaponica</i>	Adult	Thailand	149
	Brown-spotted locust	<i>Cyrtacanthacris tatarica</i>	Adult	Thailand	89
Lepidoptera	African migratory locust	<i>Locusta migratoria</i>	Adult	Netherlands	179
Hymenoptera	Silkworm	<i>Bombyx mori</i>	Pupa	Thailand	94
	Weaver ant	<i>Oecophylla maragdina</i>	Adult	Australia	1272
Coleoptera	Mexican leaf cutting ant	<i>Atta Mexicana</i>	Adult	Mexico	404
	Yellow mealworm beetle	<i>Tenebrio molitor</i>	Larvae	USA	206




Fig1. Flowchart of value of insects as human food (6-8)

Proteins

Seven kinds of insects were studied for their total protein content (12). All varieties of insects that were measured had roughly the same amounts of total protein, with the exception of the protein content of the wax moth (*G. mellonella*), which was just 38.4% (13). Yellow mealworms (*T. molitor*) made up 50.7% of the other species, while African migrating locusts (*L. migratoria*) made up 62.2%. 100 insect species' protein levels were examined by Xiaoming et al (14). Table 2 shows that, by dry matter, the protein concentration varied from 13 to 77%. Furthermore, edible insects have some essential amino acids, including threonine, lysine, and tryptophan which can be insufficient in certain plant or grain proteins (15, 16). Table 3 shows the insects from different order and their nutritional content, such as protein, fat, thiamine, and riboflavin. Overall, the edible insects may be a helpful complement to diets, especially in areas where other foods may lack certain nutrients. (5,17).

Lipids

Between 10% and 60% of the dry mass in edible insects is made up of fat (Table 4). In comparison to adults, this is higher in the larval stages (14). According to study by Tzompa-Sosa et al. (20) per 100 g of Lepidopteran caterpillars contains about 8.6 to 15.2 g of the total fat content, placing them among the insects with the greatest fat content. In comparison, grasshoppers and similar orthopteran species have a fat content from 3.8 g to 5.3 g per 100 g of insects. In the insect, fat can be found in a variety of forms. The second-most significant category is phospholipids. Phospholipids make up less than 20% of fat in most cases, but this varies depending on the phases of life cycle and species of insect (20,21). The cholesterol is the highly predominant sterol in insects. The cholesterol level of the fat from two regularly eaten animals in Nigeria: the caterpillar *Imbrasia belina* and the termite *Macrotermes bellicosus* was investigated by Ekpo et al (21) and found that the lipid fraction had an average cholesterol level of 3.6%. Besides cholesterol, palatable insects may also provide other sterols such as campesterol, stigmasterol, and sitosterol (22).

Table 2: Protein content of some insect species (12)

Insect name	Order	Life Stages	Protein content (% in dry matter)
Beetles	<i>Coleoptera</i>	Larvae and Adults	23–66
Butterflies	<i>Lepidoptera</i>	Larvae and Pupae	14–68
Hemipterans	<i>Hemiptera</i>	Larvae and Adults	42–74
Homopterans	<i>Homoptera</i>	Eggs, larvae and Adults,	45–57
Hymenopterans	<i>Hymenoptera</i>	eggs, larvae, Adult and pupae	13–77
Dragonflies	<i>Odonata</i>	naiads and Adults	46–65
Orthopterans	<i>Orthoptera</i>	nymphs and Adults	23–65

Table 3: Some pertinent insects' nutritional qualities in relation to other high-protein diets (18, 19)

Order	Insect's common name	Scientific name	Protein (g/kg)	Fat (g/kg)	Calories (kcal/kg)	Thiamine (mg/kg)	Riboflavin (mg/kg)
Diptera	Blacksoldier	<i>Hermetia illucens</i>	175	140	1994	7.7	16.2
	Fly(larvae), Housefly(adults)	<i>Musca domestica</i>	197	19	918	11.3	77.2
	Housecricket (adults)						
Orthoptera	Superworm (larvae)	<i>Acheta domestica</i>	205	68	1402	0.4	34.1
		<i>Zophobas morio</i>	107	177	2423	0.6	7.5
		<i>Tenebrio molitor</i>	187	134	2056	2.4	8.1
	Mealworm(larvae), Mealworm(adult), Giant mealworm (larvae),	<i>Tenebrio molitor</i>	237	54	1378	1	8.5
Coleoptera	Waxworm(larvae) Silkworm(larvae)	<i>Tenebrio molitor</i>	184	168	2252	1.2	16.1
	Silkworm(Moth)	<i>Galleria mellonella</i>	141	249	2747	2.3	7.3
		<i>Bombyx mori</i>	93	144	674	3.3	9.4

Table 4: Percentage of fat in dry material of edible insects (12)

Order	Common name	Scientific name	Life Stage	Fat content (% in dry matter)
Coleoptera	Silkworm	<i>Bombyx mori</i>	Pupa	29
	Yellow mealworm	<i>Tenebrio molitor</i>	Larva	36
	Giant mealworm	<i>Zophobas atratus</i>	Larva	40
Hymenoptera	Western honey bee	<i>Apis mellifera</i>	Brood	31
Lepidoptera	Wax moth	<i>Galleria mellonella</i>	Caterpillar	57
Orthoptera	African migratory locust	<i>Locusta migratoria</i>	Nymph	13
	Jamaican field cricket	<i>Gryllus assimilis</i>	Nymph	34

Fibre

A substantial amount of fibre can be found in edible insects. The most prevalent type of fibre in an insect's body is insoluble chitin, which is primarily found in the exoskeleton. In commercially farmed insects 2.7 to 49.8 mg of chitin were found in per kilogram of fresh weight (i.e. equivalent to 11.6 to 137.2 mg per kg of dry matter) (5, 23). Chitin is still considered an indigestible fibre even though human stomach contents contain the enzyme chitinase (24). Additionally, insects can be a source of fibre and contain variable levels of fat, including essential fatty acids. Chitin, a component of insect exoskeletons, may also have immune-boosting and anti-allergenic properties (25, 26, 27). It implies that more investigation into the dietary and health advantages of edible insects may result in higher consumption and broader acceptance of insects as beneficial food sources (Table 5).

Minerals

When it comes to the nutritional value of minerals like calcium, phosphorus, magnesium, manganese, and copper, as well as iron, zinc, potassium, sodium, and potassium, edible insects can be an intriguing option (5). For instance, 8-20 mg per 100 g of dry matter of the grasshopper *L. migratoria* and 31-77 mg per 100 g of dry matter of the big mopane larva of the moth *Gonimbrasia belina* are combined showed high in iron content (28). When 26.5 mg per 100 g of dry matter of palm weevil larvae *Rhynchophorus phoenicis* combined with 14 mg per 100 g of dry matter of mopane caterpillars found to be an effective source of zinc (7). However, Hyun et al. (29) found that an edible grasshopper, *Oxya chinensis formosana*, have the heavy metal concentration which was minimal and safe for human consumption. It had plumbum at 0.01–0.03 mg/100

g, cadmium at 0.002–0.005 mg/100 g, arsenic at 0.07–0.17 mg/100 g, and mercury at 0.0003–0.0007 mg/100 g. All this concentration of heavy metal in these insects found to be safe of human consumption.

Vitamins

Numerous water-soluble or lipophilic vitamins are found in insects (17, 30, 31). Bukkens (17) provided a list of several insects that contain thiamine. Its concentration was 0.1 to 4 mg per 100 g of dry matter while each 100 g of dry matter of some insects contains riboflavin retinol and 6.8–8.2 g of beta-carotene. Some levels of retinol that were less than 20 g per 100 g of dry matter and -carotene that were less than 10 g per 100 g of dry matter were found in yellow mealworms *T. molitor*, superworms *Z. morio*, and house crickets *A. domesticus* (data shown in figure 2) (17, 30). Vitamins A, D, and E may be found in eggs and escamoles from the Formicidae family. They included 2.22 mg of each 100 g of alpha-tocopherol, 3.31 mg per 100 g of cholecalciferol, and 505 g per 100 g of retinol. In general, insects are high in riboflavin, pantothenic acid, and biotin, according to Rumpold and Schlüter (11). Nevertheless, these are not a dependable source of thiamin, vitamin A, vitamin C or niacin. Oonocx and Dierenfeld (31) also noted the low vitamin E contents (6–16 mg/kg DM) in the majority of the examined insect species, except *Drosophila melanogaster* and *Microcentrum rhombifolium* they have 112 and 110 mg per kg DM respectively. All specimens had low levels of retinol, a marker of vitamin A activity, but these levels varied widely between samples of 0.670–886 mg/kg DM. It is important to note that the vitamin and mineral content of wild edible insects' changes with the season and, for farm-bred species, can be controlled by feed.

Table 5: In the dry matter of edible insect's content of neutral-detergent fibre (12)

Order	common name	Scientific name	Life Stages	Fibre content
Hymenoptera	Honey bee (western)	<i>Apis mellifera</i>	Brood	11
Orthoptera	migratory locust (African)	<i>Locusta migratoria</i>	Nymph	27
	Jamaican field cricket	<i>Gryllus assimilis</i>	Nymph	8
Coleoptera	Yellow mealworm	<i>Tenebrio molitor</i>	Larva	18
	Giant mealworm	<i>Zophobas atratus</i>	Larva	17
	Silkworm	<i>Bombyx mori</i>	Pupa	14
	Wax moth	<i>Galleria mellonella</i>	Caterpillar	21




Figure 2. Nutritional evaluation of standard diets and yellow mealworm larvae. values for protein, lipids, and ashes based on dry weight. Source: Based on data from Alves et al. (32).

Insect food and its acceptance

There are several ways to prepare and eat insects, including

Whole insect: Certain insects, including crickets, are edible intact after they've been cooked. Mealworm larvae can also be eaten whole after they have been cooked (33, 34).

Dehydrated and crushed: To add protein or other nutrients to a variety of food products, such as frankfurters, insects can also be dried and crushed (Figure 3) (35).

Added fresh, without drying, to food products after killing: Mealworms can be used to make processed foods like burgers by adding them whole to beef after they have been killed but before drying (34).

As extracted nutrients: Insects nutrients can be extracted and incorporated into diet. For instance, in order to increase the protein composition of food products, Zhao et al. (36) extracted and characterised mealworm protein.



Figure 3. The freeze-dried whole yellow mealworm larvae (A) and the milled and freeze-dried larvae (B) (Photo by Xue Zhao, SLU, Sweden)

Although mealworm larvae have high protein content, but neophobia and cultural conditioning make it difficult for people to accept them as food (37). To prevent phobias that can skew the results, sensory testing on foods made from insects might need to be carried out on people from cultures where eating insects is a tradition (37,38). According to some research, adding entire mealworm larvae powder to meals may be acceptable, and flavouring mealworm-based goods may increase their attractiveness (33,34,39). However, the kind of food matrix and the

quantity of mealworm added are important factors influencing consumer approval, as adding too much mealworm to a meal in place of meat can increase off-flavour and reduce juiciness (34).

Historical perspective of taking insects as food

Ancient tribes probably consumed insects often since they had well-developed hunting skills and few advanced tools (40). However, insect-eating customs have been disappearing in many places since the introduction of farming and the domestication of animals (41). Due to evolving societal attitudes, insects have evolved from being a primary food source to snacks, gourmet delicacies, and bait (42). Nevertheless, several nations continue to this day to use insects as stable food sources (43).

- In Asia, there are some nations, like China, where there are no explicit rules for the regulation of insects, where they have long been utilised as food and feed. Canned silkworm pupae and Silkworm (*Bombyx mori*) consumed in Korea (44,45,46,47,48). Fried grasshoppers are used in the Japanese cuisine (45). In China, Silkworm (*Bombyx mori*), Tussar silkworm (*Anthera eapernyi*), Mealworm (*Tenebrio molitor*) etc. being devoured (45,46,49,50,51). In India, the pupae of the Eri silkworm (*Samiaricini*) are prized (47). In Thailand, peoples consumed Silkworm (*Bombyx mori*), Ground cricket (*Allonemobius fasciatus*), Weaver ant (*Oecophyllas maragdina*) etc. (45,48, 49, 50,51).
- The African food culture places a significant emphasis on edible insects (52). Insects from order Lepidoptera, Orthoptera, and Coleoptera are also frequently consumed (45,53). *Macrotermes spp.* (Termite), *Ruspolia nitidula* (Grasshopper), *Chaoborus edulis* (Lake fly), *Ruspolia differens* (Grasshopper) are consumed by Uganda people (54). Silk moth, termites are eaten by Congo people and in Kenya black ant (*Carebara vidua*), Grasshopper (*Ruspolia differens*), Winged termite (*Macrotermes subhylanus*) insects are also consumed (45,51,52).
- In Papua New Guinea, a common edible insect is the sago grub (*Rhynchophorus ferrugineus*) and is the centrepiece of an annual grub festival (55). According

to Meyer- Rochow and Changkija (56), mole-crickets, mantises, crickets, locusts, and even spiders are eaten locally. According to Macfarlane (57), indigenous cultures ingested a diverse range of insects from the Cossidae, Noctuidae, Cerambycidae, and bees (58, 55,56, 57). Here insects are now offered on restaurant menus (59).

- In the Americas, Women and children actively go out hunting for edible insects, which are an important source of protein for Amazonian tribes (60). Palm weevil (*Rhynchophorus palmarum*), Ant (*Liometopuma piculatum*), Stink bug (*Atiziestax coensis*) are consumed in Mexico and consumed type are larvae, adult, pupae (45,61,62). Beetles (*Caryobruchus scheelaeae*), Wasps (*Poly baignobilis*), Grasshopper (*Aidemona Azteca*) are consumed in Colombia (61,62). The Federal Food and Drug Administration (FDA) in the USA is in charge of policing the safety of animal feed in North America (63) where a list of edible insects is classified as food additives.
- A concern to consumer health and safety is posed by Bovine Spongiform Encephalopathy (BSE), which has an impact on European regulations regarding the use of insects as feed (64, 65). Yellow mealworm, lesser mealworm, house cricket, banded cricket, field cricket, black army fly, and common housefly are among the insects that can be used (66, 67, 68). Many insect species like *Apis mellifera*, *Bombyx mori*, *Acheta domesticus*, *Locust amigratoria*, *Tenebrio molitor*, *Galleria mellonella*, *Gryllus assimillis*, *Hermetia illucens* consumed in Europe (69).

Food processing of edible insects

Millions of people worldwide eat edible insects, which are high in protein, vitamins, and minerals (70). Food processing of edible insects can enhance their palatability, digestibility, and safety. Here are some common food processing methods used for edible insects:

1.Cleaning: Before processing, insects need to be cleaned to remove any dirt, debris, or unwanted material.

2.Roasting: Roasting is a common processing method that improves the flavour of insects. It involves heating the insects at high temperatures, which also helps to kill any bacteria.

3.Drying: Drying is a good way to save insects for future use. It entails drying out insects by placing them under an extended period of low heat.

4.Grinding: Using insects as an ingredient in different culinary products can be accomplished by grinding insects into a fine powder. A protein-rich item that can be utilised in baked products, snacks, and other foods is insect flour.

5.Fermentation: Enhancing the flavour and digestibility of insects through fermentation. In order to break down the insects' proteins and other nutrients, this technique includes adding healthy bacteria or yeast to the insects.

6.Extraction: By crushing or grinding the insects, it is possible to acquire insect oils and extracts. These can be incorporated into a variety of food items, such as salad dressings, sauces, and marinades.

7.Freezing: Freezing insects is a common method for preserving them for later use. Soups, stews, and stir-fries are just a few of the recipes that can benefit from using frozen insects (70). Insects are increasingly seen as the meal of the future as conventional methods of consuming animal protein are questioned (71). According to studies aiming at adapting insects for Western tastes, some insects (such as locusts and mealworms) are now sold at retail markets in the Netherlands as part of efforts to encourage entomophagy (72). To reduce visual associations and improve palatability, insects have been treated into powder or meal (73). Table 6 summarises previous studies on the useful properties of edible insects for food processing. In addition, appropriate extraction techniques, such as zymolysis, sonication, defatting, pH adjustments are being studied by Mishyna et al.;

Table 6: Studies conducted in the past on edible insects' ability to metabolise food

Order	Scientific name	Stages of insects	Observation	Reference (78)
Lepidoptera	<i>Cirina forda</i> , <i>Bombyx mori</i> L.	Larvae, Larvae	Protein solubility Oil absorption capacity Frothing stability Water absorption capacity Emulsion capability	(79)
Orthoptera	<i>Grylloides sigillatus</i> <i>Acheta domesticus</i> <i>Schistocerca gregaria</i>	Larvae, Cricket, Grasshopper	Protein solubility Water holding ability Oil holding capacity Frothing capacity Emulsion capacity Water binding capacity	[80]
Coleoptera	<i>Tenebrio molitor</i> <i>Oryctes owariensis</i> <i>Zophobas morio</i> <i>Alphitobius diaperinus</i>	Cricket, Larvae, Larvae, Larvae	Oil immersion capacity Water absorption capacity Frothing capacity Frothing stability Frothing stability	(81)
Diptera	<i>Hermetia illucens</i>	Larvae	Emulsion capacity Emulsion stability Fat binding capacity Protein solubility Frothing stability	(73)
Blattodea	<i>Blaptica dubia</i>	Cockroach	Gel formation	(82)

Nongonierma and FitzGerald; Purschke et al., and Yi et al. (74,75, 76,77). These investigations will support the usage of insects as food components in conventional dishes.

Product made from insects

In cultures where eating insects intact is common and considered a culinary tradition, the practice dates back

centuries. Eating them as snacks, stir-frying them, grilling them over coals, or adding them to stews or soups. They are occasionally ground, used as flavour, and occasionally processed into powder and blended with salts and spices (83,84). Table 7 represents the Product made from insects, their nutritional component and types of insects used.

Table 7: List of products made from insects

Product made from insects	Nutritional component	Product named	Insects used	References
Insect Protein bars	Protein	Chapul	Several bugs Cricket	
Bread	Protein Minerals Vitamin B12	Crunchy Cricket Loaf	Ground crickets Mealworms	
Burgers	Iron Calcium Manganese	Bug Burger	Bug	
Candy & chocolate covered insects	Protein Carbohydrate	Ant wafers, worm suckers, cricket and mealworm chocolate, insect marshmallows, mealworm lollipops,	Grasshopper Ant Mealworm Black soldier flies Silkworm pupae Scorpions Super worms Sago worms Japanese Cricket Ground insect Crickets Grasshoppers Silkworm pupae	
Coffee	Vitamin Magnesium Potassium Phenolic compounds	Japanese Cricket Blend coffee	Cricket	
Cookie	Carbohydrate Protein Fibre	Bitty's cookies	Bugs	
Crisps/chips	Carbohydrate Fat	American Chirps	Cricket	
Crisps bread	Vitamin B12 Omega-2 Iron	Swedish Eat.em.	Mealworm Cricket	
Ice cream	Protein	"ento milk"	black soldier fly larvae Cricket	
Meat replacement (replacing minced meat)	Protein	Sirkkis	Cricket	
Noodles	Fat Carbohydrate Dietary fibre Sodium Protein Minerals Pre-biotic fibre	Bugsfarmsudo noodles Nutribugs cricket rice noodles	Bug Cricket	(83)
Pancake mix	Protein Iron Vitamin B12 Content	ZIRP pancake mix	Cricket	
Pasta	Potassium Carbohydrate	Hoppa Foods Eat Crawlers Circle Harvest Jimini's Nutribug	Cricket Mealworms Silkworm pupae	
Pasta sauce	Protein Fat and fibre Minerals	Gryllies Tomato Sauce	Cricket bolognese	
Pâté / bread spread / butters	Protein	Crickster	Bugs	
Powder (sometimes called Flour)	Protein	Bug flour Cricket flour	Acheta domesticus Cricket Black Soldier Fly larvae	
Powder / baking flour	Calcium Vitamin B12 Fattyacids Iron Zinc	Cricket flour	Cricket Bug	
Shakes and protein powder	Vitamin B12 Protein	Cricket protein powder	Crickets Mealworms Bugs	
Protein Snacks	Protein	Insect meal	Cricket	
Roasted whole insect (with different flavours)	Protein	Chilli lime crickets	Grasshoppers Mealworms Crickets	
Sausages	Protein Fat	Sirkkanakki," or "cricket sausage	Cricket	
Shakes and smoothies	Protein	Sirkka smoothie	Crickets Mealworms	
Spices & seasoning	Protein Vitamin Essential amino acid Omega - 3	Short-Horn super seasonings	Ant Cricket	
Soft drinks	Carbohydrate Sugar	Japanese Tagame Cider	Giant Water bug	

Insect protein vs meat protein

Even more so than mammals, insects can serve as a source of protein. According to a research in the European Journal of Clinical Nutrition, insect such as crickets have a protein content of 9.96 to 35.2 grammes per 100 grammes, compared to 16.8 to 20.6 grammes in meat. However, depending on the kind of bugs fed, protein density does vary significantly. Options abound with approximately 2,100 different edible bug species available. Crickets, various ant species, and mealworms are the catalyst of insect protein movement mostly because of their high calorie and protein density (85).

Food security

Food security can benefit from edible insects, especially in areas with limited or difficult-to-access traditional food sources. Insects are a more sustainable and effective source of food since they can be raised with a fraction of the land, water, and feed resources needed for raising conventional livestock (86,87). Due to their high adaptability and ability to be raised in a range of habitats, including urban ones, insects may be able to help solve difficulties with urban food deserts and food insecurity (88). Additionally, there may be concerns related to the safety and quality of insect-based food products, particularly if they are not produced and processed under appropriate conditions. The hazards of bacterial infection cannot be completely eliminated by boiling (89, 90). During the production and storage phases, precautions must be taken. Overall, a variety of factors, such as the development of suitable production and processing methods, the availability of suitable feed sources, and the implementation of appropriate regulatory structures to guarantee the security and excellence of food items derived from insects, will determine the food security potential of edible insects (91, 92).

Consumer acceptance of edible insect

Consumer acceptance of edible insects varies depending on region and culture, with insects being a traditional food source in many parts of South America, Asia, and Africa, but not widely accepted in North America and Europe (93, 94, 95). Factors such as cultural norms, taste preferences, and perceived health and environmental benefits can influence acceptance. Efforts to increase consumer acceptance include education campaigns to promote the nutritional and sustainability benefits of insect consumption (96, 97).

Future direction

A growing global population's nutritional needs may be met by edible insects in a different and sustainable way (98, 99). Although it is unknown how much agony and pain insects experience, they appear to pose little difficulties for animal welfare. Multidisciplinary research on edible insects is being conducted, for instance, at the Pan-African International Centre of Insect Physiology and Ecology (ICIPE) in Nairobi, Kenya. A scientific research organisation, ICIPE's primary goals are to improve the health and food security of those who live in tropical areas. ICIPE creates and implements innovative approaches and

technologies for managing insects that are practical, efficient, non-harmful, economical for resource-strapped communities, and acceptable from a cultural and social standpoint (100). In order to assure the necessary investments and corporate structures, comprehensive legal frameworks for the production and trade of insect products are needed on a national and worldwide basis.

Conclusion

In conclusion, edible insects may supply significant quantities of proteins, lipids, vitamins, and minerals with significant economic and environmental benefits. Entomophagy holds the answer to meeting the world's expanding nutrient needs. Insects can be used in a variety of besides being consumed as foods and snacks. The thorough study of insects has led to the development of numerous contemporary items. Eating edible insects is becoming more and more common in different countries worldwide. In addition to eating insects for food, people do so for entertainment. Still, there are worries about the potential security and health hazards associated with employing edible insects. Currently, the market for edible insects is not in line with the advantages they may offer. To entice and reassure, production and promotion strategies are suggested. It is advised to use semi-cultivation because it can successfully increase the output of some insects. Both insect and insect products should be of high quality, hence farming and processing practices need to be standardised. For effective collaboration and increased earnings, communications between farmers and industries are encouraged. The creation of new insect products, advancements in cultivation, and production optimisation are anticipated to lead to agricultural industrial integration.

Acknowledgements

We acknowledge with thanks to all the members of Laboratory of Parasitology, Vector Biology, Nanotechnology for their technical and general support.

Author's contributions

I.P. and A.G. wrote the main manuscript text and J.K. prepared tables and figures. A.R. supervised the whole work. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest: Authors do not have any conflict of interests to declare.

Ethical issues: None.

References

1. Sponheimer M, Ruiters D, Lee-Thorp J, Spath A. Sr/Ca and early hominin diets revisited: new data from modern and fossil tooth enamel. *J Hum Evol.* 2005; 48(2): 147-56. <https://doi.org/10.1016/j.jhevol.2004.09.003>.

2. Jongema Y. World List of Edible Insects. Laboratory of Entomology, Wageningen University, Wageningen. 2015. <http://www.wageningenur.nl/en/Expertise-Services/Chair-groups/Plant-Sciences/Laboratory-of-Entomology/Edible-insects/Worldwide-species-list.htm>.
3. Dreon AL, Paoletti MG. The wild food (plants and insects) in Western Friuli local knowledge (Friuli-Venezia Giulia, North Eastern Italy), *Contrib. Nat. Hist.* 2009; 12: 461–488.
4. Hardy A, Benford D, Notebom H, Halldorsson T, Schlatter J, Solecki R, Jeger M, Knutsen H. Risk profile related to production and consumption of insects as food and feed, *EFSA J.* 2015; 13 (10). <https://doi.org/10.2903/j.efsa.2015.4257>.
5. Van HA, Itterbeeck JV, Klunder H, Mertens, Halloran E, Muir G, Vantomme P. *Edible Insects. Future Prospects for Food and Feed Security*, FAO, Rome. 2013. (201 pp.)
6. Daylan A, Sosa T, Gent U, Fogliano V. Potential of insect-derived ingredients for food applications. In *Insect Physiology and Ecology*, Intechopen: London, UK. 2017: 215–231.
7. Kourimska L, Adamkova A. Review article: Nutritional and sensory quality of edible insects. *NFS J.* 2016; 4: 22–26. 4. [10.1016/j.nfs.2016.07.001](https://doi.org/10.1016/j.nfs.2016.07.001).
8. Van HA. Potential of insects as food and feed in assuring food security. *Annu. Rev. Entomol.* 2013; 58: 563–583. <https://doi.org/10.1146/annurev-ento-120811-153704>.
9. Finke M, Oonincx DG. Insects as food for insectivores. *Mass Production of Beneficial Organisms : Invertebrates and Entomopathogens*, Elsevier, New York. 2014: 583-616. doi: 10.1016/B978-0-12-391453-8.00017-0.
10. Payne C, Scarborough P, Rayner M, Nonaka K. Are edible insects more or less 'healthy' than commonly consumed meats? A comparison using two nutrient profiling models developed to combat over- and undernutrition. *Eur J Clin Nutr.* 2016; 70(3): 285-91. Doi: 10.1038/ejcn.2015.149.
11. Rumpold BA, Schluter OK. Nutritional composition and safety aspects of edible insects. *Mol Nutr Food Res.* 2013; 57(5): 802-23. Doi: 10.1002/mnfr.201200735.
12. Bednarova M. Possibilities of Using Insects as Food in the Czech Republic Dissertation thesis Mendel University, Brno. 2013: 50–92.
13. Ramos EJ, Pino JM, Prado EE, Perez MA, Otero JL, De GL. Nutritional value of edible insects from the state of Oaxaca, Mexico, *J. Food Compos. Anal.* 1997; 10: 142–157.
14. Xiaoming C, Ying F, Hong Z. Review of the nutritive value of edible insects. *Edible insects and other invertebrates in Australia: future prospects*, Proceedings of a Workshop on Asia-Pacific Resources and their Potential for Development. 2008: 85–92.
15. Finke MD. Nutrient content of insects, in: J.L. Capinera (Ed.), *Encyclopedia of Entomology*, Kluwer Academic, Dordrecht, London. 2004: 1562–1575.
16. Sogbesan A, Ugwumba A. Nutritional evaluation of termite (*Macrotermes subhyalinus*) meal as animal protein supplements in the diets of *Heterobranchus longifilis*, *Turk. J. Fish. Aquat. Sci.* 2008; 8: 149–157.
17. Bukkens S. Insects in the Human Diet: Nutritional Aspects, in: M.G. Paoletti (Ed.), *Ecological Implications of Mini live stock; Role of Rodents, Frogs, Snails, and Insects for Sustainable Development*. Science Publishers, New Hampshire 2005: 545–577.
18. Finke MD. Complete nutrient composition of commercially raised invertebrates used as food for insectivores. *Zoo Biol.* 2002; 21: 269–285. doi: 10.1002/zoo.10031.
19. Shockley M, Dossey AT. Insects for Human Consumption. In: Morales-Ramos, J.A., Guadalupe Rojas, M. and Shapiro-Ilan, D.I., Eds., *Mass Production of Beneficial Organisms: Invertebrates and Entomopathogens*, Academic Press, Cambridge, MA. 2014: 617-652. <https://doi.org/10.1016/B978-0-12-391453-8.00018-2>.
20. Tzompa SD, Yi L, Valenberg H, Boekel M, Lakemond C. Insect lipid profile: aqueous versus organic solvent-based extraction methods. *Food Res. Int.* 2014; 62: 1087–1094. doi:10.1016/j.foodres.2014.05.052.
21. Ekpo KE, Onigbinde AO, Asia IO. Pharmaceutical potentials of the oils of some popular insects consumed in southern Nigeria, *Afr. J. Pharm. Pharmacol.* 2009; 3(2): 51–57. <https://doi.org/10.5897/AJPP.9000216>.
22. Sabolova M, Adamkova A, Kourimska L, Chrpova D, Panek J. Minor lipophilic compounds in edible insects, *Potravinarstvo.* 2016; 10(1). Doi:10.5219/605.
23. Finke MD. Estimate of chitin in raw whole insects, *Zoo Biol.* 2007; 26: 105–115.
24. Paoletti MG, Norberto L, Damini R, Musumeci S. Human gastric juice contains chitinase that can degrade chitin. *Ann. Nutr. Metab.* 2007; 51(3): 244–251. doi: 10.1159/000104144.
25. Muzzarelli R, Terbojevich M, Muzzarelli C, Miliani M, Francescangeli O. Partial depolymerization of chitosan with the aid of papain, in: R.A.A. Muzzarelli (Ed.), *Chitin Enzymology*, Atec, Grottammare. 2001: 405–414.
26. Goodman W.G. Chitin: a magic bullet? *Food Insects Newsletter.* 1989; 2(3): 6–7.
27. Muzzarelli RA. Chitins and chitosans as immunoadjuvants and non-allergenic drug carriers. *Mar. Drugs.* 2010; 8(2): 292–312. Doi: 10.3390/md8020292.
28. Oonincx DG, Itterbeeck J, Heetkamp MJ, Van BH, Loon JJ, Huis A. An exploration on greenhouse gas and ammonia production by insect species suitable for animal or human consumption. *PLoS One.* 2010; 5(12). doi: 10.1371/journal.pone.0014445.
29. Hyun SH, Kwon KH, Park KH, Jeong HC, Kwon O, Tindwa H, Han YS. Evaluation of nutritional status of an edible grasshopper, *Oxyachinensis formosana*. *Entomol. Res.* 2012; 42(5): 284–290. Doi:10.1111/j.1748-5967.2012.00469.x
30. Finke MD. Complete nutrient composition of commercially raised invertebrates used as food for insectivores. *Zoo Biology.* 2002; 21(3): 269–285. doi: 10.1002/zoo.10031.
31. Oonincx DG, Dierenfeld ES. An investigation into the chemical composition of alternative invertebrate prey. *Zoo Biol.* 2012; 31(1): 40–54. doi: 10.1002/zoo.20382.
32. Alves AV, Sanjinez EJ, Linzmeier AM, Cardoso CA, Macedo ML. Food Value of Mealworm Grown on *Acrocomia aculeata* Pulp Flour. *PLoS One.* 2016; 11(3): e0151275. doi: 10.1371/journal.pone.0151275.
33. Sheppard B, Frazer P. Comparing social and intellectual appeals to reduce disgust of eating crickets. *Stud. Arts Humanit.* 2015; 1(2): 4–23. DOI:10.18193/sah.v1i2.29.
34. Megido RC, Gierts C, Blecker C, Brostaux Y, Haubruge E, Alabi T, Francis F. Consumer acceptance of insect-based alternative meat products in Western countries. *Food Qual. Prefer.* 2016; 52: 237–243. Doi: 10.1016/j.foodqual.2016.05.004.
35. Choi YS, Kim TK, Choi HD, Park JD, Sung JM, Jeon KH, Paik HD, Kim YB. Optimization of Replacing Pork Meat with Yellow Worm (*Tenebrio molitor* L.) for Frankfurters. *Korean J Food Sci Anim Resour.* 2017; 37(5): 617–625. doi: 10.5851/kosfa.2017.37.5.617.
36. Zhao X, Vazquez GJ, Johansson D, Landberg R, Langton M. Yellow mealworm protein for food purposes. Extraction and functional properties. *PLoS ONE.* 2016; 11(2), e0147791. <https://doi.org/10.1371/journal.pone.0147791>.
37. Hamerman EJ. Cooking and disgust sensitivity influence preference for attending insect-based food events. *Appetite.*

- 2016;96: 319-326. Doi: 10.1016/j.appet.2015.09.029.
38. Barbera F, Vermeau F, Amato M, Grunert K. Understanding Westerners' disgust for the eating of insects: The role of food neophobia and implicit associations. *Food Qual. Prefer.* 2018; 64: 120–125. Doi:10.1016/j.foodqual.2017.10.002.
 39. Aguilar ED, Lopez MG, Escamilla SC, Barba AP. Characteristics of maize flour tortilla supplemented with ground *Tenebriomolitor* larvae. *J Agric Food Chem.* 2002; 50(1):192-5. Doi: 10.1021/jf010691y.
 40. Valadez AR. La domesticación animal UNAM, Instituto de Investigaciones Antropológicas, Plaza y Valdés Editores, México City. 2003.
 41. Abrams HL. The preference for animal protein and fat: A cross-cultural survey. Temple University Press; Philadelphia, PA, USA. 1987.
 42. Smeathman H. Some account of the termites, which are found in Africa and other hot climates. In a letter from Mr. Henry Smeathman, of Clement's Inn, to Sir Joseph Banks, Bart. P. R. S. *Philos Trans R Soc Lond.* 1781; 71: 17810033. doi: 10.1098/rstl.1781.0033.
 43. Lahteenmaki A, Grmelova N, Henault L, Deschamps MH, Vandenberg GW, Zhao A, Zhang Y, Yang B, Neman V. Laws of the European Union, United States, Canada, Mexico, Australia, and China. *Eur. Food Feed Law Rev. EFFL* 2017; 12(1): 22–36.
 44. Pemberton RW. The revival of rice-field grasshoppers as human food in South Korea. *Pan-Pac Entomol.* 1994; 70: 323–327.
 45. DeFoliart GR. Insects as food: why the western attitude is important. *Annu Rev Entomol.* 1999; 44: 21-50. Doi: 10.1146/annurev.ento.44.1.21. PMID: 9990715.
 46. Feng Y, Chen XM, Zhao M, He Z, Sun L, Wang CY, Ding WF. Edible insects in China: Utilization and prospects. *Insect Sci.* 2018; 25: 184–198. Doi: 10.1111/1744-7917.12449.
 47. Peigler RS. Wild silks of the world. *Am Entomol.* 1993; 39: 151–162. Doi:10.1093/ae/39.3.151.
 48. Sungpuag P, Puwastien P. Nutritive value of unconventional protein source: Insect. *Pochanagan Sarn.* 1983; 1: 5–12.
 49. Yhoun-aree J. Edible insects in Thailand: Nutritional values and health concerns. In: Durst PB, Johnson DV, Leslie RN, Shono K, editors. *In Forest insects as food: Humans bite back* Food and Agriculture Organization of the United Nations; Bangkok, Thailand. 2010: 201–216.
 50. Chakravorty J. Diversity of edible insects and practices of entomophagy in India: An overview. *J Biodivers Biopros Dev.* 2014; 1(3): 1-6. doi: 10.4172/2376-0214.1000124.
 51. Raheem D, Carrascosa C, Oluwole OB, Nieuwland M, Saraiva A, Millan R, Raposo A. Traditional consumption of and rearing edible insects in Africa, Asia and Europe. *Crit Rev Food Sci Nutr.* 2018; 59: 2169–2188. Doi: 10.1080/10408398.2018.1440191.
 52. Mutungi C, Irungu FG, Nduko J, Mutua F, Affognon H, Nakimbugwe D, Ekesi S, Fiaboe K. Postharvest processes of edible insects in Africa: A review of processing methods, and the implications for nutrition, safety and new products development. *Crit Rev Food Sci Nutr.* 2019; 59: 276–298. Doi: 10.1080/10408398.2017.1365330.
 53. Kelemu S, Niassy S, Torto B, Fiaboe K, Affognon H, Tonnang H, Maniania NK, Ekesi S. African edible insects for food and feed: Inventory, diversity, commonalities and contribution to food security. *J Insect Food Feed.* 2015; 1: 103–119. Doi: 10.3920/JIFF2014.0016.
 54. Mbabazi M. Dietary contribution of grasshoppers (*Ruspolianitidula*) and white ants (*Macrotermes bellicosus*) and influence of processing methods on their nutrient composition. Makerere Univ, Kampala, Uganda. 2011. <http://hdl.handle.net/10570/3546>.
 55. Mercer CWL. Insects as food in Papua New Guinea. In *Proceedings of Invertebrates Farming Seminar.* Institute of Tropical Medicine; Antwerp, Belgium. 1993: 157–162.
 56. Meyer VB, Changkija S. Uses of insects as human food in Papua New Guinea, Australia, and North East India: Cross cultural considerations and cautious conclusions. *Ecol Food Nutr.* 1997; 36(2-4): 159–185. Doi: 10.1080/03670244.1997.9991513.
 57. Macfarlane WV. Aboriginal desert hunter-gatherers in transition. *Symposium on the Nutrition of Aborigines, Commonwealth Scientific and Industrial Research Organisation; Melbourne.* 1978: 49–62.
 58. Yen AL. Edible insects and other invertebrates in Australia: Future prospects. In: Durst PB, Johnson DV, Leslie RN, Shono K, editors. *In Forest insects as food: Humans bite back.* FAO; Bangkok, Thailand. 2010: 65–84.
 59. Irvine G. Putting insects on the Australian menu. *Food Aust.* 1989; 41(1): 565–566.
 60. Dufour DL. Insects as food: A case study from the northwest Amazon. *American Anthropol.* 1987; 89: 383–397. Doi: 10.1525/aa.1987.89.2.02a00070.
 61. Ramos BJ. The importance of edible insects in the nutrition and economy of people of the rural areas of Mexico. *Ecol Food Nutr.* 1997; 36: 347–366. Doi: 10.1080/03670244.1997.9991524.
 62. Ruddle K. The human use of insects: Examples from the Yukpa. *Biotropica.* 1973; 5(2): 94–101. Doi: 10.2307/2989658.
 63. Lahteenmaki A, Henault L, Marimuthu SB, Talibov S, Allen RN, Neman V, Vandenberg GW, Jozefiak D. The impact of the insect regulatory system on the insect marketing system. *J. Insects Food Feed.* 2018; 4(3): 187–198. <https://doi.org/10.3920/JIFF2017.0073>.
 64. Marberg A, van Kranenburg H, Korzilius H. The big bug: The legitimization of the edible insect sector in the Netherlands. *Food Policy.* 2017; 71(C): 111–123. Doi: 10.1016/j.foodpol.2017.07.008.
 65. Belluco S, Halloran A, Ricci A. New protein sources and food legislation: the case of edible insects and EU law. *Food Sec.* 2017; 9: 803–814. <https://doi.org/10.1007/s12571-017-0704-0>.
 66. EC Council. Regulation (EC) No 999/2001 of the European Parliament and of the Council of 22 May 2001 laying down rules for the prevention, control and eradication of certain transmissible spongiform encephalopathies. *Off. J. Eur. Union.* 2001; L147: 1–40.
 67. Regulation (EU) No 2017/893 of 24 May 2017 Amending Annexes I and IV to Regulation (EC) No 999/2001 of the European Parliament and of the Council and Annexes X, XIV and XV to Commission Regulation (EU) No 142/2011 as Regards the Provisions on Processed Animal Protein. Available online: <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32017R0893> (accessed on 12 February 2019).
 68. Regulation (EU) No 2017/1017 of 15 June 2017 Amending Regulation (EU) No 68/2013 on the Catalogue of Feed Materials. Available online: <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32017R1017> (accessed on 12 February 2019).
 69. Comby B. *Delicieux insectes. Les proteines du futur*, 1st ed., Editions Jouvence: Geneve, Switzerland. 1990: 105–111.
 70. Kim TK, Cha JY, Yong HI, Jang HW, Jung S, Choi YS. Application of Edible Insects as Novel Protein Sources and Strategies for Improving Their Processing. *Food Sci Anim Resour.* 2022; 42(3): 372–388. Doi: 10.5851/kosfa.2022.e10.
 71. Sogari G. Entomophagy and Italian consumers: An exploratory analysis. *Prog Nutr.* 2015; 17: 311–316.

72. Schosler H, Boer J, Boersema JJ. Can we cut out the meat of the dish? Constructing consumer-oriented pathways towards meat substitution. *Appetite*. 2012; 58(1): 39-47. Doi: 10.1016/j.appet.2011.09.009.
73. Bubler S, Rumpold BA, Jander E, Rawel HM, Schluter OK. Recovery and techno-functionality of flours and proteins from two edible insect species: Meal worm (*Tenebrio molitor*) and black soldier fly (*Hermetia illucens*) larvae. *Heliyon*. 2016; 2(12): e00218. Doi: 10.1016/j.heliyon.2016.e00218.
74. Mishyna M, Martinez JI, Chen J, Benjamin O. Extraction, characterization and functional properties of soluble proteins from edible grasshopper (*Schistocerca gregaria*) and honey bee (*Apis mellifera*). *Food Res Int*. 2019; 116: 697-706. Doi: 10.1016/j.foodres.2018.08.098.
75. Nongonierma AB, Fitzgerald RJ. Unlocking the biological potential of proteins from edible insects through enzymatic hydrolysis: A review. *Innov Food Sci Emerg Technol*. 2017; 43: 239-252. Doi: 10.1016/j.ifset.2017.08.014.
76. Purschke B, Meinschmidt P, Hom C, Rieder O, Jager H. Improvement of techno-functional properties of edible insect protein from migratory locust by enzymatic hydrolysis. *Eur Food Res Technol*. 2018; 244: 999-1013. Doi: 10.1007/s00217-017-3017-9.
77. Yi L, Boekel JS, Boeren S, Lakemond MM. Protein identification and *in vitro* digestion of fractions from *Tenebrio molitor*. *Eur Food Res Technol*. 2016; 242: 1285-1297. Doi: 10.1007/s00217-015-2632-6.
78. Omotoso OT. Nutritional quality, functional properties and anti-nutrient compositions of the larva of *Cirina fforda* (Westwood) (Lepidoptera: Saturniidae). *J Zhejiang Univ Sci B*. 2006; 7(1): 51-55. Doi: 10.1631/jzus.2006.B0051.
79. Zielinska E, Karas M, Baraniak B. Comparison of functional properties of edible insects and protein preparations thereof. *LWT-Food Sci Technol*. 2018; 91(4): 168-174. Doi: 10.1016/j.lwt.2018.01.058.
80. Osasona AI, Olaofe O. Nutritional and functional properties of *Cirina fforda* larva from Ado-Ekiti, Nigeria. *Afr J Food Sci*. 2010; 4: 775-777.
81. Assielou B, Due EA, Koffi MD, Dabonne S, Kouame PL. *Oryctesowariensis* larvae as good alternative protein source: Nutritional and functional properties. *Annu Res Rev Biol*. 2015; 8: 1-9. Doi: 10.9734/ARRB/2015/19093.
82. Yi L, Lakemond MM, Sagis MC, Eisner V, Huis A, Boekel JS. Extraction and characterisation of protein fractions from five insect species. *Food Chem*. 2013 Dec; 141(4): 3341-3348. Doi: 10.1016/j.foodchem.2013.05.115.
83. Food Products Made from Insects – Insect Gourmet – Your Guide to Edible Insects. www.insectgourmet.com
84. Tang C, Yang D, Liao H. Edible insects as a food source: a review. *Food Prod Process and Nutr* 1, 2019 Nov; 8. <https://doi.org/10.1186/s43014-019-0008-1>.
85. Payne CL, Scarborough P, Rayner M, Nonaka K. Are edible insects more or less 'healthy' than commonly consumed meats? A comparison using two nutrient profiling models developed to combat over- and under nutrition. *Eur J Clin Nutr*. 2016 Mar; 70(3): 285-91. Doi: 10.1038/ejcn.2015.149.
86. Van Huis A, Van Itterbeeck J, Klunder H, Mertens E, Halloran A, Muir G. *Edible insects: future prospects for food and feed security*. Rome: Food and Agriculture Organization of the United Nations. 2013. 187 p. (FAO forestry paper; 171).
87. Premalatha M, Abbasi T, Abbasi T. Energy-efficient food production to reduce global warming and ecodegradation: the use of edible insects. *Renew. Sust. Energ. Rev*. 2011, 15(9): 4357-4360. Doi: 10.1016/j.rser.2011.07.115.
88. Lloyd SJ, Kovats RS, Chalabi Z. Climate change, crop yields, and undernutrition: development of a model to quantify the impact of climate scenarios on child undernutrition. *Environ Health Perspect*. 2011 Dec; 119(12): 1817-23. Doi: 10.1289/ehp.1003311.
89. Ramos-Elorduy J, Moreno MP, Prado EE, Perez MA, Otero JL, Guevara OL. Nutritional value of edible insects from the state of Oaxaca, Mexico. *Journal of Food Composition and Analysis*. 1997; 10(2): 142-157. Doi: 10.1006/jfca.1997.0530.
90. Payne CLR, Van Itterbeeck J. Ecosystem services from edible insects in agricultural systems: A review. *Insects*. 2017 Feb; 8(1). <https://doi.org/10.3390/insects8010024>.
91. Klunder HC, Wolkers-Rooijackers J, Korpela JM, Nout MJR. Microbiological aspects of processing and storage of edible insects. *Food Control*. 2012 Aug; 26(2): 628-631. <https://doi.org/10.1016/j.foodcont.2012.02.013>.
92. Choo J. Potential ecological implications of human entomophagy by subsistence groups of the Neotropics. *Terrestrial Arthropod Reviews*. 2008 June; 1(1): 81-93. Doi: 10.1163/187498308X345442.
93. Rozin P, Fallon AE. A perspective on disgust. *Psychol. Rev.*, 1987, 94(1): 23-41. Doi: 10.1037/0033-295X.94.1.23.
94. Hartmann C, Siegrist M. Consumer perception and behaviour regarding sustainable protein consumption: A systematic review. *Trends in Food Science & Technology*. 2017 Mar; 61: 11-25. Doi: 10.1016/j.tifs.2016.12.006.
95. Vane-Wright RI. Why not eat insects? *Bull. Entomol. Res*. 1991 Feb; 81(01): 1-4. Doi: 10.1017/S0007485300053165.
96. Mignon J. L'entomophagie: une question de culture? *Tropicicultura*. 2002 Jan; 20(3): 151-155.
97. Mela DJ. Food choice and intake: the human factor. *Proc Nutr Soc*. 1999 Aug; 58(3): 513-21. Doi: 10.1017/s0029665199000683.
98. Sun-Waterhouse D, Waterhouse GIN, You L, Zhang J, Liu Y, Ma L, Gao J, Dong Y. Transforming insect biomass into consumer wellness foods: A review. *Food Res Int*. 2016 Nov; 89(Pt 1): 129-151. Doi: 10.1016/j.foodres.2016.10.001.
99. Mlcek J, Rop O, Borkovcova M, Bednarova M. A Comprehensive Look at the Possibilities of Edible Insects as Food in Europe – A Review. *Polish Journal of Food and Nutrition Sciences*. 2014 Sep; 64(3): 147-157. Doi: 10.2478/v10222-012-0099-8.
100. International Centre of Insect Physiology and Ecology. <http://www.icipe.org>. 2021.